

(19) World Intellectual Property  
Organization  
International Bureau



539,956  
10/539956



(43) International Publication Date  
15 July 2004 (15.07.2004)

PCT

(10) International Publication Number  
**WO 2004/058181 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K** (74) Agent: **WILSON, Mark, B.**; Fulbright & Jaworski L.L.P., Suite 2400, 600 Congress Avenue, Austin, TX 78701 (US).
- (21) International Application Number: PCT/US2003/041182 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 22 December 2003 (22.12.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/435,077 20 December 2002 (20.12.2002) US (84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: US 60/435,077 (CON) Filed on 20 December 2002 (20.12.2002)
- (71) Applicant (*for all designated States except US*): **BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th Street, Austin, TX 78701 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **NORRIS, Steven, J.** [US/US]; 6043 Beaudry Dr., Houston, TX 77035 (US).
- Published:**  
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: VMP-LIKE SEQUENCES OF PATHOGENIC *BORRELIA* SPECIES AND STRAINS

(57) Abstract: The present invention relates to DNA sequences encoding Vmp-like polypeptides of pathogenic *Borrelia*, the use of the DNA sequences in recombinant vectors to express polypeptides, the encoded amino acid sequences, application of the DNA and amino acid sequences to the production of polypeptides as antigens for immunoprophylaxis, immunotherapy, and immunodiagnosis. Also disclosed are the use of the nucleic acid sequences as probes or primers for the detection of organisms causing Lyme disease, relapsing fever, or related disorders, and kits designed to facilitate methods of using the described polypeptides, DNA segments and antibodies.

**BEST AVAILABLE COPY**

WO 2004/058181 A2

## DESCRIPTION

### VMP-LIKE SEQUENCES OF PATHOGENIC *BORRELLIA* SPECIES AND STRAINS

#### BACKGROUND OF THE INVENTION

5           The government owns rights in the present invention pursuant to grant number AI37277 from the National Institutes of Health .

##### A.     Field of the Invention

10           The invention relates to the field of molecular biology; in particular, to immunogenic compositions and recombinant VMP-like genes useful for treatment and diagnosis of Lyme disease. Also included are methods for the determination of virulence factors in Lyme disease.

##### B.     Description of Related Art

15           Lyme disease is a bacterial infection caused by pathogenic spirochetes of the genus *Borrelia*. The infection can occur in humans, dogs, deer, mice and other animals, and is transmitted by arthropod vectors, most notably ticks of the genus *Ixodes*. *Borrelia burgdorferi*, the most common cause of Lyme disease in North America, was first cultured in 1982. *B. garinii*  
20           and *B. afzelii* are the most common infectious agents of Lyme disease in Europe, and another species, *B. japonicum*, has been described in Japan. These organisms are closely related and cause similar manifestations with multiple stages: an expanding rash at the site of the tick bite (erythema migrans); fever, lymphadenopathy, fatigue, and malaise; effects of disseminated infection, including carditis, meningoradiculitis, and polyarthrititis; and chronic manifestations  
25           including arthritis and neurologic disorders.

          Lyme disease is often difficult to diagnose because of shared manifestations with other disorders, and it can also be refractory to treatment during late stages of the disease. It is most common in areas such as suburban regions of upstate New York and Connecticut, where large populations of deer and white-footed mice serve as the principal mammalian hosts and reservoirs  
30           of infection. Approximately 20,000 cases of Lyme disease in humans are reported per year in the United States, and it is also a significant veterinary problem due to a high infection rate of dogs and other domestic animals in endemic regions.

          The pathogenic *Borrelia* that cause Lyme disease are able to persist for years in patients or animals despite the presence of an active immune response. Antigenic variation is a

mechanism by which members of the genus *Borrelia* may be able to evade the host immune response (Zhang, 1997; Wang 2003). Antigenic variation has been defined as changes in the structure or expression of antigenic proteins that occurs during infection at a frequency greater than the usual mutation rate (Borst and Geaves, 1987; Robertson and Meyer, 1992; Seifert and So, 1988).

Relapsing fever is another disease caused by pathogenic *Borrelia*. It has both epidemic and endemic forms. The epidemic form is caused by *B. recurrentis* and is transmitted between humans by lice. It was a major source of morbidity and mortality during World War I, but has been rare since then due largely to public health measures. Endemic relapsing fever is an epizootic infection caused by several *Borrelia* species, including *B. hermsii*. It occurs sporadically among hunters, spelunkers, and others who come in contact with infected soft-bodied ticks of the genus *Ornithodoros*. Relapsing fever is characterized by two or more episodes or "relapses" of high bacteremia (up to  $10^8$ /ml). The first wave of infection is caused by *Borreliae* expressing a certain Variable Major Protein (VMP) on their surface (e.g. Vmp21). The gene encoding this VMP is located at a promoter site in the expression plasmid, whereas over 24 nonexpressed copies of different VMP genes are present on the so-called silent plasmid. When the host develops antibodies against the expressed VMP, the organisms of that serotype are destroyed and the patient improves. However, a small proportion of organisms have undergone antigenic switching to a different serotype. Nonreciprocal recombination occurs between the expression plasmid and the silent plasmid, resulting in the insertion of a different VMP gene in the expression site (e.g., Vmp7). The organisms expressing Vmp7 are not affected by the anti-Vmp21 antibodies, and therefore multiply in the host and cause a second episode of the disease. Up to five of these 3-5 day episodes can occur, separated by 1-2 week intervals.

Such well-demarcated episodes of infection do not occur during Lyme disease, and fewer organisms are present in the blood at any stage. However, there are reasons to suspect that similar mechanisms of antigenic variation may occur in *B. afzelii* and other Lyme disease *Borreliae* such as *B. garinii* and *B. burgdorferi*. The infection, if untreated, commonly persists for months to years despite the occurrence of host antibody and cellular responses; this observation indicates effective evasion of the immune response. Lyme disease may be disabling (particularly in its chronic form), and thus there is a need for effective therapeutic and prophylactic treatment.

Genetic loci analogous to the VMP antigenic variation system have been detected in North American and European Lyme disease *Borrelia* by Southern hybridization and PCR

analysis (Iyer *et al.*, 2000; Wang *et al.*, 2001). In addition, sequences from fragments of *vls* (VMP-like sequence) silent cassettes have been reported for the *Borrelia burgdorferi* strains 297 and N40, and the *Borrelia garinii* strains Ip90 and A87S (Kawabata *et al.*, 1998; Liang and Philipp, 1999; Wang *et al.*, 2001), (S. Feng and S. W. Barthold, unpublished data). VMP-like sequences of *B. burgdorferi* have been described and patented in U.S. Patent No. 6,437,116.

Open reading frames in a *B. burgdorferi* plasmid that encode hypothetical proteins resembling the VMP proteins of relapsing fever organisms have been identified (Zhang *et al.*, 1997). The inventors have found that the presence of the plasmid containing these VMP-like sequences in *B. burgdorferi* clones correlates strongly with infectivity (Zhang *et al.*, 1997; Purser and Norris, 2000; Labandeira-Ray *et al.*, 2001). Thus it is likely that the proteins encoded by the VMP-like sequences are important in immunoprotection and pathogenesis. Proteins encoded by the VMP-like sequences of *B. burgdorferi* may provide protection when used either alone or in combination with other antigens. They may also be useful for immunodiagnosis.

Greater than 90% of Lyme disease patients beyond the erythema migrans stage from North America and Europe express antibodies against VlsE (Lawrenz *et al.*, 1999; Liang *et al.*, 1999; Liang *et al.*, 2000). In addition, mice infected experimentally with *Borrelia afzelii* and *Borrelia garinii* strains express anti-VlsE antibodies (Liang *et al.*, 2000). Finally, a protein product of ~35 kDa expressed by *Borrelia garinii* Ip90 reacts with antibodies against IR6, a peptide corresponding to invariant region 6 of the VlsE cassette region (Liang *et al.*, 1999a). Portions of several *vls* silent cassettes from *Borrelia garinii* strain A87S have been published (Wang *et al.*, 2001). Further, several amino acid sequences of *Borrelia garinii* Ip90 have been previously characterized by Liang *et al.* (1999a).

There is a commercial demand for vaccines and diagnostic kits for Lyme disease, both for human and veterinary use. Several companies have active research and development programs in these areas.

## SUMMARY OF THE INVENTION

Partial and complete DNA sequences have been determined for several recombinant clones containing DNA encoding VMP-like sequences. The identification and characterization of these sequences now allows: (1) identification of the expressed gene(s) or DNA segments containing open reading frames in several *Borreliae*; (2) expression of these gene(s) by a recombinant vector in a host organism such as *E. coli*; (3) immunization of laboratory animals



with the resulting polypeptide, and determination of protective activity against *Borreliae* infection; (4) use of antibodies against the expressed protein to identify the reactive polypeptide(s) in *Borreliae* cells; (5) use of the expressed protein(s) to detect antibody responses in infected humans and animals; (6) determination of the presence, sequence differences, and expression of the VMP-like DNA sequences in other Lyme disease *Borreliae*.

The invention is contemplated to be useful in the immunoprophylaxis, diagnosis, or treatment of Lyme disease, relapsing fever, or related diseases in humans or animals. It is expected that recombinant or native proteins expressed by the VMP-like genes (or portions thereof) will be useful for (a) immunoprophylaxis against Lyme disease, relapsing fever, or related disorders in humans and animals; (b) immunotherapy of existing Lyme disease, relapsing fever, or related illnesses, by way of immunization or injection of antibodies directed against VMP-like proteins; and (c) immunodiagnosis of Lyme disease, relapsing fever, or related diseases, including their use in kits in which the VMP-like proteins are the sole antigen or one of multiple antigens. The DNA may be employed in: (a) production of recombinant DNA plasmids or other vectors capable of expressing recombinant polypeptides; and (b) design and implementation of nucleic acid probes or oligonucleotides for detection and/or amplification of VMP-like sequences. The latter is expected to have application in the diagnosis of infection with *Borrelia* organisms.

Another aspect of the invention is the method for identification of possible virulence factors. This approach entails subtractive hybridization of target DNA from high infectivity organisms with driver DNA from low-infectivity strains or clones. This procedure greatly enriches for sequences which differ between the high- and low-infectivity strains and thus may encode proteins important in virulence. Of particular utility is the use of closely related isogenic clones that differ in their infectivity; in this case, the DNA differences should be restricted more stringently to those related to infectivity.

The invention is considered to include DNA segments corresponding to 10, 20, 30, and 40 base pairs of the VMP-like sequences; DNA segments inclusive of the entire open reading frames of the VMP-like sequences; shorter DNA segments containing portions of these open reading frames; amino acid sequences corresponding to both conserved and variable regions of the VMP-like sequences; recombinant vectors encoding an antigenic protein corresponding to the above amino acid sequences; recombinant cells where extrachromosomal DNA expresses a polypeptide encoded by the DNA encoding *Borrelia* VMP-like sequences; a recombinant *Borreliae* or *E. coli* cell containing the DNA encoding VMP-like sequences; methods of

preparing transformed bacterial host cells using the DNA encoding the VMP-like polypeptides; methods using the plasmid or another vector to transform the bacterial host cell to express *Borreliae* polypeptides encoded by the DNA sequences; and methods for immunization of humans or animals with the native *Borreliae* polypeptides, polypeptides expressed by recombinant cells that include DNA encoding the VMP-like polypeptides, or synthetic peptides that include VMP-like sequences.

Also included in the invention are primer sets capable of priming amplification of the VMP-like DNA sequences; kits for the detection of *Borreliae* nucleic acids in a sample, the kits containing a nucleic acid probe specific for the VMP-like sequences, together with a means for detecting a specific hybridization with the probe; kits for detection of antibodies against the VMP-like sequences of *Borreliae* and kits containing a native, recombinant, or synthetic VMP-like polypeptide, together with means for detecting a specific binding of antibodies to the antigen.

A preferred embodiment of the present invention is an isolated nucleic acid comprising a nucleotide sequence that encodes an antigenic peptide of *Borrelia garinii* or *B. afzelii*. More preferably, the present invention provides an isolated nucleic acid that encodes at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

In another embodiment, the present invention provides an isolated nucleic acid comprising 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300,

400, 500 or more contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96. Further, the invention contemplates any range derivable between any of the above-described integers.

In yet another embodiment, the isolated nucleic acid comprises a complement to or a degenerate variant of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500 or more contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96. Further, the invention contemplates any range derivable between any of the above-described integers.

In some embodiments the isolated nucleic acid is a DNA molecule. In other embodiments the isolated nucleic acid is an RNA molecule.

In certain embodiments the invention provides an isolated nucleic acid obtained by amplification from a template nucleic acid using a primer selected from the group consisting of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101,

SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:107.

The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, enhancers, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like.

A preferred embodiment of the present invention is an isolated polypeptide comprising 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

In one aspect, the present invention provides for an isolated polypeptide or an isolated nucleic acid encoding a polypeptide having between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and 90%; or even more preferably between about 90% and about 99% of amino acids that are identical to the amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or fragments thereof. The percent identity or homology is determined with regard to the length of the relevant amino acid sequence. Therefore, if a polypeptide of the present invention is comprised within a larger polypeptide, the percent homology is determined with regard only to the portion of the polypeptide that

corresponds to the polypeptide of the present invention and not the percent homology of the entirety of the larger polypeptide.

In addition, the present invention encompasses fragments of polypeptides or nucleic acids encoding fragments of polypeptides that have between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and 90%; or even more preferably between about 90% and about 99% of amino acids that are identical to the amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 even if the particular fragment itself does not have between about 70% and about 75%; or more preferably between about 75% and about 80%; or more preferably between about 80% and 90%; or even more preferably between about 90% and about 99% amino acid homology with the polypeptides of the present invention.

In another embodiment the invention provides an isolated polypeptide that binds immunologically with antibodies raised against an antigenic polypeptide of *Borrelia garinii* or *B. afzelii*. In a preferred embodiment the antibodies are raised against an antigenic polypeptide comprising at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 75, 100, 125, 150, 175, 200 or more contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97. Further, the invention contemplates any range derivable between any of the above-described integers.

The polypeptides of the present invention may be fused with other proteins or peptides. Such fusion polypeptides may be useful for purification or immunodetection purposes, for

example. In a preferred embodiment the polypeptides of the invention are expressed as fusions with  $\beta$ -galactosidase, avidin, ubiquitin, *Schistosoma japonicum* glutathione S-transferase, multiple histidines, epitope-tags and the like.

Another aspect of the invention comprises vectors that comprise a nucleic acid encoding all or part of a polypeptide of the present invention. The vectors may, for example, be cloning or expression vectors.

In certain embodiments, it is contemplated that particular advantages will be gained by positioning the nucleic acid sequences of the present invention under the control of a promoter. The promoter may be the promoter that is normally associated with the nucleic acid sequence in its natural environment or it may be a recombinant or heterologous promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with a *vls* gene in its natural environment. Such promoters may include those normally associated with other *Borrelia* polypeptide genes, or promoters isolated from any other bacterial, viral, eukaryotic, or mammalian cell. Naturally, it will be important to employ a promoter that effectively directs the expression of the nucleic acid in the particular cell being used.

The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level or regulated expression of the introduced nucleic acid. In preferred embodiments the promoters are lac, T7, Ara, CMV, RSV LTR, the SV40 promoter alone, or the SV40 promoter in combination with the SV40 enhancer.

Another embodiment is a method of preparing a protein composition comprising growing a recombinant host cell comprising a vector that encodes a polypeptide of the present invention under conditions permitting nucleic acid expression and protein production followed by recovering the protein so produced. The host cell, conditions permitting nucleic acid expression, protein production and recovery, will be known to those of skill in the art, in light of the present disclosure of the *vls* gene. The recombinant host cell may be a prokaryotic cell or a eukaryotic cell.

VMP-like related proteins and functional variants are also considered part of the invention. Thus it is expected that truncated and mutated versions of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ

ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 will afford more convenient and effective forms of polypeptides for treatment regimens. Thus, any functional version of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97, such as truncated species or homologs, and mutated versions of VMP-like protein are considered as part of the invention.

Another aspect of the invention comprises the recombination of the 14 silent *vls* cassettes of *B. afzelii* in numerous combinations, providing for example a cocktail of peptide compositions for use as immunogens to develop vaccines for use in Lyme disease and related conditions. Likewise, the 11 silent *vls* cassettes of *B. garinii* and the 15 silent *vls* cassettes of *B. burgdorferi* may be recombined in numerous combinations. It is further contemplated by the present invention that these cassettes may be recombined among strains, as well as species of *Borrelia*, providing a cocktail of peptide compositions for use as immunogens to develop vaccines for use in Lyme disease and related conditions.

Pharmaceutical compositions prepared in accordance with the present invention find use in preventing or ameliorating conditions associated with *Borrelia* infections, particularly Lyme disease.

Such methods generally involve administering a pharmaceutical composition comprising an effective amount of a VMP-like antigen of *Borrelia*, such as SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85,

SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or various epitopes thereof.

In certain embodiments of the invention a vaccine may comprise a polynucleotide encoding an antigenic polypeptide. In more specific embodiments the polynucleotide may have a  
5 sequence of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64,  
10 SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96 or various fragments thereof. The vaccines of the present invention may comprise multiple polypeptides and/or polynucleotides.

15 It will also be understood that, if desired, the nucleic acid segment or gene encoding a VMP-like protein could be administered in combination with further agents, such as, proteins or polypeptides or various pharmaceutically active agents. There is virtually no limit to other components which may also be included, given that the additional agents do not cause a significant adverse effect upon contact with the target cells or host tissues.

20 Therapeutic kits comprising a polypeptide or nucleic acid of the present invention comprise another aspect of the invention. Such kits will generally contain, in suitable container means, a pharmaceutically acceptable formulation of a polypeptide or nucleic acid of the present invention. The kit may have a single container means that contains a polypeptide or nucleic acid of the present invention or it may have distinct container means for the polypeptide or nucleic  
25 acid of the present invention and other reagents that may be included within such kits.

The components of the kit may be provided as liquid solution(s), or as dried powder(s). When the components are provided in a liquid solution, the liquid solution is an aqueous solution, with a sterile aqueous solution being particularly preferred. When reagents or components are provided as a dry powder, the powder can be reconstituted by the addition of a  
30 suitable solvent. It is envisioned that the solvent may also be provided in another container means.

In another embodiment, the invention provides diagnostic kits. The diagnostic kits may comprise reagents for detecting VMP-like polypeptides or anti-VMP-like antibodies in a sample,



such as required for immunoassay. The immunodetection reagent will typically comprise a label associated with the antibody or antigen, or associated with a secondary binding ligand. Exemplary ligands might include a secondary antibody directed against the first antibody or antigen or a biotin or avidin (or streptavidin) ligand having an associated label. Of course, a number of exemplary labels are known in the art and all such labels may be employed in connection with the present invention. The kits may contain antibody-label conjugates either in fully conjugated form, in the form of intermediates, or as separate moieties to be conjugated by the user of the kit.

The container means will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which the antigen or antibody may be placed, and preferably suitably aliquoted. Where a second binding ligand is provided, the kit will also generally contain a second vial or other container into which this ligand or antibody may be placed. The kits of the present invention will also typically include a means for containing the antibody, antigen, and reagent containers in close confinement for commercial sale. Such containers may include injection or blow-molded plastic containers into which the desired vials are retained.

In another aspect, the present invention contemplates an antibody that is immunoreactive with a polypeptide of the invention. An antibody can be a polyclonal or a monoclonal antibody. In a preferred embodiment, an antibody is a monoclonal antibody.

Antibodies, both polyclonal and monoclonal, specific for VMP-like polypeptides and particularly those represented by SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97 or variants and epitopes thereof, may be prepared using conventional immunization techniques, as will be generally known to those of skill in the art.

In related embodiments, the invention provides methods of using the antibodies of the invention. In preferred embodiments, the antibodies may be used in immunochemical procedures, such as ELISA and Western blot methods. In other embodiments, the antibodies

may be used in purifying native or recombinant VMP-like polypeptides, inhibition studies, and immunolocalization studies.

Table 1 below provides the SEQ ID NO, the GenBank accession number, if any, and a brief description of the sequences described herein.

TABLE 1

SEQ ID NO.	GENBANK NO.	DESCRIPTION
SEQ ID NO:1	U76405	<i>B. burgdorferi</i> vlsE gene allele vlsE1
SEQ ID NO:2	AAC45733	Translation of <i>B. burgdorferi</i> vlsE1 gene
SEQ ID NO:3	L04788	<i>B. hermsii</i> vmp17 gene
SEQ ID NO:4	AAA22963	Translation of <i>B. hermsii</i> vmp17 gene
SEQ ID NO:5	AY100629	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2622 vlsE
SEQ ID NO:6	AAM77200	Translation of AY100629
SEQ ID NO:7	AY100630	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2624a vlsE
SEQ ID NO:8	AAM77201	Translation of AY100630
SEQ ID NO:9	AY100631	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2624b vlsE
SEQ ID NO:10	AAM77202	Translation of AY100631
SEQ ID NO:11	AY100632	RT-PCR product of <i>B. afzelii</i> strain ACAI clone 2625 vlsE
SEQ ID NO:12	AAM77203	Translation of AY100632
SEQ ID NO:13	AY100634	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 17 vlsE
SEQ ID NO:14	AAM77204	Translation of AY100634
SEQ ID NO:15	AY100635	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 20 vlsE
SEQ ID NO:16	AAM77205	Translation of AY100635
SEQ ID NO:17	AY100636	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 21 vlsE
SEQ ID NO:18	AAM77206	Translation of AY100636
SEQ ID NO:19	AY100637	RT-PCR product of <i>B. garinii</i> strain Ip90 clone 23 vlsE
SEQ ID NO:20	AAM77207	Translation of AY100637
SEQ ID NO:21	N/A	Primer 4540 (Wang <i>et al.</i> , 2001)
SEQ ID NO:22	N/A	Primer 4548 (Wang <i>et al.</i> , 2001)
SEQ ID NO:23	N/A	Primer 4545 (Wang <i>et al.</i> , 2001)
SEQ ID NO:24	N/A	Primer 4587 (Wang <i>et al.</i> , 2001)
SEQ ID NO:25	N/A	Primer 4588 (Wang <i>et al.</i> , 2001)
SEQ ID NO:26	N/A	Primer 4470 (Wang <i>et al.</i> , 2001)
SEQ ID NO:27	N/A	Primer 4471 (Wang <i>et al.</i> , 2001)
SEQ ID NO:28	AY100633	<i>B. garinii</i> vls silent cassette locus
SEQ ID NO:29	AY100633	<i>B. garinii</i> upstream ORF
SEQ ID NO:30	AAN87823	Translation of <i>B. garinii</i> upstream ORF
SEQ ID NO:31	AY100633	<i>B. garinii</i> 5' vlsE homolog
SEQ ID NO:32	AAN87824	Translation of <i>B. garinii</i> 5' vlsE homolog
SEQ ID NO:33	AY100633	<i>B. garinii</i> vls1
SEQ ID NO:34	AAN87825	Translation of <i>B. garinii</i> vls1
SEQ ID NO:35	AY100633	<i>B. garinii</i> vls2
SEQ ID NO:36	AAN87826	Translation of <i>B. garinii</i> vls2
SEQ ID NO:37	AY100633	<i>B. garinii</i> vls3
SEQ ID NO:38	AAN87827	Translation of <i>B. garinii</i> vls3

SEQ ID NO.	GENBANK NO.	DESCRIPTION
SEQ ID NO:39	AY100633	<i>B. garinii</i> vls4
SEQ ID NO:40	AAN87828	Translation of <i>B. garinii</i> vls4
SEQ ID NO:41	AY100633	<i>B. garinii</i> vls5
SEQ ID NO:42	AAN87829	Translation of <i>B. garinii</i> vls5
SEQ ID NO:43	AY100633	<i>B. garinii</i> vls6
SEQ ID NO:44	AAN87830	Translation of <i>B. garinii</i> vls6
SEQ ID NO:45	AY100633	<i>B. garinii</i> vls7
SEQ ID NO:46	AAN87831	Translation of <i>B. garinii</i> vls7
SEQ ID NO:47	AY100633	<i>B. garinii</i> vls8
SEQ ID NO:48	AAN87832	Translation of <i>B. garinii</i> vls8
SEQ ID NO:49	AY100633	<i>B. garinii</i> vls9
SEQ ID NO:50	AAN87833	Translation of <i>B. garinii</i> vls9
SEQ ID NO:51	AY100633	<i>B. garinii</i> vls10
SEQ ID NO:52	AAN87834	Translation of <i>B. garinii</i> vls10
SEQ ID NO:53	AY100633	<i>B. garinii</i> vls11
SEQ ID NO:54	AAN87835	Translation of <i>B. garinii</i> vls11
SEQ ID NO:55	AY100633	<i>B. garinii</i> truncated gene
SEQ ID NO:56	AAN87823	Translation of <i>B. garinii</i> truncated gene
SEQ ID NO:57	AY100628	vls silent cassette locus of <i>B. afzelii</i>
SEQ ID NO:58	AY100628	<i>B. afzelii</i> vls1
SEQ ID NO:59	AAN87809	Translation of <i>B. afzelii</i> vls1
SEQ ID NO:60	AY100628	<i>B. afzelii</i> vls2
SEQ ID NO:61	AAN87810	Translation of <i>B. afzelii</i> vls2
SEQ ID NO:62	AY100628	<i>B. afzelii</i> vls3
SEQ ID NO:63	AAN87811	Translation of <i>B. afzelii</i> vls3
SEQ ID NO:64	AY100628	<i>B. afzelii</i> vls4
SEQ ID NO:65	AAN87812	Translation of <i>B. afzelii</i> vls4
SEQ ID NO:66	AY100628	<i>B. afzelii</i> vls5
SEQ ID NO:67	AAN87813	Translation of <i>B. afzelii</i> vls5
SEQ ID NO:68	AY100628	<i>B. afzelii</i> vls6
SEQ ID NO:69	AAN87814	Translation of <i>B. afzelii</i> vls6
SEQ ID NO:70	AY100628	<i>B. afzelii</i> vls7
SEQ ID NO:71	AAN87815	Translation of <i>B. afzelii</i> vls7
SEQ ID NO:72	AY100628	<i>B. afzelii</i> vls8
SEQ ID NO:73	AAN87816	Translation of <i>B. afzelii</i> vls8
SEQ ID NO:74	AY100628	<i>B. afzelii</i> vls9a
SEQ ID NO:75	AAN87817	Translation of <i>B. afzelii</i> vls9a
SEQ ID NO:76	AY100628	<i>B. afzelii</i> vls10
SEQ ID NO:77	AAN87818	Translation of <i>B. afzelii</i> vls10
SEQ ID NO:78	AY100628	<i>B. afzelii</i> vls11
SEQ ID NO:79	AAN87819	Translation of <i>B. afzelii</i> vls11
SEQ ID NO:80	AY100628	<i>B. afzelii</i> vls12
SEQ ID NO:81	AAN87820	Translation of <i>B. afzelii</i> vls12
SEQ ID NO:82	AY100628	<i>B. afzelii</i> vls13
SEQ ID NO:83	AAN87821	Translation of <i>B. afzelii</i> vls13
SEQ ID NO:84	AY100628	<i>B. afzelii</i> vls14
SEQ ID NO:85	AAN87822	Translation of <i>B. afzelii</i> vls14

SEQ ID NO.	GENBANK NO.	DESCRIPTION
SEQ ID NO:86	AY100628	<i>B. afzelii</i> conserved protein
SEQ ID NO:87	AAN87823	Translation of <i>B. afzelii</i> conserved protein
SEQ ID NO:88	N/A	Nucleotides 1-2775 of AY100633 ( <i>B. garinii</i> )
SEQ ID NO:89	N/A	Nucleotides 3823-5897 of AY100633 ( <i>B. garinii</i> )
SEQ ID NO:90	N/A	Fragment of <i>B. garinii</i> vls5
SEQ ID NO:91	N/A	Amino acids 1-184 of AAN87829 ( <i>B. garinii</i> )
SEQ ID NO:92	N/A	Fragment of <i>B. garinii</i> vls8
SEQ ID NO:93	N/A	Amino acids 56-195 of AAN87832 ( <i>B. garinii</i> )
SEQ ID NO:94	N/A	Expressed ORF in pBG-10-1
SEQ ID NO:95	N/A	Protein sequence expressed by pBG-10-1
SEQ ID NO:96	N/A	Expressed ORF in pBA-13-1
SEQ ID NO:97	N/A	Protein sequence expressed by pBA-13-1
SEQ ID NO:98	N/A	Primer
SEQ ID NO:99	N/A	Primer
SEQ ID NO:100	N/A	Primer
SEQ ID NO:101	N/A	Primer
SEQ ID NO:102	N/A	Primer
SEQ ID NO:103	N/A	Primer
SEQ ID NO:104	N/A	Primer
SEQ ID NO:105	N/A	Primer
SEQ ID NO:106	N/A	17-bp direct repeat of <i>B. burgdorferi</i>
SEQ ID NO:107	N/A	EcoRI linker

## BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B. Arrangement of *vls* silent cassette regions of *B. garinii* Ip90 and *B. afzelii* ACAI. The orientation of the silent cassettes is indicated by a dashed arrow. Direct repeats are indicated by heavily weighted lines between silent cassettes. The location and orientation of conserved hypothetical protein genes are indicated at the 3' end of each locus. Restriction sites used for cloning and sequencing are also shown. (FIG. 1A) *B. garinii* Ip90. The cross-hatched bar indicates the location of P7-1 clone (Liang and Philipp, 1999) in the *vls* locus of Ip90. The locations of the telomeric repeat sequences (TRS) and the *vlsE*-like sequence are shown. (FIG. 1B) *B. afzelii* ACAI. The location and orientation of the *vls* cassettes and other features of this region are indicated as described above.

FIGS. 2A-2B. Alignment of predicted amino acid sequences of *vls* silent cassettes of *B. afzelii* ACAI (FIG. 2A) and *B. garinii* Ip90 (FIG. 2B) with the cassette region of *B. burgdorferi* B31 *vlsE*. Alignment for *B. afzelii* ACAI is based on cassette 1 and for *B. garinii* Ip90 based on cassette 10. The underlined residues at the end of cassette 9 in panel A are a continuation of the cassette following a frameshift. Identical amino acid sequences are shown as periods. The variable regions are indicated by shaded boxes and the lines under the shaded boxes represent the corresponding variable regions of *B. burgdorferi* B31. Gaps and predicted stop codons are indicated as dashes and asterisks, respectively.

FIG. 3. RT-PCR of *vlsE* sequences, using RNA from *B. afzelii* ACAI (lanes 1 and 2) and *B. garinii* Ip90 (lanes 3 and 4) as template. Lanes 2 and 4, with reverse transcriptase; lanes 1 and 3, controls without reverse transcriptase. DNA marker sizes (bp) are indicated on the left.

FIGS. 4A-4B. Alignment of the predicted amino acid sequences based on RT-PCR products from *vlsE* variants of *B. afzelii* ACAI (FIG. 4A) and *B. garinii* Ip90 (FIG. 4B). Alignments for *B. afzelii* ACAI and *B. garinii* Ip90 are based on the sequences of clones 2622 and 17, respectively. The variable regions labeled VR-I through VR-VI (FIG. 4A) and VR-II through VR-V (FIG. 4B) are indicated by the shaded regions. Only portions of VR-I and VR-VI are shown for ACAI. Identical amino acid sequences and gaps are shown as periods and dashes, respectively. Solid and dotted bars indicate the predicted minimum and maximum possible recombination events, respectively, resulting in the given *vlsE* variant. Solid lines indicate 100% sequence identity between the given position in the variant and silent cassette(s) indicated. Dashed lines mark the limits of maximum recombination. Asterisks above certain residues indicate sites of possible point mutations, as explained in the text. In regions where more than

one silent cassette matches the variant amino acid sequence, the matches were further analyzed at the nucleotide level.

FIG. 5. Hybridization of plasmid DNA of *B. afzelii* ACAI and *B. garinii* Ip90 with pJRZ53 probe. Lane 1, ACAI plasmid DNA; lane 2, ACAI plasmid DNA digested with EcoRI; lane 3, Ip90 plasmid DNA; and lane 4, Ip90 plasmid DNA digested with EcoRI. The size of EcoRI fragments containing *vls* sequences are indicated by arrows at left.

FIG. 6. Reactivity of human Lyme disease serum pool and a normal human serum pool with recombinant *Borrelia afzelii* Vls protein VLS-BA13.

FIG. 7. Effect of VLS-BA13 protein concentration on enzyme immunoassay reactivity of serum pools from Lyme disease human subjects and normal human subjects.

FIG. 8. Reactivity of mouse anti-*Borrelia burgdorferi* serum and normal mouse serum with recombinant *Borrelia afzelii* Vls protein VLS-BA13. The reactivity of normal mouse serum was below background levels.

FIG. 9. Effect of VLS-BA13 protein concentration on enzyme immunoassay reactivity of mouse anti-*B. burgdorferi* antiserum and normal mouse serum. The reactivity of normal mouse serum was below background levels.

FIG. 10. Reactivity of human Lyme disease serum pool and a normal human serum pool with recombinant *Borrelia garinii* Vls protein VLS-BG10.

FIG. 11. Effect of VLS-BG10 protein concentration on enzyme immunoassay reactivity of serum pools from Lyme disease human subjects and normal human subjects.

FIG. 12. Reactivity of mouse anti-*Borrelia burgdorferi* serum and normal mouse serum with recombinant *Borrelia garinii* Vls protein VLS-BG10. The reactivity of normal mouse serum was below background levels.

FIG. 13. Effect of VLS-BG10 protein concentration on enzyme immunoassay reactivity of mouse anti-*B. burgdorferi* antiserum and normal mouse serum. The reactivity of normal mouse serum was below background levels.

## DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The present work discloses the identification and characterization of an elaborate genetic system in the Lyme disease spirochete *Borrelia* that promotes extensive antigenic variation of a surface-exposed lipoprotein, VlsE.

Hybridization with the *B. burgdorferi* B31 *vls* silent cassette sequence in recombinant plasmid pJRZ53 was used in identifying the plasmids and DNA fragments containing *vls* sequences in *B. garinii* Ip90 and *B. afzelii* ACAI. The pJRZ53 probe hybridized exclusively to plasmids with an approximate size of 28 kb in both ACAI and Ip90. DNA fragments from these *B. garinii* Ip90 and *B. afzelii* ACAI plasmids were inserted into a recombinant lambda bacteriophage vector (lambda-DashI) and sequenced. The results showed *B. garinii* Ip90 to consist of 11 *vls* silent cassettes and *B. afzelii* ACAI of 14 *vls* silent cassettes.

With the exception of the junctions at *vls3/4* and *vls6/7*, the 11 *vls* silent cassettes of Ip90 are flanked by 18 bp direct repeat sequences in the 6 kb region. However, several of these cassettes (*vls1*, *vls4*, *vls6*, and *vls11*) are truncated (189 to 288 bp in length) relative to the other, full-length cassettes ranging in size from 573 to 594 bp. Unlike Ip90 and B31, the ACAI *vls* locus is located on an internal *EcoRI* fragment of a 28-kb linear plasmid, and its location relative to the plasmid telomeres is not known. The ACAI *vls* locus contained 13 complete and 1 partial silent cassettes with each cassette being flanked by an 18 bp direct repeat sequence.

These silent cassettes share 90% to 97% nucleotide sequence identity with one another within the Ip90 *vls* locus and 84% to 91% within the ACAI *vls* locus. Amino acid similarity to the B31 silent cassettes ranges from 51% to 62% for the Ip90 *vls* silent cassettes and from 50% to 65% for the ACAI *vls* silent cassettes. The nucleotide sequence and predicted amino acid sequence of *vlsE* in *B. burgdorferi* is provided in SEQ ID NO:1 and SEQ ID NO:2, respectively. The *vlsE* expression sites of Ip90 and ACAI have not been isolated, but transcripts of *vlsE* have been detected by reverse transcriptase PCR for both Ip90 and ACAI. In addition, the occurrence of sequence variation within the *vlsE* cassette region of these transcripts was verified. Mice infected experimentally with *B. garinii* and *B. afzelii* strains have been shown to express anti-VlsE antibodies (Liang *et al.*, 2000a). Additionally, a protein product of ~35 kDa expressed by *B. garinii* Ip90 reacts with antibodies against IR6, a peptide corresponding to invariant region 6 of the VlsE cassette region (Liang *et al.*, 1999a). The characteristics of the *vls* loci present in *B. garinii* Ip90 and *B. afzelii* ACAI are therefore similar to those found in *B. burgdorferi* B31.

Genetic variation involved in multi-gene families has been described in several other pathogenic microorganisms (Borst and Geaves, 1987; Borst *et al.*, 1995; Donelson, 1995). In the



context of combinatorial recombination, the genetic variation at the *vlsE* site is similar to that of the pilin-encoding genes of *Neisseria gonorrhoeae* (Seifert and So, 1988). The gonococcal pilus is primarily composed of repeating subunits of an 18-kilodalton pilin protein and is required for adherence of the bacterium to a variety of human cells (Swanson and Koomey, 1989). While the complete pilin genes are expressed only at two expression sites (*pilE1* and *pilE2*), multiple silent copies (*pilS*) containing portions of the pilin genes are found over a wide range on the gonococcal chromosome (Haas and Meyer, 1986). Through multiple combinatorial recombination events, a single gonococcal clone expressing one pilin stereotype can give rise to a large number of progeny that express antigenically distinctive pilin variants (Meyer *et al.*, 1982; Hagblom *et al.*, 1985; Segal *et al.*, 1986). The recombination between the expression and silent loci occurs predominantly through a non-reciprocal gene conversion mechanism (Haas and Meyer, 1986; Koomey *et al.*, 1987).

The coding sequences of the *Neisseria* pilin variants are divided into constant, semi-variable, and hypervariable regions (Haas and Meyer, 1986), which are analogous to the conserved and variable regions of the *vls* cassettes. The constant regions and a conserved DNA sequence (Sma/Cla repeat) located at the 3' end of all pilin loci are thought to pair the regions involved in recombination events (Wainwright *et al.*, 1994). In this context, the 18-bp direct repeats and the conserved regions of the *vls* cassettes in *B. garinii* and *B. afzelii* may play a similar role in recombination events. The silent loci of gonococcal pilin genes contain different regions of the complete pilin genes (Haas and Meyer, 1986), whereas the silent *vls* cassettes of *Borrelia* represent only the central cassette region of the *vlsE* gene.

Non-reciprocal recombinations also occur between the expressed and the silent genes encoding variant surface glycoproteins (VsGs) in African trypanosomes (Donelson, 1995). Based on similarities between the *vls* locus and the multi-gene families of the other pathogenic microorganisms and experimental data (Zhang and Norris, 1998b), it is likely that a unidirectional gene conversion mechanism is also active in the *vls* locus. The exact mechanism of *vls* recombination remains to be determined.

Variation of *Borreliae* surface proteins such as VlsE may also affect the organism's virulence and its ability to adapt to different micro-environments during infection of the mammalian host. Recent studies of a *Borrelia turicatae* mouse infection model that resembles Lyme disease showed that one serotype expressing VmpB exhibited more severe arthritic manifestations, whereas another expressing VmpA had more severe central nervous system involvement (Cadavid *et al.*, 1994). The numbers of *Borreliae* present in the joints and blood of

serotype B-infected mice were much higher than those of mice infected with serotype A, consistent with a relationship between Vmp serotype and disease severity (Pennington *et al.*, 1997). Antigenic variation of *Neisseria pilin* (Lambden *et al.*, 1980; Rudel *et al.*, 1992; Nassif *et al.*, 1993; Jonsson *et al.*, 1994) and Opa proteins (Kupsch *et al.*, 1993) is known to affect adherence of the organisms to human leukocytes and epithelial cells.

#### A. Antigenic variation in *B. hermsii*

A complex antigenic variation mechanism has been characterized in *Borrelia hermsii*, a relative of *B. afzelii* and *B. garinii* that causes relapsing fever (Balmelii and Piffatetti, 1996; Barbour, 1993; Donelson, 1995). Surface-exposed lipoproteins called variable major proteins (Vmps) are encoded by homologous genes located in 28- to 32-kb linear plasmids with covalently closed telomeres (Barbour and Garon, 1987; Kitten and Barbour, 1990). The *vmp* genes have been subdivided into two groups: small and large (Restrepo *et al.*, 1992). Large *vmp* genes such as *vmp7* and *vmp17* and small *vmp* genes such as *vmp1* and *vmp3* are approximately 1 kb and 0.6 kb in size, respectively. Each organism contains both small and large *vmp* genes in an unexpressed (silent) form in the so-called storage plasmids (Plasterk *et al.*, 1985). Only one *vmp* gene located near one of the telomeres of a different plasmid (called the expression plasmid) is expressed in each organism (Kitten and Barbour, 1990; Barbour *et al.*, 1991a). The nucleotide sequence and predicted amino acid sequence of an expressed *vmp* gene of *B. hermsii* are provided in SEQ ID NO:3 and SEQ ID NO:4, respectively. Antigenic variation occurs when the expressed *vmp* is replaced completely or partially by one of the silent *vmp* genes at the telomeric expression site through interplasmic recombination (Meier *et al.*, 1985; Plasterk *et al.*, 1985; Barbour *et al.*, 1991b), intraplasmic recombination (Restrepo *et al.*, 1994), and post-switch rearrangement (Restrepo and Barbour, 1994). The antigenic switch occurs spontaneously at a frequency of  $10^{-3}$  to  $10^{-4}$  per generation (Stoenner *et al.*, 1982).

#### B. Identification of *vls*

The present invention discloses a repetitive DNA sequence ~500 bp in length, which is present in multiple, nonidentical copies in a 28-kb linear plasmid of infectious *Borrelia burgdorferi*, *Borrelia garinii*, and *Borrelia afzelii*, the causative agents of Lyme disease. These DNA sequences encode polypeptides that have sequence similarity to the Variable Major Proteins (VMPs) of relapsing fever *Borreliae* (such as *B. hermsii*). VMPs are highly antigenic surface proteins, which the relapsing fever *Borreliae* are able to change through a genetic recombination mechanism, thereby evading the immune response. Antibodies against a

particular VMP protein are protective, resulting in rapid clearance of bacteria of the corresponding serotype. In *Borrelia burgdorferi*, *Borrelia garinii*, and *Borrelia afzelii*, VMP-like sequences (vls) are present on a 28-kb linear plasmid, and this plasmid appears to encode virulence factor(s) required for infectivity.

## 5 C. ELISAs

ELISAs may be used in conjunction with the invention. In an ELISA assay, proteins or peptides incorporating *Borrelia* Vls antigenic sequences are immobilized onto a selected surface, preferably a surface exhibiting a protein affinity such as the wells of a polystyrene microtiter plate. The antigenic proteins or peptides may be isolated or comprised within larger polypeptides. For example, an antigenic Vls peptide may be comprised within a larger polypeptide that also includes a moiety that is useful for anchoring the polypeptide to the selected surface. The anchoring moiety may be an amino acid sequence. Virtually any amino acid sequence may be added to the antigenic Vls sequence so long as it does not confound the results of the ELISA assay. Those of skill in the art would know how to select amino acid sequences that are antigenically neutral with regard to antibodies in the biological sample (including, but not limited to, whole blood, plasma, serum, cerebrospinal fluid, other body fluids, or tissue extracts) that is being tested.

After washing to remove incompletely adsorbed material, it is desirable to bind or coat the assay plate wells with a nonspecific protein that is known to be antigenically neutral with regard to the biological sample such as bovine serum albumin (BSA), casein or solutions of powdered milk. This allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antibodies in the biological sample onto the surface.

After binding of antigenic material to the well, coating with a non-reactive material to reduce background, and washing to remove unbound material, the immobilizing surface is contacted with the antisera or clinical or biological sample to be tested in a manner conducive to immune complex (antigen/antibody) formation. Such conditions preferably include diluting the sample with diluents such as BSA, solution or phosphate buffered saline (PBS)/Tween®. These added agents also tend to assist in the reduction of nonspecific background. The layered biological sample preparation is then allowed to incubate in the well for from about 1 to about 4 hr, at temperatures preferably on the order of about 25° to about 37°C. Following incubation with the diluted or undiluted biological sample, the antisera-contacted surface is washed so as to remove non-immunocomplexed material. A preferred washing procedure includes washing with a solution such as PBS/Tween®.

Following formation of specific immunocomplexes between the test sample and the bound antigen, and subsequent washing, the occurrence and even amount of immunocomplex formation may be determined by subjecting same to a second antibody having specificity for the first. To provide a detecting means, the second antibody will preferably have an associated enzyme that will generate a color development upon incubating with an appropriate chromogenic substrate. Thus, for example, one will desire to contact and incubate the antisera-bound surface with a urease, alkaline phosphatase or peroxidase-conjugated anti-human IgG for a period of time and under conditions which favor the development of immunocomplex formation (e.g., incubation for 2 hr at room temperature in a PBS-containing solution such as PBS/Tween®).

After incubation with the second enzyme-tagged antibody, and subsequent to washing to remove unbound material, the amount of label is quantified by incubation with a chromogenic substrate such as urea and bromocresol purple or 2,2'-azino-di-(3-ethyl-benzthiazoline)-6-sulfonic acid (ABTS) and H<sub>2</sub>O<sub>2</sub>, in the case of peroxidase as the enzyme label. Quantitation is then achieved by measuring the degree of color generation, e.g., using a visible spectrum spectrophotometer.

Alternatively, the ELISA assay may be performed where antibodies that bind immunologically to *Borrelia* Vls antigenic sequences are immobilized onto a selected surface. After binding of the antibody to the surface, coating with a non-reactive material to reduce background, and washing to remove unbound material, the immobilizing surface is contacted with the biological sample to be tested in a manner conducive to immune complex (antigen/antibody) formation. Following formation of specific immunocomplexes between the test sample and the bound antibody, and subsequent washing, immunocomplex formation may be determined using a second, labeled antibody. This approach enables the detection of an antigen in a biological sample.

#### D. Epitopic Core Sequences

The present invention is also directed to protein or peptide compositions, free from total cells and other peptides, which comprise a purified protein or peptide which incorporates an epitope that is immunologically cross-reactive with one or more anti-*Borrelia* VMP-like antibodies.

As used herein, the term "incorporating an epitope(s) that is immunologically cross-reactive with one or more anti-VMP-like antibodies" is intended to refer to a peptide or protein antigen which includes a primary, secondary or tertiary structure similar to an epitope located within a *Borrelia* VMP-like polypeptide. The level of similarity will generally be to such a degree that polyclonal antibodies directed against the *Borrelia* VMP-like polypeptide will also bind to, react with, or otherwise recognize, the cross-reactive peptide or protein antigen. Various immunoassay

methods may be employed in conjunction with such antibodies, such as, for example, Western blotting, ELISA, RIA, and the like, all of which are known to those of skill in the art.

The identification of *Borrelia* VMP-like epitopes, and/or their functional equivalents, suitable for use in vaccines is a relatively straightforward matter. For example, one may employ the methods of Hopp, as taught in U.S. Patent 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. The methods described in several other papers, and software programs based thereon, can also be used to identify epitopic core sequences (see, for example, Jameson and Wolf, 1988; Wolf *et al.*, 1988; U.S. Patent Number 4,554,101). The amino acid sequence of these "epitopic core sequences" may then be readily incorporated into peptides, either through the application of peptide synthesis or recombinant technology.

Preferred peptides for use in accordance with the present invention will generally be on the order of about 5 to about 50 amino acids in length, and more preferably about 8 to about 40 amino acids in length. Such peptides may be isolated or comprised within a larger polypeptide. It is proposed that shorter antigenic *Borrelia* VMP-like-derived peptide sequences will provide advantages in certain circumstances, for example, in the preparation of vaccines or in immunologic detection assays. Exemplary advantages include the ease of preparation and purification, the relatively low cost and improved reproducibility of production, and advantageous biodistribution.

It is proposed that particular advantages of the present invention may be realized through the preparation of synthetic peptides which include modified and/or extended epitopic/immunogenic core sequences which result in a "universal" epitopic peptide directed to *Borrelia* VMP-like and *Borrelia* VMP-like-related sequences. It is proposed that these regions represent those which are most likely to promote T-cell or B-cell stimulation in an animal, and, hence, elicit specific antibody production in such an animal.

An epitopic core sequence, as used herein, is a relatively short stretch of amino acids that is "complementary" to, and therefore will bind, antigen binding sites on *vls* protein-specific antibodies. Additionally or alternatively, an epitopic core sequence is one that will elicit antibodies that are cross-reactive with antibodies directed against the peptide compositions of the present invention. It will be understood that in the context of the present disclosure, the term "complementary" refers to amino acids or peptides that exhibit an attractive force towards each other. Thus, certain epitope core sequences of the present invention may be operationally defined in terms of their ability to compete with or perhaps displace the binding of the desired protein antigen with the corresponding protein-directed antisera.

In general, the size of the polypeptide antigen is not believed to be particularly crucial, so long as it is at least large enough to carry the identified core sequence or sequences. The smallest useful core sequence expected by the present disclosure would generally be on the order of about 5 amino acids in length, with sequences on the order of 8 or 25 being more preferred. Thus, this size will generally correspond to the smallest peptide antigens prepared in accordance with the invention. However, the size of the antigen may be larger where desired, so long as it contains a basic epitopic core sequence.

The identification of epitopic core sequences is known to those of skill in the art, for example, as described in U.S. Patent 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. Moreover, numerous computer programs are available for use in predicting antigenic portions of proteins (see *e.g.*, Jameson and Wolf, 1988; Wolf *et al.*, 1988). Computerized peptide sequence analysis programs (*e.g.*, DNASTar® software, DNASTar, Inc., Madison, Wisc.) may also be useful in designing synthetic *Borrelia* VMP-like peptides and peptide analogs in accordance with the present disclosure. In addition, epitope mapping may be performed, in which overlapping peptides corresponding to all regions of the protein are synthesized and tested for reactivity with antibodies directed against *vls* sequences. Reactivity of serum from animals or humans infected with Lyme disease *Borrelia*, and nonreactivity with serum from animals or patients that do not have Lyme disease would help to define those peptides that react sensitively and specifically with antibodies against Lyme disease *Borrelia*.

An epitopic core sequence may be comprised within a larger polypeptide. For example, an epitopic core sequence of the present invention may be comprised in a larger polypeptide, which also comprises a moiety that is useful for anchoring the polypeptide to the selected surface. The anchoring moiety may be an amino acid sequence.. These polypeptides would be particularly useful in the various immunoassay methods of the present invention. In a particular example, a peptide or polypeptide of the present invention may have a cysteine added at one end of the amino acid sequence to permit the addition of biotin. The biotinylated peptides or polypeptides could then be captured on streptavidin-coated surfaces. Those of skill in the art would know how to identify which polypeptides react sensitively and specifically with antibodies against Lyme disease *Borrelia*. For example, reactivity of serum from animals or humans infected with Lyme disease *Borrelia*, and nonreactivity with serum from animals or patients that do not have Lyme disease would help to define those polypeptides that react sensitively and specifically with antibodies against Lyme disease *Borrelia*.

Syntheses of epitopic sequences, or peptides which include an antigenic epitope within their sequence, are readily achieved using conventional synthetic techniques such as the solid phase method (*e.g.*, through the use of commercially available peptide synthesizer such as an Applied Biosystems Model 430A Peptide Synthesizer). Peptide antigens synthesized in this manner may then be aliquoted in predetermined amounts and stored in conventional manners, such as in aqueous solutions or, even more preferably, in a powder or lyophilized state pending use.

In general, due to the relative stability of peptides, they may be readily stored in aqueous solutions for fairly long periods of time if desired, *e.g.*, up to six months or more, in virtually any aqueous solution without appreciable degradation or loss of antigenic activity. However, where extended aqueous storage is contemplated it will generally be desirable to include agents including buffers such as Tris or phosphate buffers to maintain a pH of about 7.0 to about 7.5. Moreover, it may be desirable to include agents which will inhibit microbial growth, such as sodium azide or Merthiolate. For extended storage in an aqueous state it will be desirable to store the solutions at 4°C, or more preferably, frozen. Of course, where the peptides are stored in a lyophilized or powdered state, they may be stored virtually indefinitely, *e.g.*, in metered aliquots that may be rehydrated with a predetermined amount of water (preferably distilled) or buffer prior to use.

#### **E. Antibodies**

Means for preparing and characterizing antibodies are well known in the art (See, *e.g.*, Harlow and Lane, 1988; incorporated herein by reference). An antibody can be a polyclonal or a monoclonal antibody.

The methods for generating monoclonal antibodies (mAbs) generally begin along the same lines as those for preparing polyclonal antibodies. Briefly, a polyclonal antibody is prepared by immunizing an animal with an immunogenic composition in accordance with the present invention and collecting antisera from that immunized animal. A wide range of animal species can be used for the production of antisera. Typically the animal used for production of anti-antisera is a rabbit, a mouse, a rat, a hamster, a guinea pig or a goat. Because of the relatively large blood volume of rabbits, a rabbit is a preferred choice for production of polyclonal antibodies.

As is well known in the art, a given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide immunogen to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin or rabbit serum albumin can also be used as carriers. Means for conjugating a

polypeptide to a carrier protein are well known in the art and include glutaraldehyde, *m*-maleimidobenzoyl-*N*-hydroxysuccinimide ester, carbodiimide and bis-biazotized benzidine.

As is also well known in the art, the immunogenicity of a particular immunogen composition can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Exemplary and preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvant and aluminum hydroxide adjuvant.

The amount of immunogen composition used in the production of polyclonal antibodies varies upon the nature of the immunogen as well as the animal used for immunization. A variety of routes can be used to administer the immunogen (subcutaneous, intramuscular, intradermal, intravenous and intraperitoneal). The production of polyclonal antibodies may be monitored by sampling blood of the immunized animal at various points following immunization. A second, booster, injection may also be given. The process of boosting and titering is repeated until a suitable titer is achieved. When a desired level of immunogenicity is obtained, the immunized animal can be bled and the serum isolated and stored, and/or the animal can be used to generate mAbs.

mAbs may be readily prepared through use of well-known techniques, such as those exemplified in U.S. Patent 4,196,265, incorporated herein by reference. Typically, this technique involves immunizing a suitable animal with a selected immunogen composition, *e.g.*, a purified or partially purified LCRF protein, polypeptide or peptide. The immunizing composition is administered in a manner effective to stimulate antibody producing cells. Rodents such as mice and rats are preferred animals, however, the use of rabbit, sheep, or frog cells is also possible. The use of rats may provide certain advantages (Goding, 1986), but mice are preferred, with the BALB/c mouse being most preferred as this is most routinely used and generally gives a higher percentage of stable fusions.

Following immunization, somatic cells with the potential for producing antibodies, specifically B-lymphocytes (B-cells), are selected for use in the mAb generating protocol. These cells may be obtained from biopsied spleens, tonsils or lymph nodes, or from a peripheral blood sample. Spleen cells and peripheral blood cells are preferred, the former because they are a rich source of antibody-producing cells that are in the dividing plasmablast stage, and the latter because peripheral blood is easily accessible. Often, a panel of animals will have been immunized and the spleen of animal with the highest antibody titer will be removed and the spleen lymphocytes



obtained by homogenizing the spleen with a syringe. Typically, a spleen from an immunized mouse contains approximately  $5 \times 10^7$  to  $2 \times 10^8$  lymphocytes.

The antibody-producing B lymphocytes from the immunized animal are then fused with cells of an immortal myeloma cell, generally one of the same species as the animal that was immunized. Myeloma cell lines suited for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas).

Any one of a number of myeloma cells may be used, as are known to those of skill in the art (Goding, 1986; Campbell, 1984). For example, where the immunized animal is a mouse, one may use P3-X63/Ag8, X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0 Bul; for rats, one may use R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210; and U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6 are all useful in connection with human cell fusions.

One preferred murine myeloma cell is the NS-1 myeloma cell line (also termed P3-NS-1-Ag4-1), which is readily available from the NIGMS Human Genetic Mutant Cell Repository by requesting cell line repository number GM3573. Another mouse myeloma cell line that may be used is the 8-azaguanine-resistant mouse murine myeloma SP2/0 non-producer cell line.

Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually comprise mixing somatic cells with myeloma cells in a 2:1 ratio, though the ratio may vary from about 20:1 to about 1:1, respectively, in the presence of an agent or agents (chemical or electrical) that promote the fusion of cell membranes. Fusion methods using Sendai virus have been described (Kohler and Milstein, 1975; 1976), and those using polyethylene glycol (PEG), such as 37% (v/v) PEG, by Gefer *et al.*, (1977). The use of electrically induced fusion methods is also appropriate (Goding, 1986).

Fusion procedures usually produce viable hybrids at low frequencies, about  $1 \times 10^{-6}$  to  $1 \times 10^{-8}$ . However, this does not pose a problem, as the viable, fused hybrids are differentiated from the parental, unfused cells (particularly the unfused myeloma cells that would normally continue to divide indefinitely) by culturing in a selective medium. The selective medium is generally one that contains an agent that blocks the *de novo* synthesis of nucleotides in the tissue culture media. Exemplary and preferred agents are aminopterin, methotrexate, and azaserine. Aminopterin and methotrexate block *de novo* synthesis of both purines and pyrimidines, whereas azaserine blocks only purine synthesis. Where aminopterin or methotrexate is used, the media is supplemented with

hypoxanthine and thymidine as a source of nucleotides (HAT medium). Where azaserine is used, the media is supplemented with hypoxanthine.

The preferred selection medium is HAT. Only cells capable of operating nucleotide salvage pathways are able to survive in HAT medium. The myeloma cells are defective in key enzymes of the salvage pathway, *e.g.*, hypoxanthine phosphoribosyl transferase (HPRT), and they cannot survive. The B-cells can operate this pathway, but they have a limited life span in culture and generally die within about two weeks. Therefore, the only cells that can survive in the selective media are those hybrids formed from myeloma and B-cells.

This culturing provides a population of hybridomas from which specific hybridomas are selected. Typically, selection of hybridomas is performed by culturing the cells by single-clone dilution in microtiter plates, followed by testing the individual clonal supernatants (after about two to three weeks) for the desired reactivity. The assay should be sensitive, simple and rapid, such as radioimmunoassays, enzyme immunoassays, cytotoxicity assays, plaque assays, dot immunobinding assays, and the like.

The selected hybridomas would then be serially diluted and cloned into individual antibody-producing cell lines, which clones can then be propagated indefinitely to provide mAbs. The cell lines may be exploited for mAb production in two basic ways. A sample of the hybridoma can be injected (often into the peritoneal cavity) into a histocompatible animal of the type that was used to provide the somatic and myeloma cells for the original fusion. The injected animal develops tumors secreting the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can then be tapped to provide mAbs in high concentration. The individual cell lines could also be cultured *in vitro*, where the mAbs are naturally secreted into the culture medium from which they can be readily obtained in high concentrations. mAbs produced by either means may be further purified, if desired, using filtration, centrifugation and various chromatographic methods such as HPLC or affinity chromatography.

#### **F. Immunoprecipitation**

The antibodies of the present invention are particularly useful for the isolation of antigens by immunoprecipitation. Immunoprecipitation involves the separation of the target antigen-antibody complexes from a complex mixture, and is used to discriminate or isolate minute amounts of protein. For the isolation of membrane proteins cells must be solubilized into detergent micelles. Nonionic detergents are preferred, since other agents, such as bile salts, precipitate at acid pH or in the presence of bivalent cations.

In an alternative embodiment the antibodies of the present invention are useful for the close juxtaposition of two antigens. This is particularly useful for increasing the localized concentration of antigens, e.g., enzyme-substrate pairs.

#### G. Western Blots

The compositions of the present invention will find great use in immunoblot or western blot analysis. The anti-*Borrelia* VMP-like antibodies may be used as high-affinity primary reagents for the identification of proteins immobilized onto a solid support matrix, such as nitrocellulose, nylon or combinations thereof. In conjunction with immunoprecipitation, followed by gel electrophoresis, these may be used as a single step reagent for use in detecting antigens against which secondary reagents used in the detection of the antigen cause an adverse background. This is especially useful when the antigens studied are immunoglobulins (precluding the use of immunoglobulins binding bacterial cell wall components), the antigens studied cross-react with the detecting agent, or they migrate at the same relative molecular weight as a cross-reacting signal.

Immunologically-based detection methods for use in conjunction with Western blotting include enzymatically-, radiolabel-, or fluorescently-tagged secondary antibodies against the toxin moiety are considered to be of particular use in this regard.

#### H. Vaccines

An important aspect of the invention is the recognition that *Borrelia* VMP-like sequences recombine at the *vlsE* site, with the result that antigenic variation is virtually limitless. Multiclonal populations therefore can exist in an infected patient so that immunological defenses are severely tested if not totally overwhelmed. Thus there is now the opportunity to develop more effective combinations of immunogens for protection against *Borrelia* infections or as preventive inoculations such as in the form of cocktails of multiple antigenic variants based on a series of combinatorial VMP-like antigens.

VMP-like protein preparations may be administered in several ways, either locally or systemically in pharmaceutically acceptable formulations. Amounts appropriate for administration are determined on an individual basis depending on such factors as age and sex of the subject, as well as physical condition and weight. Such determinations are well within the skill of the practitioner in the medical field.

Other methods of administration may include injection of *Borrelia* VMP-like DNAs into vaccine recipients (human or animal) driven by an appropriate promoter such as CMV, (so called

DNA vaccines). Such preparations could be injected subcutaneously or intramuscularly, administered orally, or introduced into the skin on metal particles propelled by high-pressure gas. DNA vaccination techniques are currently well past the initial development stage and have shown promise as vaccination strategies.

5 The present invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared most directly from immunogenic *Borrelia* VMP-like peptides prepared in a manner disclosed herein. Preferably the antigenic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a  
10 desired vehicle.

The preparation of vaccines which contain *Borrelia* VMP-like peptide or polypeptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Patents 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all incorporated  
15 herein by reference. Typically, such vaccines are prepared as injectables. Either as liquid solutions or suspensions: solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain auxiliary substances  
20 such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines.

Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Vaccines may also be administered orally. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol,  
25 lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10 to about 95% of active ingredient, preferably about 25 to about 70%.

The *Borrelia* VMP-like-derived peptides or polypeptides of the present invention may be  
30 formulated into the vaccine as neutral or salt forms. It is anticipated that many VMP-like-derived peptides or polypeptides with different sequences could be incorporated into a single vaccine, in effect producing a combinatorial vaccine. Pharmaceutically-acceptable salts, include the acid addition salts (formed with the free amino groups of the peptide) and those which are formed with

inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including, *e.g.*, the capacity of the individual's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size of the host.

Various methods of achieving adjuvant effect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70° to about 101°C for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin treated (Fab) antibodies to albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of Gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed.

In many instances, it will be desirable to have multiple administrations of the vaccine, usually not exceeding six vaccinations, more usually not exceeding four vaccinations and preferably one or more, usually at least about three vaccinations. The vaccinations will normally be at from two to twelve week intervals, more usually from three to five week intervals. Periodic boosters at intervals of 1-5 years, usually three years, will be desirable to maintain protective levels

of the antibodies. The course of the immunization may be followed by assays for antibodies for the supernatant antigens. The assays may be performed by labeling with conventional labels, such as radionucleotides, enzymes, fluorescents, and the like. These techniques are well known and may be found in a wide variety of patents, such as U.S. Patent Nos. 3,791,932; 4,174,384 and 3,949,064, as illustrative of these types of assays.

## I. Nucleic Acids

The present invention provides the nucleotide sequences of the *vls* gene in *B. garinii* and *B. afzelii*. It is contemplated that the isolated nucleic acids of the present invention may be put under the control of a promoter. The promoter may be the promoter that is naturally associated with the *vls* gene or it may be a recombinant or heterologous promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with a DNA segment encoding a *Borrelia* VMP-like peptide in its natural environment. Such promoters may include promoters normally associated with other genes, and/or promoters isolated from any viral, prokaryotic (e.g., bacterial), eukaryotic (e.g., fungal, yeast, plant, or animal) cell.

Naturally, it will be important to employ a promoter that effectively directs the expression of the DNA segment in the cell type, organism, or even animal, chosen for expression. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook *et al.*, 2001. The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins or peptides. Appropriate promoter/expression systems contemplated for use in high-level expression include, but are not limited to, the *Pichia* expression vector system (Pharmacia LKB Biotechnology), a baculovirus system for expression in insect cells, or any suitable yeast or bacterial expression system.

In connection with expression embodiments to prepare recombinant proteins and peptides, it is contemplated that longer DNA segments will most often be used, with DNA segments encoding the entire peptide sequence being most preferred. However, it will be appreciated that the use of shorter DNA segments to direct the expression of *Borrelia* VMP-like peptides or epitopic core regions, such as may be used to generate anti-*Borrelia* VMP-like antibodies, also falls within the scope of the invention. DNA segments that encode *Borrelia* VMP-like peptide antigens from about 10 to about 100 amino acids in length, or more preferably, from about 20 to about 80 amino acids in length, or even more preferably, from about 30 to about 70 amino acids in length are contemplated to be particularly useful.

In addition to their use in directing the expression of *Borrelia* VMP-like peptides of the present invention, the nucleic acid sequences contemplated herein also have a variety of other uses. For example, they also have utility as probes or primers in nucleic acid hybridization embodiments. As such, it is contemplated that nucleic acid segments that comprise a sequence region that consists of at least about a 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, an about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 nucleotide long contiguous DNA segment of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96 will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200, 300, 400, 500, (including all intermediate lengths) and those up to and including full-length sequences will also be of use in certain embodiments.

The ability of such nucleic acid probes to specifically hybridize to *Borrelia* VMP-like-encoding sequences will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are envisioned, including the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

Nucleic acid molecules having sequence regions consisting of contiguous nucleotide stretches of about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 150, 175, 200,

300, 400, 500 or more, identical or complementary to the DNA sequence of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96, are particularly contemplated as hybridization probes for use in, *e.g.*, Southern and Northern blotting. Smaller fragments will generally find use in hybridization embodiments, wherein the length of the contiguous complementary region may be varied, such as between about 10-14 and up to about 100 nucleotides, but larger contiguous complementary stretches may be used, according to the length complementary sequences one wishes to detect.

The use of a hybridization probe of about 14 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules having contiguous complementary sequences over stretches greater than 14 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of about 15 to about 20 contiguous nucleotides, or even longer where desired.

Of course, fragments may also be obtained by other techniques such as, *e.g.*, by mechanical shearing or by restriction enzyme digestion. Small nucleic acid segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as PCR<sup>TM</sup>, by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular biology.

Accordingly, the nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of DNA fragments. Depending on the application envisioned, one will desire to employ varying conditions of hybridization to achieve varying degrees of selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the



hybrids, *e.g.*, conditions of high stringency where one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50°C to about 70°C. Such selective conditions tolerate little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating *Borrelia* VMP-like-encoding DNA segments. Detection of DNA segments *via* hybridization is well-known to those of skill in the art, and the teachings of U.S. Patents 4,965,188 and 5,176,995 (each incorporated herein by reference) are exemplary of the methods of hybridization analyses. Teachings such as those found in the texts of Maloy *et al.*, 1994; Segal, 1976; Prokop, 1991; and Kuby, 1994, are particularly relevant.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template or where one seeks to isolate *Borrelia* VMP-like-encoding sequences from related species, functional equivalents, or the like, less stringent hybridization conditions will typically be needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ conditions such as about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

In certain embodiments, it will be advantageous to employ nucleic acid sequences of the present invention in combination with an appropriate means, such as a label, for determining hybridization. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of giving a detectable signal. In preferred embodiments, one will likely desire to employ a fluorescent label or an enzyme tag, such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmental undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

In general, it is envisioned that the hybridization probes described herein will be useful both as reagents in solution hybridization as well as in embodiments employing a solid phase. In embodiments involving a solid phase, the test DNA (or RNA) is adsorbed or otherwise affixed to a

selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to specific hybridization with selected probes under desired conditions. The selected conditions will depend on the particular circumstances based on the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, *etc.*). Following washing of the hybridized surface so as to remove nonspecifically bound probe molecules, specific hybridization is detected, or even quantitated, by means of the label.

Isolated nucleic acids encoding *vls* or *vls*-related genes are contemplated to be particularly useful in connection with this invention. Any recombinant *vls* combining any of the *vlsE* expression site loci and/or silent *vls* cassette would likewise be very useful with the methods of the invention.

Isolation of the DNA encoding VMP-like polypeptides allows one to use methods well known to those of skill in the art, and as herein described, to make changes in the codons for specific amino acids such that the codons are "preferred usage" codons for a given species. Thus for example, preferred codons will vary significantly for bacterial species as compared with mammalian species; however, there are preferences even among related species. Shown below is a preferred codon usage table for humans. Isolation of spirochete DNA encoding VMP-like proteins will allow substitutions for preferred human codons, although expressed polypeptide product from human DNA is expected to be homologous to bacterial VMP-like proteins and so would be expected to be structurally and functionally equivalent to VMP-like proteins isolated from a spirochete. However, substitutions of preferred human codons may improve expression in the human host, thereby improving the efficiency of potential DNA vaccines. This method may also be useful in achieving improved expression of the recombinant VMP-like protein in *E. coli* or any of a variety of prokaryotic and eukaryotic cells.

**TABLE 2**  
**Codon Frequency in Homo sapiens**

Codon	$\nu^b$	Total # <sup>a</sup>	Codon	$\nu^b$	Total # <sup>a</sup>	Codon	$\nu^b$	Total # <sup>a</sup>	Codon	$\nu^b$	Total # <sup>a</sup>
UUU	16.6	72711	UCU	14.0	62953	UAU	12.3	55039	UGU	9.5	42692
UUC	21.4	95962	UCC	17.7	79482	UAC	17.0	76480	UGC	12.8	57368
UUA	6.3	28202	UCA	10.7	48225	UAA	0.7	2955	UGA	1.2	5481
UUG	11.5	51496	UCG	4.4	19640	UAG	0.5	2181	UGG	13.5	59982
CUU	11.7	52401	CCU	16.7	74975	CAU	9.6	43193	CGU	4.6	20792
CUC	19.5	87696	CCC	20.0	89974	CAC	14.6	65533	CGC	11.0	49507
CUA	6.3	28474	CCA	16.2	72711	CAA	11.4	51146	CGA	5.9	26551
CUG	40.6	182139	CCG	6.9	30863	CAG	33.7	151070	CGG	11.3	50682
AUU	15.7	70652	ACU	12.8	57288	AAU	16.6	74401	AGU	11.1	49894
AUC	23.7	106390	ACC	21.1	94793	AAC	21.1	94725	AGC	19.1	85754
AUA	6.7	30139	ACA	14.7	66136	AAA	23.2	104221	AGA	10.8	48369
AUG	22.6	101326	ACG	6.7	30059	AAG	33.9	152179	AGG	10.9	48882
GUU	10.6	47805	GCU	18.7	83800	GAU	22.0	98712	GCU	11.2	50125
GUC	15.6	70189	GCC	29.2	130966	GAC	27.0	121005	GGC	24.0	107571
GUA	6.6	29659	GCA	15.3	68653	GAA	27.8	124852	GGA	16.9	75708
GUG	30.0	134750	GCG	7.5	33759	GAG	40.8	182943	GGG	16.7	74859

Coding GC 52.96% 1st letter GC 55.98% 2nd letter GC 42.29% 3rd letter GC 60.60%

<sup>a</sup> Total 4489120

<sup>b</sup>  $\nu$  = Frequency per 1000

The definition of a "VMP-like sequence" or "VMP-related gene" as used herein, is a gene that hybridizes, under relatively stringent hybridization conditions (see, *e.g.*, Sambrook *et al.*, 2001), to DNA sequences presently known to include related gene sequences.

5 To prepare a VMP-like gene segment or cDNA one may follow the teachings disclosed herein and also the teachings of any patents or scientific documents specifically referenced herein. One may obtain a rVMP- or other related-encoding DNA segments using molecular biological techniques, such as polymerase chain reaction (PCR™) or screening of a cDNA or genomic library, using primers or probes  
10 with sequences based on the above nucleotide sequence. Such single- or double-stranded DNA segments may be readily prepared by, for example, directly synthesizing the fragments by chemical means, by application of nucleic acid reproduction technology, such as the PCR™ technology of U.S. Patents 4,683,195 and 4,683,202 (herein incorporated by reference). The practice of these techniques is a  
15 routine matter for those of skill in the art, as taught in various scientific texts (see *e.g.*, Sambrook *et al.*, 2001), incorporated herein by reference. Certain documents further particularly describe suitable mammalian expression vectors, *e.g.*, U.S. Patent 5,168,050, incorporated herein by reference. The VMP-like genes and DNA segments that are particularly preferred for use in certain aspects of the present methods are  
20 those encoding VMP-like and VMP-related polypeptides.

It is also contemplated that one may clone other additional genes or cDNAs that encode a VMP-like or VMP-related peptide, protein or polypeptide. The techniques for cloning DNA molecules, *i.e.*, obtaining a specific coding sequence from a DNA library that is distinct from other portions of DNA, are well known in the  
25 art. This can be achieved by, for example, screening an appropriate DNA library which relates to the cloning of a *vls* gene such as from the variable region of that gene. The screening procedure may be based on the hybridization of oligonucleotide probes, designed from a consideration of portions of the amino acid sequence of known DNA sequences encoding related *Borrelia* proteins. The operation of such screening  
30 protocols is well known to those of skill in the art and are described in detail in the scientific literature, for example, see Sambrook *et al.*, 2001.

Techniques for introducing changes in nucleotide sequences that are designed to alter the functional properties of the encoded proteins or polypeptides are well known in the art, *e.g.*, U.S. Patent 4,518,584, incorporated herein by reference, which techniques are also described in further detail herein. Such modifications include the deletion, insertion or substitution of bases, which may or may not result in changes in the amino acid sequence. Changes may be made to increase the activity of a protein, to increase its biological stability or half-life, to change its glycosylation pattern, and the like. All such modifications to the nucleotide sequences are encompassed by this invention.

**I. Biological Functional Equivalents**

Modification and changes may be made in the structure of the peptides of the present invention and DNA segments which encode them and still obtain a functional molecule that encodes a protein or peptide with desirable characteristics. The following is a discussion based upon changing the amino acids of a protein to create an equivalent, or even an improved, second-generation molecule. The amino acid changes may be achieved by changing the codons of the DNA sequence, according to the following codon table:

TABLE 3

Amino Acids				Codons				
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Cysteine	Cys	C	UGC	UGU				
Aspartic acid	Asp	D	GAC	GAU				
Glutamic acid	Glu	E	GAA	GAG				
Phenylalanine	Phe	F	UUC	UUU				
Glycine	Gly	G	GGA	GGC	GGG	GGU		
Histidine	His	H	CAC	CAU				
Isoleucine	Ile	I	AUA	AUC	AUU			
Lysine	Lys	K	AAA	AAG				
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU
Methionine	Met	M	AUG					
Asparagine	Asn	N	AAC	AAU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Glutamine	Gln	Q	CAA	CAG				
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Valine	Val	V	GUA	GUC	GUG	GUU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				

For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporate herein by reference). It is accepted that the relative  
5 hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

Each amino acid has been assigned a hydropathic index on the basis of their  
10 hydrophobicity and charge characteristics (Kyte and Doolittle, 1982), these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

15 It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.*, still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydropathic indices are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and those  
20 within  $\pm 0.5$  are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Patent 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological  
25 property of the protein.

As detailed in U.S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0  $\pm$  1); glutamate (+3.0  $\pm$  1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5  $\pm$  1); alanine (-0.5); histidine (-0.5); cysteine (-1.0);  
30 methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular,

an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

#### 10 J. Site-Specific Mutagenesis

Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, through specific mutagenesis of the underlying DNA. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 1 to 10 residues on both sides of the junction of the sequence being altered.

In general, the technique of site-specific mutagenesis is well known in the art, as exemplified by various publications. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double



stranded vector which includes within its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I  
5 Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

10 The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may  
15 be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

#### H. Expression of VMP-like Proteins

A particular aspect of this invention provides novel ways in which to utilize VMP-like DNA segments and recombinant vectors comprising *vls* DNA segments. As is well known to those of skill in the art, many such vectors are readily available,  
20 one particular detailed example of a suitable vector for expression in mammalian cells is that described in U. S. Patent 5,168,050, incorporated herein by reference. However, there is no requirement that a highly purified vector be used, so long as the coding segment employed encodes a VMP-like protein and does not include any coding or regulatory sequences that would have an adverse effect on cells. Therefore,  
25 it will also be understood that useful nucleic acid sequences may include additional residues, such as additional non-coding sequences flanking either of the 5' or 3' portions of the coding including, for example, promoter regions, or may include various internal sequences, *i.e.*, introns, which are known to occur within genes.

After identifying an appropriate VMP-encoding gene or DNA molecule, it may  
30 be inserted into any one of the many vectors currently known in the art, so that it will direct the expression and production of the VMP-like protein when incorporated into a host cell. In a recombinant expression vector, the coding portion of the DNA

segment is positioned under the control of a promoter. The promoter may be in the form of the promoter which is naturally associated with a VMP-encoding gene, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment, for example, using recombinant cloning and/or PCR™ technology, in connection with the compositions disclosed herein.

The use of recombinant promoters to achieve protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook *et al.*, (2001).

For the expression of VMP-like proteins, once a suitable (full-length if desired) clone or clones have been obtained, whether they be cDNA based or genomic, one may proceed to prepare an expression system for the recombinant preparation of VMP-like proteins. The engineering of DNA segment(s) for expression in a prokaryotic or eukaryotic system may be performed by techniques generally known to those of skill in recombinant expression. It is believed that virtually any expression system may be employed in the expression of VMP-like proteins.

VMP-like proteins may be successfully expressed in eukaryotic expression systems, however, it is also envisioned that bacterial expression systems may be preferred for the preparation of VMP-like proteins for all purposes. The DNA or cDNA encoding VMP-like proteins may be separately expressed in bacterial systems, with the encoded proteins being expressed as fusions with beta-galactosidase, ubiquitin, *Schistosoma japonicum* glutathione S-transferase, green fluorescent protein, polyhistidine and the like. It is believed that bacterial expression will ultimately have advantages over eukaryotic expression in terms of ease of use and quantity of materials obtained thereby.

It is proposed that transformation of host cells with DNA segments encoding VMP-like proteins will provide a convenient means for obtaining VMP-like peptides. Both cDNA and genomic sequences are suitable for eukaryotic expression, as the host cell will, of course, process the genomic transcripts to yield functional mRNA for translation into protein.

It is similarly believed that almost any eukaryotic expression system may be utilized for the expression of VMP-like proteins, *e.g.*, baculovirus-based, glutamine synthase-based or dihydrofolate reductase-based systems could be employed. However, in preferred embodiments, it is contemplated that plasmid vectors incorporating an origin of replication and an efficient eukaryotic promoter, as exemplified by the eukaryotic vectors of the pCMV series, such as pCMV5, will be of most use.

For expression in this manner, one would position the coding sequences adjacent to and under the control of the promoter. It is understood in the art that to bring a coding sequence under the control of such a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame of the protein between about 1 and about 50 nucleotides "downstream" of (*i.e.*, 3' of) the chosen promoter.

Where eukaryotic expression is contemplated, one will also typically desire to incorporate into the transcriptional unit which includes VMP-like protein, an appropriate polyadenylation site (*e.g.*, 5'-AATAAA-3') if one was not contained within the original cloned segment. Typically, the poly A addition site is placed about 30 to 2000 nucleotides "downstream" of the termination site of the protein at a position prior to transcription termination.

Translational enhancers may also be incorporated as part of the vector DNA. Thus the DNA constructs of the present invention should also preferable contain one or more 5' non-translated leader sequences which may serve to enhance expression of the gene products from the resulting mRNA transcripts. Such sequences may be derived from the promoter selected to express the gene or can be specifically modified to increase translation of the RNA. Such regions may also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence (Griffiths, *et al.*, 1993).

Such "enhancer" sequences may be desirable to increase or alter the transcription of translational efficiency of the resultant mRNA. The present invention is not limited to constructs where the enhancer is derived from the native 5'-

nontranslated promoter sequence, but may also include non-translated leader sequences derived from other non-related promoters such as other enhancer transcriptional activators or genes.

5 It is contemplated that virtually any of the commonly employed host cells can be used in connection with the expression of VMPs in accordance herewith. Examples include cell lines typically employed for eukaryotic expression such as 239, AtT-20, HepG2, VERO, HeLa, CHO, WI 38, BHK, COS-7, RIN and MDCK cell lines.

10 It is contemplated that VMP-like protein may be "overexpressed", *i.e.*, expressed in increased levels relative to its natural expression in *Borrelia* cells, or even relative to the expression of other proteins in a recombinant host cell containing VMP-encoding DNA segments. Such overexpression may be assessed by a variety of methods, including radio-labeling and/or protein purification. However, simple and direct methods are preferred, for example, those involving SDS/PAGE and protein  
15 staining or Western blotting, followed by quantitative analyses, such as densitometric scanning of the resultant gel or blot. A specific increase in the level of the recombinant protein or peptide in comparison to the level in natural VMP-producing animal cells is indicative of overexpression, as is a relative abundance of the specific protein in relation to the other proteins produced by the host cell and, *e.g.*, visible on a  
20 gel.

As used herein, the term "engineered" or "recombinant" cell is intended to refer to a cell into which a recombinant gene, such as a gene encoding a VMP-like peptide has been introduced. Therefore, engineered cells are distinguishable from naturally occurring cells which do not contain a recombinantly introduced gene.  
25 Engineered cells are thus cells having a gene or genes introduced through the hand of man. Recombinantly introduced genes will either be in the form of a cDNA gene (*i.e.*, they will not contain introns), a copy of a genomic gene, or will include genes positioned adjacent to a promoter not naturally associated with the particular introduced gene.

It will be understood that recombinant VMP-like proteins may differ from naturally produced VMP-like proteins in certain ways. In particular, the degree of post-translational modifications, such as, for example, lipidation, glycosylation and phosphorylation may be different between the recombinant VMP-like and the VMP-like polypeptide purified from a natural source, such as *Borrelia*.

After identifying an appropriate DNA molecule by any or a combination of means as described above, the DNA may then be inserted into any one of the many vectors currently known in the art and transferred to a prokaryotic or eukaryotic host cell where it will direct the expression and production of the so-called "recombinant" version of the protein. The recombinant host cell may be selected from a group consisting of *S. mutans*, *E. coli*, *S. cerevisiae*, *Bacillus* sp., *Lactococci* sp., *Enterococci* sp., or *Salmonella* sp. In certain preferred embodiments, the recombinant host cell will have a *recA* phenotype.

Where the introduction of a recombinant version of one or more of the foregoing genes is required, it will be important to introduce the gene such that it is under the control of a promoter that effectively directs the expression of the gene in the cell type chosen for engineering. In general, one will desire to employ a promoter that allows constitutive (constant) expression of the gene of interest. The use of these constitutive promoters will ensure a high, constant level of expression of the introduced genes. The level of expression from the introduced genes of interest can vary in different clones, probably as a function of the site of insertion of the recombinant gene in the chromosomal DNA. Thus, the level of expression of a particular recombinant gene can be chosen by evaluating different clones derived from each transfection study; once that line is chosen, the constitutive promoter ensures that the desired level of expression is permanently maintained. It may also be possible to use promoters that are subject to regulation, such as those regulated by the presence of lactose analog or by the expression of bacteriophage T7 DNA polymerase.

Technology for introduction of DNA into cells is well-known to those of skill in the art. Five general methods for delivering a gene into cells have been described: (1) chemical methods (Graham and VanDerEb, 1973); (2) physical methods such as microinjection (Capecchi, 1980), electroporation (Wong and Neumann, 1982; Fromm *et al.*, 1985) and the gene gun (Yang *et al.*, 1990); (3) viral vectors (Clapp, 1993;

Danos and Heard, 1992; Eglitis and Anderson, 1988); (4) receptor-mediated mechanisms (Wu *et al.*, 1991; Curiel *et al.*, 1991; Wagner *et al.*, 1992); and (5) direct injection of purified DNA into human or animals.

#### G. Liposomes and Nanocapsules

5       The formation and use of liposomes is generally known to those of skill in the art (see for example, Couvreur *et al.*, 1991 which describes the use of liposomes and nanocapsules in the targeted antibiotic therapy of intracellular bacterial infections and diseases). Recently, liposomes were developed with improved serum stability and circulation half-times of substances, including DNA (Gabizon and Papahadjopoulos, 10       1988; Allen and Choun, 1987). The following is a brief description of this and other DNA delivery modes.

Nanocapsules can generally entrap compounds in a stable and reproducible way (Henry-Michelland *et al.*, 1987). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1  $\mu\text{m}$ ) should be 15       designed using polymers able to be degraded *in vivo*. Biodegradable polyalkylcyanoacrylate nanoparticles that meet these requirements are contemplated for use in the present invention, and such particles may be easily made, as described (Couvreur *et al.*, 1984; 1988).

Liposomes are formed from phospholipids that are dispersed in an aqueous 20       medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs). MLVs generally have diameters ranging from 25  $\mu\text{m}$  to 4  $\mu\text{m}$ . Sonication of MLVs results in the formation of small unilamellar vesicles (SUVs) with diameters in the range of 200 to 500  $\text{\AA}$ , containing an aqueous solution in the core.

25       In addition to the teachings of Couvreur *et al.* (1991), the following information may be utilized in generating liposomal formulations. Phospholipids can form a variety of structures other than liposomes when dispersed in water, depending on the molar ratio of lipid to water. At low ratios the liposome is the preferred structure. The physical characteristics of liposomes depend on pH, ionic strength and 30       the presence of divalent cations. Liposomes can show low permeability to ionic and

polar substances, but at elevated temperatures undergo a phase transition which markedly alters their permeability. The phase transition involves a change from a closely packed, ordered structure, known as the gel state, to a loosely packed, less-ordered structure, known as the fluid state. This occurs at a characteristic phase-  
5 transition temperature and results in an increase in permeability to ions, sugars and drugs.

Liposomes interact with cells *via* four different mechanisms: Endocytosis by phagocytic cells of the reticuloendothelial system such as macrophages and neutrophils; adsorption to the cell surface, either by nonspecific weak hydrophobic or  
10 electrostatic forces, or by specific interactions with cell-surface components; fusion with the plasma cell membrane by insertion of the lipid bilayer of the liposome into the plasma membrane, with simultaneous release of liposomal contents into the cytoplasm; and by transfer of liposomal lipids to cellular or subcellular membranes, or *vice versa*, without any association of the liposome contents. It often is difficult to  
15 determine which mechanism is operative and more than one may operate at the same time.

#### **L. Pharmaceutical Compositions**

The pharmaceutical compositions disclosed herein may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be  
20 enclosed in hard or soft shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least  
25 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of the unit. The amount of active compounds in such therapeutically useful compositions is such that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following:  
30 a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the

like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

The active compounds may also be administered parenterally or intraperitoneally. Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium



chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

5 Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of  
10 sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and  
15 absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

20 The phrase "pharmaceutically acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human. The preparation of an aqueous composition that contains a protein as an active ingredient is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions; solid  
25 forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified.

The composition can be formulated in a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example,  
30 hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or

ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

## EXAMPLES

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

### Example 1

#### **Experimental Procedures**

##### **Bacterial strains**

5        *B. garinii* Ip90 was initially isolated from ticks collected in eastern Russia (Kriuchechnikov *et al.*, 1988). *B. afzelii* ACAI was cultured from a patient in Sweden with acrodermatitis chronica atrophicans (Asbrink *et al.*, 1984). Both strains were graciously provided by Dr. Alan Barbour, University of California at Irvine School of Medicine, and had been passed through C3H/HeN mice to assure infectivity. Strains  
10        were passaged in vitro fewer than 5 times following mouse infection.

##### **DNA cloning and sequencing**

Plasmid DNA was purified from the *Borrelia* strains as described previously (Purser and Norris, 2000).  $\lambda$  DASH II libraries of plasmid DNA fragments were  
15        prepared as described by Zhang *et al.* (Zhang *et al.*, 1997), with minor modifications. Thirty micrograms of plasmid DNA was treated with 30 units of mung bean nuclease at 30°C for 30 min to hydrolyze hairpin loops in telomeres, and an *EcoRI* linker (5'-CCGGAATTCCGG-3'; SEQ. ID. NO:107) was then ligated to the treated plasmid DNA using T<sub>4</sub> DNA ligase at 15°C overnight. This preparation was then digested to  
20        completion with *EcoRI*, and the resulting DNA fragments were fractionated by agarose gel electrophoresis. *EcoRI*-treated DNA fragments ranging in size from 8 kb to 25 kb were used to create libraries in *EcoRI* pre-treated  $\lambda$  DASH II vector arms as described in the manufacturer's instructions (Stratagene, La Jolla, CA, USA). Recombinant phages were screened by plaque hybridization using *B. burgdorferi* B31  
25        *vls* silent cassette clone pJRZ53 (Zhang *et al.*, 1997) as probe; hybridization with pJRZ53 was confirmed by secondary phage plaque screening as well as Southern blot hybridization. Selected phage clones were expanded, phage were purified, and DNA was prepared by standard techniques (Sambrook and Russell, 2000). The  $\lambda$  phage clones Ip90.1A1 and ACAI.2A1, each containing a 15 kb borrelia DNA insert, were  
30        selected for analysis.

To sequence the DNA insert of Ip90.1A1, the phage DNA was digested with *EcoRI* and *HindIII* and a 6kb *EcoRI/HindIII* fragment containing *vls*-like sequence was then cloned into pBluescript II SK(-) (Stratagene). The plasmid DNA of the

pBluescript clone was digested with *EcoRI* and *HindIII*, and the 6 kb DNA fragment was isolated by agarose gel electrophoresis followed by electroelution, partially digested with DNase I and cloned into *EcoRV* treated pBluescript II SK (-) to create random DNase I library as described previously (Zhang *et al.*, 1997). Clones with insert DNA ranging in size from 500 to 1,000 bp from the DNase I library were selected for sequencing using primers specific for the vector T7 and T3 sequences. To facilitate sequencing of the ACAI.2A1 clone, the phage DNA was treated with *XbaI* and *EcoRI*, and one 8kb *EcoRI/XbaI* fragment containing *vls*-like sequence was isolated from an agarose gel. This 8kb *EcoRI/XbaI* fragment was digested separately with *RsaI* and *PstI* and then cloned into pBluescript II SK (-) to generate *RsaI* and *PstI* libraries. Clones from both libraries were selected for sequencing at the Department of Microbiology and Molecular Genetics Sequencing Facility. Primer walking and PCR (see below) were utilized as needed to fill gaps, establish clone order, and confirm and extend the sequences. DNA sequences were assembled using DNASTAR software (DNASTAR, Inc., Madison, WI).

### Southern hybridization

Fifty nanograms of DNA was digested with the indicated restriction enzymes, subjected to agarose electrophoresis in 1X TAE buffer at 100V for 2 hr, and transferred to Amersham Hybond N<sup>+</sup> membranes using standard alkaline transfer techniques (Sambrook and Russell, 2000). Hybridization with pJRZ53 as probe was performed by enhanced chemiluminescence techniques following the manufacturer's protocol (Amersham Gene Images, Amersham, Piscataway, NJ, USA).

### PCR and RT-PCR

PCR was utilized to amplify *vls* sequences beyond the end of the 8kb *EcoRI/XbaI* fragment from ACAI, and thereby extend the sequence beyond the cloned region. The specific primer 4540 (5'-CCA GCA AAC AAC TTC CCC GCC-3' – SEQ ID NO:21), based on a variable region, and the nonspecific primer 4548 (5'- ATC CTT AAA CTC CGC CCC ATC ATC-3' – SEQ ID NO:22), based on an invariant 5' region of the *vls* silent cassettes of ACAI, were used as primers. Primer 4545 (5'- GAG TGC TGT GGA GAG TGC TGT TGA TGA G-3' – SEQ ID NO:23), based on

the direct repeat sequence, was also used in some PCR studies. *B. afzelii* ACAI plasmid DNA was used as the template in these reactions.

RT-PCR was used to detect transcription of *vlsE* in *B. garinii* Ip90 and *B. afzelii* ACAI. Forward primer 4587 (5'-GGG GAT AAA GGG GAT TGT TGAT GCT GC-3' – SEQ ID NO:24) and reverse primer 4588 (5'-GCA AAC TGC CCA TCC TTA GCC ATT CC-3' – SEQ ID NO:25) were designed based on the invariable regions of *vls* silent cassettes of Ip90; the forward primer 4470 (5'-AAG GGG ATT GCG AAG GGG ATA AAG G-3' – SEQ ID NO:26) and reverse primer 4471 (5'-TTA GCA GCA AACTTT CCA TCC TTA GCC-3' – SEQ ID NO:27) were used for ACAI. Total RNA was isolated from late log-phase cultures of Ip90 and ACAI using an RNA purification kit (Amersham). RT-PCR was carried out using the Promega Access RT-PCR kit according to manufacturer's instructions. Briefly, reverse transcription was carried out for 50 min at 48°C followed by an initial denaturation at 94°C for 3 min, and 30 cycles consisting of denaturation at 94°C for 30 sec, annealing at 68°C for 1.5 min, and extension at 68°C for 1.5 min.

#### Cloning and sequencing *vlsE* RT-PCR products

As mentioned above, both *B. afzelii* ACAI and *B. garinii* Ip90 used in these studies were first cloned by colony formation and then passaged through mice. To determine whether *vlsE* sequence variation was present following mouse infection, *B. afzelii* ACAI was grown from a frozen stock and cloned by colony formation on BSKY plates (Dever *et al.*, 1992). RT-PCR of individual clones was performed as described in a previous section, and cDNA was ligated into pCR 2.1 TOPO TA cloning vector (Invitrogen, Carlsbad, CA, USA). Each *vlsE* variant was sequenced with the M13 forward and reverse primers. *B. garinii* Ip90 RNA was isolated from an uncloned population following mouse infection, and thus contained a mixture of variants. RT-PCR and cDNA cloning were performed using the method described for ACAI. Sequences were aligned with the multiple alignment program (Smith *et al.*, 1996). The alignment output was formatted using Boxshade 3.21 (Hofmann and Baron, 1996).

### Accession numbers

The sequence of the *vls* silent cassette region of *B. afzelii* ACAI is provided at the United States National Center for Biomedical Information with GenBank accession number AY100628 (SEQ ID: NO:57). The *B. garinii* Ip90 silent cassette  
5 region is listed as AY100633 (SEQ ID NO:28). The RT-PCR product sequences obtained are listed as AY100629 – AY100632 (SEQ ID: NOS:5-12) and AY100634 – AY100637 (SEQ ID NOS:13-20) for ACAI and Ip90, respectively.

### Example 2

10

#### Identification of *vls* loci in *B. garinii* Ip90 and *B. afzelii* ACAI

Hybridization with the *B. burgdorferi* B31 *vls* silent cassette sequence in recombinant plasmid pJRZ53 was used as a means of identifying the plasmids and DNA fragments containing *vls* sequences in *B. garinii* Ip90 and *B. afzelii* ACAI. The  
15 pJRZ53 probe hybridized exclusively to plasmids with an approximate size of 28 kb in both ACAI and Ip90. Following treatment of plasmid preparations with restriction enzymes, the major hybridizing DNA segments were identified as a 15 kb *EcoRI* fragment of ACAI DNA and a 20kb *EcoRI* fragment of Ip90 plasmid DNA. Libraries  
20 of plasmid DNA *EcoRI* fragments were prepared in Lambda Dash II using a technique that permits the cloning of telomere-containing as well as internal fragments through treatment of the hairpin loop telomeres with mung bean nuclease followed by ligation with *EcoRI* linkers (Zhang *et al.*, 1997). The phage libraries were screened by hybridization with pJRZ53, and clones Ip90.1A1 and ACAI.2A1, each containing 15  
25 kb of insert DNA, were used for further analysis.

25

### Example 3

#### Organization of *vls* silent cassette loci

The overall organization of the *vls* silent cassette loci of Ip90 and ACAI is  
30 shown in FIG. 1. As was the case in *B. burgdorferi* B31, the silent cassette loci in each strain was composed of a contiguous array of multiple cassettes. The loci in Ip90 and ACAI consisted largely of contiguous, uninterrupted open reading frames, with

one frameshift present at the 3' end of cassette 9 in ACAI. The B31 *vls* silent cassette locus contained one stop codon and two frame shifts (Zhang *et al.*, 1997).

#### Example 4

5

##### **Structure of the Ip90 *vls* silent cassette locus**

In Ip90, the *vls* array consisted of 11 regions with homology to the *vls* cassettes of B31 (FIG. 1A). With the exception of the junctions at *vls3/4* and *vls6/7*, the 11 *vls* silent cassettes are flanked by 18 bp direct repeat sequences in the 6 kb region. However, several of these cassettes (*vls1*, *vls4*, *vls6*, and *vls11*) were truncated (189 to 288 bp in length) relative to the other, full-length cassettes ranging in size from 573 to 594 bp. By comparison with the *vls* expression cassette of B31, cassette 1 is truncated at the 3' region, containing only 92 amino acid codons; cassette 4 lacks 125 codons in its 5' region; cassette 6 contains only 89 codons and is missing most of the 3' region; and cassette 11 has 86 codons, but is missing the 3' region. A portion of the silent cassette locus from the last 3 bp of cassette 5 to the first 165bp of cassette 8 is identical to the P7-1 clone previously characterized by Liang *et al.* (Liang and Philipp, 1999) (FIG. 1A). The 3' end of the Ip90 silent cassette locus possessed a truncated pseudogene of a conserved hypothetical protein belonging to gene family 144 of *B. burgdorferi* B31 (TIGR, 2002).

20

The 5' end of the locus also contained a region homologous to the 5', unique (non-cassette) portion of B31 expression site, *vlsE* (FIG. 1A). However, this gene segment is lacking a promoter region and the first 59 codons of *vlsE*, and also contains segments that are non-homologous to B31 *vlsE*. Therefore, this '*vlsE*-like' sequence appears to be a pseudogene, although it is in frame with the cassette 1 of the *vls* silent cassette array and could conceivably encode a *vlsE*-like product. It is of interest to note that *vlsE* of *B. burgdorferi* B31 is located close to the telomere of lp28-1, but is oriented in the opposite direction (i.e. is transcribed toward the telomere) relative to the *vlsE*-like sequence of Ip90. In addition, the reading frame of the *vls* silent cassette array in Ip90 runs away from, rather than toward (as is the case with the silent cassettes in B31), the nearest telomere (FIG. 1) (Zhang *et al.*, 1997). Therefore, the B31 and Ip90 versions of the silent cassette loci have likely undergone large-scale rearrangements during evolution from a common ancestral organism, and it is unlikely

30

that the Ip90 *vlsE*-like pseudogene evolved directly from a functional telomeric copy of *vlsE*. Based on other evidence, we believe that a functional *vlsE* gene is located elsewhere on the 28 kb plasmid of Ip90 (see below).

Portions of several *vls* silent cassettes from *Borrelia garinii* strain A87S were published previously (Wang *et al.*, 2001). Each putative silent cassette in the longest available A87S sequence (GenBank Accession No. AF274070) was compared to its corresponding cassette among the Ip90 silent cassettes. The A87S sequence shared only 63 to 68% nucleotide identity to Ip90 sequences, and amino acid similarity ranged from 51 to 57%. An amino acid alignment between the A87S and Ip90 silent cassettes reveals that the heterogeneity exists largely within invariable region 1 (IR1), found upstream of VR-I (data not shown). There are also considerable differences in IR4 and IR6, but to a lesser extent when compared to IR1. The sequence differences between the *vls* silent cassettes sequences of Ip90 and A87S indicates that a considerable degree of heterogeneity exists among *vls* sequences within this species, as also appears to be the case with *Borrelia burgdorferi* strains (Iyer *et al.*, 2000).

An unusual feature of the Ip90 telomere region upstream of the *vls* cassettes is the presence of a set of 6 complete and 1 partial copies of a 41 bp direct repeat sequence. The telomere itself was identified by its location in the lambda clone insert next to the *EcoRI* linker used to clone mung bean nuclease-treated telomere regions. Because mung bean nuclease potentially could remove terminal nucleotides as well as disrupting the hairpin loop 5'-3' bond, it is not known whether this sequence represents the absolute end of the telomere sequence. The telomeric repeat sequences (TRS) begin 52 bp from the end of the telomere sequence, and are present as six 41-bp repeats (TRS-A through TRS-F) followed by a 32-bp truncated version of the 41-bp sequence (TRS-G) in a contiguous array. These direct repeats differ at only one position in TRS-B, and are otherwise identical. The telomeric direct repeat has no significant homology with *vls* sequences or any other *borrelia* sequence reported previously. Although the direct repeats obviously arose through duplication events, their origin and significance are unknown at this time.

30



### Example 5

#### **Structure of the ACAI *vls* silent cassette locus**

The overall arrangement of the *B. afzelii* ACAI *vls* silent cassette locus is depicted in FIG. 1B. Unlike Ip90 and B31, the ACAI *vls* locus was located on an internal *EcoRI* fragment of a 28-kb linear plasmid, and its location relative to the plasmid telomeres is not known. The ACAI *vls* locus contains 13 complete and 1 partial silent cassettes and each cassette is also flanked by an 18 bp direct repeat sequence. Twelve of the cassettes appear to represent 'full-length' sequences (ranging from 591 to 630 bp in length), whereas cassette 11 contains an internal deletion and cassette 14 has an internal deletion and a short, 3' truncation relative to the other cassette sequences (FIG. 1B). The 3' end of the silent cassette locus is demarcated by a complete copy of a conserved hypothetical protein gene belonging to gene family 57 of *B. burgdorferi* B31 (TIGR, 2002). We were unable to obtain additional sequence 5' of cassette 1, and it is possible that additional *vls* sequences are localized upstream of the region we have characterized thus far.

### Example 6

#### **Direct repeats in the silent cassette loci**

In *B. burgdorferi* B31, both the central cassette of *vlsE* and the homologous *vls* silent cassettes are flanked by a 17 bp direct repeat sequence (5'-TGAGGGGGCTATTAAGG-3' (SEQ ID NO:106)). This sequence is generally well-conserved in the *vlsE* expression site and the silent cassettes; it is absent from the 5'-truncated cassette 1, and only 10 of 17 nucleotides are present at the junction between *vls9* and *vls10* (Zhang *et al.*, 1997). Based on the location and high degree of conservation of the 17 bp direct repeat, it was hypothesized previously that these sequences may play an important role in the *vls* gene conversion process. However, the 17 bp sequence is not highly conserved in the *B. garinii* Ip90 and *B. afzelii* ACAI *vls* silent cassette sequences (data not shown). A comparison of 17 bp consensus sequences from Ip90 and ACAI to the B31 17 bp sequence shows that the Ip90 and ACAI sequences are more similar to each other than to the B31 sequence. Nevertheless, the higher degree of variability in the Ip90 and ACAI 17 bp sequences

compared to the B31 sequence suggests that the 17 bp sequence is not as important in the gene conversion process as previously thought (Zhang *et al.*, 1997).

### Example 7

5

#### **Similarity of *vls* silent cassette loci**

Alignment of the *vls* cassette sequences from Ip90, ACAI, and B31 indicates a high degree of sequence conservation both within and between each strain (FIG. 2). The Ip90 cassettes share 90 to 97% nucleotide sequence identity with one another, whereas the ACAI silent cassettes have from 84 to 91% nucleotide sequence identity (data not shown). The Ip90 *vls* silent cassettes are also highly homologous with *B. burgdorferi* *vls* sequences; for example, sequence identities with the B31 allele *vlsE1* (Zhang *et al.*, 1997) range from 64% to 73% on the nucleotide level and from 53% to 62% in predicted amino acid sequence (FIG. 2A). The identities between the ACAI *vls* silent cassettes and B31 *vlsE1* likewise range from 65% to 73% on the nucleotide level and from 50% to 65% in predicted amino acid sequence (FIG. 2B). Each complete silent cassette of Ip90 and ACAI contains six variable regions interspersed by six invariable regions similar to those found in the *vls* sequences of B31 (FIG. 2).

SEQ ID NO:28 is the *B. garinii* Ip90 *vls* locus silent cassette nucleic acid sequence. SEQ ID NO:30 is a translation of an upstream open reading frame of SEQ ID NO:28, which is contiguous with the open reading frame of the silent cassettes of the *B. garinii* Ip90 *vls* locus. SEQ ID NO:32 is a translation of a *vlsE*-like sequence of SEQ ID NO:28. SEQ ID NOS:33-54 are nucleotide and amino acid sequences of silent cassette Nos. 1-11 of the *B. garinii* Ip90 *vls* locus as set forth in FIG. 2B. SEQ ID NO:55 and 56 are the nucleotide and amino acid sequences of a truncated pseudogene in the *B. garinii* Ip90 *vls* locus with 85% similarity to amino acids 70-140 of the *B. burgdorferi* B31 ORF-10 predicted product, GenBank Accession No. AA 34908.

SEQ ID NO:57 is the *B. afzelii* ACAI *vls* silent cassette locus nucleic acid sequence. SEQ ID NOS:58-85 are the nucleotide and amino acid sequences of silent cassette Nos. 1-14 of the *B. afzelii* ACAI silent cassette locus as set forth in FIG. 2A. SEQ ID NOS:86 and 87 are the nucleotide and amino acid sequences of a portion of the *B. afzelii* ACAI *vls* silent cassette locus which encodes a member of protein family

PF02414, a conserved hypothetical protein family thought to be involved in *Borrelia* plasmid partitions of replication.

### Example 8

#### 5 **Transcription of *vlsE* of *B. garinii* Ip90 and *B. afzelii* ACAI**

We have thus far been unsuccessful in cloning a complete *vlsE* expression site from either Ip90 or ACAI using a variety of approaches (data not shown). To determine whether *vls* expression sites are present in Ip90 and ACAI, RT-PCR was carried out using total RNA from in vitro cultured *B. garinii* Ip90 and *B. afzelii* ACAI.

10 Primers corresponding to invariant regions in the *vls* silent cassette regions of each organism were utilized. We observed a positive RT-PCR result in ethidium bromide-stained agarose gels for both *B. garinii* Ip90 and *B. afzelii* ACAI, but no products were observed if reverse transcriptase was omitted in the RT reaction (FIG. 3). The RT-PCR products containing *vls*-like sequence were confirmed by sequencing,

15 confirming that both organisms have *vls* expression sites. In *B. burgdorferi* B31, *vlsE* is located only 160 bp from the *vls* silent cassette array (Hudson *et al.*, 2001; Zhang *et al.*, 1997). Based on our studies, the *vls* expression sites of ACAI and Ip90 do not appear to be located in close proximity to the *vls* silent cassettes.

20

### Example 9

#### **Sequence analysis of *vlsE* variants of *B. afzelii* ACAI and *B. garinii* Ip90**

Both ACAI and Ip90 were passaged through mice prior to analysis. In previous studies with *B. burgdorferi* B31, extensive sequence variation due to

25 apparent gene conversion events occurred within the *vlsE* cassette region during mouse infection (Zhang and Norris, 1998a, b). To determine whether similar sequence variation occurred in ACAI and Ip90, individual RT-PCR products from each mouse-passaged strain were cloned and sequenced.

An alignment of the predicted *VlsE* protein sequences of ACAI and Ip90

30 (FIG. 4) demonstrated that sequence variation occurred within each strain. Moreover, the changes observed were consistent with gene conversion involving segments of the

silent cassettes, as had been seen previously with B31. As with B31, the sequence differences were predictably localized primarily within the variable regions.

Using the sequences from the silent cassettes of each organism (FIG. 2), we determined the silent cassette sequences that were most likely involved in the gene conversion events within ACAI and Ip90 *plsE* genes (FIG. 4). The theoretical minimum and maximum recombination events are indicated by solid and dotted lines, respectively. In FIG. 4A, silent cassette amino acid sequences matching regions of each variant are noted for all ACAI *plsE* variants except clone 2622. The variation seen in clones 2624a and 2624b can be attributed to two silent cassettes each. In clone 2624a, *pls8* matched the sequence found in a portion of variable region I (VR-I) and the entire sequence within VR-II, while *pls7* matched the sequence found in VR-III, VR-IV, and VR-V. In clone 2624b, *pls10* matched the sequence found in a portion of VR-I and the entire sequence within VR-II and VR-III, while *pls12* matched the sequence found in VR-IV and VR-V. While both *pls5* and *pls6* match large portions of sequence in clone 2625, it seems more likely that *pls5* was exclusively involved in the gene conversion events leading to the variation seen in clone 2625 since it contains sequence identity to VR-II, VR-III, VR-IV, and VR-V. It was difficult to ascertain which silent cassettes most likely contributed to the variation seen in clone 2622. Most silent cassettes matches spanned only a few residues in clone 2622. The nature of the sequence in clone 2622 suggests that it may be an artifactual PCR product.

Minimal recombination regions, indicated by solid lines in FIG. 4, were defined as the range of a *plsE* RT-PCR product sequence that matched only a single silent cassette sequence. These commonly extend over several variable regions, as was also the case with *B. burgdorferi* B31 in previous studies (Zhang *et al.*, 1997). In some cases, there are two or more silent cassettes that contain the same sequence within the same range. Therefore, it is only possible to predict the most likely silent cassette sequences involved (Indest *et al.*, 2001). Maximum recombination regions commonly extend from a variable region and continue into the flanking invariant region of the corresponding matching silent cassette (FIG. 4). The extension of the maximum recombination region ends at the first position of sequence non-identity between the *plsE* sequence of the clone and the given silent cassette. The degree of variation appears to be less than observed previously with *B. burgdorferi* B31, but an

analysis of *vlsE* at different times during mammalian infection (Zhang and Norris, 1998b) is required to provide an accurate measure of the kinetics.

There are two instances of what we believe to be point mutations in the Ip90 clones (FIG. 4B). The first instance lies two residues upstream of VR-II in clone 21, where there is an arginine residue not encoded in the silent cassettes. We believe a point mutation was responsible for changing the AAG codon in the silent cassettes to AGG in clone 21. The second example of a possible point mutation is the lone threonine after VR-V in clone 20. All of the silent cassette sequences possess a GCT codon at that position, while ACT is present in clone 20.

In conclusion, our results verify previous indications that both *B. garinii* and *B. afzelii* contain plasmid-encoded *vls* silent cassette loci similar to those of *B. burgdorferi*. In addition, RT-PCR results indicate that a *vls* product is expressed by both species, and that sequence variation occurs and hence may contribute to antigenic variation. Taken together, these and previous findings confirm that the *vls* sequence variation system is a common feature of Lyme disease borrelia, and hence is likely to be important in the pathogenesis of these organisms.

### Example 10

#### **Reactivity of Sera from Human Lyme Disease Patients and Infected Mice with *Borrelia afzelii* Protein.**

A recombinant DNA vector comprising a nucleotide sequence encoding the predicted amino acid sequence of the *B. afzelii* ACA-I *vls* cassette 13 (SEQ ID NOs:96 and 97) has been constructed. Briefly, DNA containing the coding sequence of the cassette region was amplified using a two-step polymerase chain reaction (PCR) method. During the first amplification, specific primers flanking the *B. afzelii* ACA-I *vls* cassette 13 (5'-CGGAATTC<sup>ACTCGCCTTACTATTATC</sup>-3' (SEQ ID NO:98) and 5'-CGGGATCCGAGAGTGCTGTTGATGAGGTT-3' (SEQ ID NO:99)) were used with *B. afzelii* ACA-I DNA as template to amplify a fragment containing the desired cassette. Then a second PCR was performed using primers specific for the cassette region itself (5'-CGGGATCC<sup>AAGAGTGCTGTGGATGAGGCTAGCAAG</sup>-3' (SEQ ID NO:100) and 5'-TTCTGCAGCACACTCGCCTTACTATTATCATTAGC-3' (SEQ ID

NO:101)) and the purified product of the first reaction as the DNA template. The two primers contained *Bam*HI and *Pst*I sites, respectively (underlined); the PCR product was treated with these two enzymes and ligated into the expression vector pQE30 cut with the same two enzymes. The sequence of the insert was analyzed and found to be the correct sequence. The resulting recombinant plasmid, pBA-13-1 was used to transform *E. coli* cells, and expression was induced by incubation of a transformed *E. coli* clone to 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) for 3 hours. The *E. coli* cells were lysed by sonication and centrifuged to remove cellular debris. The recombinant, His6-tagged protein (VLS-BA13) was purified by liquid chromatography over a nickel affinity column, elution of bound protein with imidazole, and further purification using a heparin-Sepharose column. The purity of the protein was determined to be >90% by sodium-dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and the concentration determined by a Bradford protein assay.

The purified recombinant protein VLS-BA13 was tested for reactivity with antibodies from humans using a pool of sera from patients fulfilling CDC criteria for Lyme disease, acquired in the North Central United States. A pool of negative control sera was obtained from human blood donors in Houston, Texas. Enzyme-linked immunosorbent assays (ELISAs) were performed as described (Lawrenz *et al.*, 1999), except that protein and serum concentrations were varied to determine the optimal concentrations. As shown in FIG. 6, VLS-BA13 protein (50 nanograms per well) consistently yielded higher absorbance readings with the Lyme disease serum pool than with the normal serum pool, up to a serum dilution of 1:6400. Differences in absorbance between the two serum preparations (1:200 dilution) were observed with VLS-BA13 protein concentrations as low as 3.13 nanograms per well (FIG. 7). Very similar results were obtained with sera from mice infected experimentally with *Borrelia burgdorferi* and sera from uninfected mice (FIGs. C and D). Taken together, these results provide evidence that amino acid sequences corresponding to *B. afzelii* Vls protein sequences react in a specific and sensitive manner with serum antibodies from Lyme disease patients or from *B. burgdorferi* infected mice.

### Example 11

#### **Reactivity of Sera from Human Lyme Disease Patients and Infected Mice with *Borrelia garinii* Protein.**

A recombinant DNA vector comprising a nucleotide sequence encoding the predicted amino acid sequence of the *B. garinii* Ip90 *vls* cassette 10 (SEQ ID NOs:94 and 95) has been constructed. Briefly, DNA containing the coding sequence of the cassette region was amplified using a two-step polymerase chain reaction (PCR) method. During the first amplification, specific primers flanking the *B. garinii* Ip90 *vls* cassette 10 (5'-CGGGATCCGCTGTTGGGAGTYGCAAC-3' (SEQ ID NO:102) and 5'-AACTGCAGATTATCATGAGCAGCATCCTTC-3' (SEQ ID NO:103)) were used with *B. garinii* Ip90 DNA as template to amplify a fragment containing the desired cassette. Then a second PCR was performed using primers specific for the cassette region itself (5'- CGGGATCCAAGGGGACTGTTAAGAATGCTGTTG-3' (SEQ ID NO:104) and 5'-TTCTGCAGATGATTATCATGAGCAGCATCCTTCA-3'(SEQ ID NO:105)) and the purified product of the first reaction as the DNA template. The two primers contained *Bam*HI and *Pst*I sites, respectively (underlined); the PCR product was treated with these two enzymes and ligated into the expression vector pQE30 cut with the same two enzymes. The sequence of the insert was analyzed and found to be the correct sequence. The resulting recombinant plasmid, pBG-10-1 was used to transform *E. coli* cells, and expression was induced by incubation of a transformed *E. coli* clone to 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) for 3 hours. The *E. coli* cells were lysed by sonication and centrifuged to remove cellular debris. The recombinant, His6-tagged protein (VLS-BG10) was purified by liquid chromatography over a nickel affinity column, elution of bound protein with imidazole, and further purification using a heparin-Sepharose column. The purity of the protein was determined to be >90% by sodium-dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and the concentration determined by a Bradford protein assay.

The purified recombinant protein VLS-BG10 was tested for reactivity with antibodies from humans using a pool of sera from patients fulfilling CDC criteria for Lyme disease, acquired in the North Central United States. A pool of negative control

sera was obtained from human blood donors in Houston, Texas. Enzyme-linked immunosorbent assays (ELISAs) were performed as described (Lawrenz *et al.*, 1999), except that protein and serum concentrations were varied to determine the optimal concentrations. In the examples shown, the antigen (VLS-BG10) was used to coat the wells, and the measured parameter was the amount of antibody bound as determined by addition of either goat anti-human IgG (alkaline phosphatase conjugate) or goat anti-mouse IgG (alkaline phosphatase conjugate), followed by washing and addition of a suitable substrate. As shown in FIG. 10, VLS-BG10 protein (10 nanograms per well) consistently yielded higher absorbance readings with the Lyme disease serum pool than with the normal serum pool, up to a serum dilution of 1:6400. Differences in absorbance between the two serum preparations (1:200 dilution) were observed with VLS-BG10 protein concentrations as low as 0.031 micrograms per well (FIG. 11). Very similar results were obtained with sera from mice infected experimentally with *Borrelia burgdorferi* and sera from uninfected mice (FIGs. 12 and 13). Taken together, these results provide evidence that amino acid sequences corresponding to *B. garinii* Vls protein sequences react in a specific and sensitive manner with serum antibodies from Lyme disease patients or from *B. burgdorferi* infected mice.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.



### REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

5

U.S. Patent 3,791,932

U.S. Patent 3,949,064

U.S. Patent 4,174,384

U.S. Patent 4,196,265

10

U.S. Patent 4,518,584

U.S. Patent 4,554,101

U.S. Patent 4,578,770

U.S. Patent 4,596,792

U.S. Patent 4,599,230

15

U.S. Patent 4,599,231

U.S. Patent 4,601,903

U.S. Patent 4,608,251

U.S. Patent 4,683,195

U.S. Patent 4,683,202

20

U.S. Patent 5,155,022

U.S. Patent 5,168,050

U.S. Patent 5,178,859

U.S. Patent 5,187,065

U.S. Patent 5,217,872

25

U.S. Patent 5,246,844

U.S. Patent 5,279,938

U.S. Patent 5,283,175

U.S. Patent 5,324,630

U.S. Patent 5,385,826

30

U.S. Patent 5,403,718

U.S. Patent 5,434,077

U.S. Patent 5,436,000

U.S. Patent 6,437,116

- Altschul, Gish, Miller, Myers, Lipman, "Basic local alignment search tool," *J. Mol. Biol.*, 215:403-410, 1990.
- 5 Asbrink, Hederstedt, Hovmark, "The spirochetal etiology of erythema chronicum migrans *Afzelius*," *Acta Derm. Venerol.*, 64:291-295, 1984.
- Balmelii and Piffatetti, "Analysis of the genetic polymorphism of *Borrelia burgdorferi* sensu lato by multilocus enzyme electrophoresis," *Int. J. Syst. Bacteriol.*, 46:167-172, 1996.
- 10 Barbour, "Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent," *J. Clin. Microbiol.*, 42:475-478, 1988.
- Barbour, "Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent," *J. Clin. Microbiol.*, 26:475-478, 1988.
- Barbour, "Linear DNA of *Borrelia* species and antigenic variation," *Trends*  
15 *Microbiol.*, 1:236-239, 1993.
- Barbour and Garon, "Linear plasmids of the bacterium *Borrelia burgdorferi* have covalently closed ends," *Science*, 237:409-411, 1987.
- Barbour, Burman, Carter, Kitten, Bergstrom, "Variable antigen genes of the relapsing fever agent *Borrelia hermsii* are activated by promoter addition," *Mol.*  
20 *Microbiol.*, 5:489-493, 1991a.
- Barbour, Carter, Burman, Freitag, Garon, Bergstrom, "Tandem insertion sequence-like elements define the expression site for variable antigen genes of *Borrelia hermsii*," *Infect. Immun.*, 59:390-397, 1991b.
- Barbour *et al.*, "Structural analysis of the variable major proteins of *Borrelia hermsii*,"  
25 *J. Exp. Med.*, 158:2127-2140, 1983.
- Barbour *et al.*, "Variable major proteins of *Borrelia hermsii*," *J. Exp. Med.*, 156:1312-1324, 1982.
- Barstad *et al.*, "Variable major proteins of *Borrelia hermsii*. Epitope mapping and partial sequence analysis of CNBr peptides," *J. Exp. Med.*, 161:1302-1314,  
30 1985.
- Barthold, "Antigenic stability of *Borrelia burgdorferi* during chronic infections of immunocompetent mice," *Infect. Immun.*, 61:4955-4961, 1993.

- Barthold, Moody, Beck, "Susceptibility of laboratory rats to isolates of *Borrelia burgdorferi* from different geographic areas," *Am. J. Trop. Med. Hyg.*, 42:596-600, 1990.
- 5 Borst and Geaves, "Programmed gene rearrangements altering gene expression," *Science*, 235:658-667, 1987.
- Borst, Bitter, McCulloch, Leeuwen, Rudenko, "Antigenic variation in malaria," *Cell*, 82:104, 1995.
- Burgdorfer, Barbour, Hayes, Benach, Grunwaldt, Davis, "Lyme disease, a tick-borne spirochetosis?," *Science*, 216: 1317-1319, 1982.
- 10 Carroll and Gheradini, "Membrane protein variations associated with *in vitro* passage of *Borrelia burgdorferi*," *Infect. Immun.*, 64:392-398, 1996.
- Carter *et al.*, "A family of surface-exposed proteins of 20 kilodaltons in the genus *Borrelia*," *Infect. Immun.*, 62:2792-2799, 1994.
- Casjens, Delange III, Ley, Rosa, Huang, "Linear chromosomes of Lyme disease agent spirochetes: genetic diversity and conservation of gene order," *J. Bacteriol.*, 15 177:2769-2780, 1995.
- Demolis, Mallet, Bussereau, Jacquet, "Improved strategy for large-scale DNA sequencing using DNase I cleavage for generating random subclones," *Biotechniques*, 18:453-457, 1995.
- 20 Dever, Jorgensen, Barbour, "In vitro antimicrobial susceptibility testing of *Borrelia burgdorferi*: a microdilution MIC method and time-kill studies," *J. Clin. Microbiol.*, 30:2692-2697, 1992.
- Donelson, "Mechanisms of antigenic variation in *Borrelia hermsii* and African trypanosomes," *J. Biol. Chem.*, 270:7783-7786, 1995.
- 25 Fuchs, Jauris, Lottspeich, Preacmursic, Wilskie, Soutschek, "Molecular analysis and expression of a *Borrelia burgdorferi* gene encoding a 22 kDa protein (pC) in *E. coli*," *Mol. Microbiol.*, 6:503-509, 1992.
- Haas and Meyer, "The repertoire of silent pilus genes in *Neisseria gonorrhoeae*: evidence for gene conversion," *Cell*, 44:107-115, 1986.
- 30 Hagblom, Segal, Billyard, So, "Intragenic recombination leads to pilus antigenic variation in *Neisseria gonorrhoeae*," *Nature*, 315:156-158, 1985.

- Hinnebusch, Bergstrom, Barbour, "Cloning and sequence analysis of linear plasmid telomeres of the bacterium *Borrelia burgdorferi*," *Mol. Microbiol.*, 4:811-820, 1990.
- Hofman and Baron, "Boxshade 3.21 [WWW Document},  
5 [http://www.isrec.isbsib.ch:8080/software/BOX\\_form.html](http://www.isrec.isbsib.ch:8080/software/BOX_form.html), 1996.
- Hudson, Frye, Quinn, Gherardini, "Increased expression of *Borrelia burgdorferi* vlsE in response to human endothelial cell membrane, *Mol. Microbiol.*, 41:229-239, 2001.
- Hughes and Johnson, "Methylated DNA in *Borrelia* species," *J. Bacteriol.*, 172:6602-  
10 6604, 1990.
- Indest, Howell, Jacobs, School-Meker, Norris, Phillipp, "Analysis of *Borrelia burgdorferi* vlsE gene expression and recombination in the tick vector," *Infect. Immun.*, 69:7083-7090, 2001.
- Johnson *et al.*, "Infection of Syrian hamsters with Lyme disease spirochetes," *J. Clin.*  
15 *Microbiol.*, 20:1099-1101, 1984.
- Jonsson, Ilver, Falk, Pepose, Normark, "Sequence changes in the pilus subunit lead to tropism variation of *Neisseria gonorrhoeae* to human tissue," *Mol. Microbiol.*, 13:403-416, 1994.
- Kitten and Barbour, "Juxtaposition of expressed variable antigen genes with a  
20 conserved telomere in the bacterium *Borrelia hermsii*," *Proc. Natl. Acad. Sci. USA*, 87:6077-6081, 1990.
- Kitten and Barbour, "The relapsing fever agent *Borrelia hermsii* has multiple copies of its chromosome and linear plasmids," *Genetics*, 132:311-324, 1992.
- Koomey, Gotschlich, Robbins, Bergstrom, Swanson, "Effects of *recA* mutations on  
25 pilus antigenic variation and phase transitions in *Neisseria gonorrhoeae*," *Genetics*, 117:391-398, 1987.
- Kriuchechnikov, Korenberg, Shcherbakov, Kovalevskii, Levin, "Identification of borrelia isolated in the USSR from *Ixodes persulcatus* Schulze ticks], *Zh Mikrobiol. Epidemiol. Immunobiol.*, 12:41-44, 1988.
- 30 Kupsch, Knepper, Kuroki, Heuer, Meyer, "Variable opacity (Opa) outer membrane proteins account for the cell tropisms displayed by *Neisseria gonorrhoeae* for human leukocytes and epithelial cells," *EMBO. J.*, 12:641-650, 1993.

- Lambden, Robertson, Watt, "Biological properties of two distinct pilus types produced by isogenic variants of *Neisseria gonorrhoeae* P9," *J. Bacteriol.*, 141:393-396, 1980.
- 5 Liang and Philipp, "Analysis of antibody response to invariable regions of VlsE, the variable surface antigen of *Borrelia burgdorferi*," *Infect. Immun.*, 67:6702-6706, 1999.
- Liang, Alvarez, Gu, Nowling, Ramamoorthy, Philipp, "An immunodominant conserved region within the variable domain of VlsE, the variable surface antigen of *Borrelia burgdorferi*," *J. Immunol.*, 163:5566-5573, 1999a..
- 10 Liang, Aberer, Cinco, Gern, Hu, Lobet, Ruscio, Voet, Jr., Weynants, Philipp, "Antigenic conservation of an immunodominant invariable region of the VlsE lipoprotein among European pathogenic genospecies of *Borrelia burgdorferi* SL," *J. Infect. Dis.*, 182:1455-1462, 2000a..
- Livey, Gibbs, Schuster, Dorner, "Evidence for lateral transfer and recombination in OspC variation in Lyme disease *Borrelia*," *Mol. Microbiol.*, 18:257-269, 1995.
- 15 Marconi, Konkell, Garon, "Variability of *osp* genes and gene products among species of Lyme disease spirochetes," *Infect. Immun.*, 61:2611-2617, 1993.
- Marconi, Samuels, Landry, Garon, "Analysis of the distribution and molecular heterogeneity of the *ospD* gene among the Lyme disease spirochetes: evidence
- 20 for lateral gene exchange," *J. Bacteriol.*, 176:4572-4582, 1994.
- Margolis *et al.*, "Homology between *Borrelia burgdorferi* OspC and members of the family of *Borrelia hermsii* variable major proteins," *Gene*, 143:105-110, 1994.
- Meier, Simon, Barbour, "Antigenic variation is associated with DNA rearrangements in a relapsing fever *Borrelia*," *Cell*, 41:403-409, 1985.
- 25 Meyer, Mlawer, So, "Pilus expression in *Neisseria gonorrhoeae* involves chromosomal rearrangement," *Cell*, 30:45-52, 1982.
- Moody *et al.*, "Lyme borreliosis in laboratory animals: effect of host species and *in vitro* passage of *Borrelia burgdorferi*," *Am. J. Trop. Med. Hyg.*, 43:87-92, 1990.
- 30 Nassif, Lowry, Stenberg, O'Gaora, Ganji, So, "Antigenic variation of pilin regulates adhesion of *Neisseria meningitidis* to human epithelial cells," *Mol. Microbiol.*, 8:719-725, 1993.

- Norris, Carter, Howell, Barbour, "Low-passage-associated proteins of *Borrelia burgdorferi* B31: characterization and molecular cloning of OspD, a surface-exposed, plasmid-encoded lipoprotein," *Infect. Immun.*, 60:4662-4672, 1992.
- Norris *et al.*, "High- and low-infectivity phenotypes of clonal populations of *in vitro*-cultured *Borrelia burgdorferi*," *Infect. Immun.*, 63:2206-2212, 1995.
- Norris *et al.*, "Low-passage-associated proteins of *Borrelia burgdorferi* B31: characterization and molecular cloning of OspD, a surface exposed, plasmid-encoded lipoprotein," *Infect. Immun.*, 60:4662-4672, 1992.
- Persing, Mathiesen, Podzorski, Barthold, "Genetic stability of *Borrelia burgdorferi* recovered from chronically infected immunocompetent mice," *Infect. Immun.*, 62:3521-3527, 1994.
- Plasterk *et al.*, "Transposition of structural genes to an expression sequence on a linear plasmid causes antigenic variation in the bacterium *Borrelia hermsii*," *Nature*, 318:257-263, 1985.
- Purser and Norris, "Correlation between plasmid content and infectivity in *Borrelia burgdorferi*," *Proc. Natl. Acad. Sci. USA*, 97:13865-13870, 2000.
- Restrepo and Barbour, "Antigen diversity in the bacterium *B. hermsii* through 'somatic' mutations in rearranged *vmp* genes," *Cell*, 78:867-876, 1994.
- Restrepo, Carter, Barbour, "Activation of a *vmp* pseudogene in *Borrelia hermsii*: an alternate mechanism of antigenic variation during relapsing fever," *Mol. Microbiol.*, 13:287-299, 1994.
- Restrepo, Kitten, Carter, Infante, Barbour, "Subtelomeric expression regions of *Borrelia hermsii* linear plasmids are highly polymorphic," *Mol. Microbiol.*, 6:3299-3311, 1992.
- Robertson and Meyer, "Genetic variation in pathogenic bacteria," *Trends Genet.*, 8:422-427, 1992.
- Rosa, Samuels, Hogan, Stevenson, Casjens, Tilly, "Directed insertion of a selectable marker into a circular plasmid of *Borrelia burgdorferi*," *J. Bacteriol.*, 178:5946-5953, 1996.
- Rosa, Schwan, Hogen, "Recombination between genes encoding major surface proteins A and B of *Borrelia burgdorferi*," *Mol. Microbiol.*, 6:3031-3040, 1992.

- Rudel, Van Putten, Gibbs, Haas, Meyer, "Interaction of two variable proteins (PilE and PilC) required for pilus-mediated adherence of *Neisseria gonorrhoeae* to human epithelial cells," *Mol. Microbiol.*, 6:3439-3450, 1992.
- 5 Sadziene, Rosa, Thompson, Hogan, Barbour, "Antibody-resistant mutations of *Borrelia burgdorferi*: *in vitro* selection and characterization," *J. Exp. Med.*, 176:799-809, 1992.
- Sambrook, Fritsch, Maniatis, "Molecular cloning: a laboratory manual," Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, 1989.
- 10 Sambrook, Fritsch, Maniatis, "Molecular cloning: a laboratory manual," Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, 2001.
- Samuels, Mach, Garon, "Genetic transformation of the Lyme disease agent *Borrelia burgdorferi* with coumarin-resistant *gyrB*," *J. Bacteriol.*, 176:6045-6049, 1994.
- 15 Schutzer, "Lyme disease: Molecular and immunologic approaches. In: *Current communications in cell and molecular biology*," J. Inglis and J.A. Witkowski, eds. (Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press).
- Schwann *et al.*, "Changes in infectivity and plasmid profile of the Lyme disease spirochete, *Borrelia burgdorferi*, as a result of *in vitro* cultivation," *Infect. Immun.*, 56:1831-1836, 1988a.
- 20 Schwann, Burgdorfer, Schrumppg, Karstens, "The urinary bladder, a consistent source of *Borrelia burgdorferi* in experimentally infected white-footed mice (*Peromyscus leucopus*)," *J. Clin. Microbiol.*, 26:893-895, 1988b.
- Schwann, Karstens, Schrumppf, Simpson, "Changes in antigenic reactivity of *Borrelia burgdorferi*, the Lyme disease spirochete, during persistent infection of mice," *Can. J. Microbiol.*, 37:450-454, 1991.
- 25 Seal, Jackson, Daniels, "Isolation of a *Pseudomonas solanacearum*-specific DNA probe by subtraction hybridization and construction of species-specific oligonucleotide primers for sensitive detection by the polymerase chain reaction," *Appl. Environ. Microbiol.*, 58:3751-3758, 1992.
- 30 Segal, Hagblom, Seifert, So, "Antigenic variation of gonococcal pilus involves assembly of separated silent gene segments," *Proc. Natl. Acad. Sci. USA*, 83:2177-2181, 1986.

- Seifert and So, "Genetic mechanisms of bacterial antigenic variation," *Microbiol. Rev.*, 52:327-336, 1988.
- Smith, Wiese, Wojzynski, Davison, Worley, "BCM Search Launcher—an integrated interface to molecular biology data base search and analysis services available on the World Wide Web, *Genome Res.*, 6:454-462, 1996.
- Steere, "Lyme disease," *N. Engl. J. Med.*, 321:586-596, 1989.
- Stevenson, Bockenstedt, Barthold, "Expression and gene sequence of outer surface protein C of *Borrelia burgdorferi* reisolated from chronically infected mice," *Infect. Immun.*, 62:3568-3571, 1994.
- Stoenner, Dodd, Larsen, "Antigenic variation of *Borrelia hermsii*," *J. Exp. Med.*, 156:1297-1311, 1982.
- Swanson and Koomey, "Mechanisms for variation of pili and outer membrane protein II in *Neisseria gonorrhoeae*," D.E. Berg and M.M. Howe, Eds. (Washington, D.C.: American Society for Microbiology).
- Thiessen *et al.*, "Evolution of the *Borrelia burgdorferi* outer surface protein OspC," *J. Bacteriol.*, 177:3036-3044, 1995.
- TIGR, The Institute for Genomic Research ([www.tigr.org](http://www.tigr.org)), 2002.
- Wainwright, Pritchard, Seifert, "A conserved DNA sequence is required for efficient gonococcal pilin antigenic variation," *Mol. Microbiol.*, 13:75-87, 1994.
- Walker, Howell, You, Hoffmaster, Heath, Weinstock, Norris, "Physical map of the genome of *Treponema pallidum* subsp. *Pallidum* (Nichols)," *J. Bacteriol.*, 177:1797-1804, 1995.
- Wang, van Dam, Dankert, "Analysis of a VMP-like sequence (vls) locus in *Borrelia garinii* and Vls homologues among four *Borrelia burgdorferi* sensu lato species," *FEMS Microbiol. Lett.*, 3199:9-45, 2001.
- Wilske, Barbour, Bergstrom, Burman, Restrepo, Rosa, Schwan, Soutschek, Wallich, "Antigenic variation and strain heterogeneity in *Borrelia* spp," *Res. Microbiol.*, 143:583-596, 1992.
- Wu and Tokunaga, "Biogenesis of lipoproteins in bacteria," *Curr. Top. Microbiol. Immunol.*, 125:127-157, 1986.
- Xu, Kodner, Coleman, Johnson, "Correlation of plasmids with infectivity of *Borrelia burgdorferi* sensu stricto type strain B31," *Infect. Immun.*, 64:3870-3876, 1996.



- Xu and Johnson, "Analysis and comparison of plasmid profile of *Borrelia burgdorferi* sensu lato strains," *J. Clin. Microbiol.*, 33:2679-2685, 1995.
- Zhang, Hardman, Barbour, Norris, "Antigenic variation in Lyme disease borreliae by promiscuous recombination of VMP-like sequence cassettes," *Cell*, 89:275-285, 1997.
- 5 Zhang and Norris, "Genetic variation of the *Borrelia burgdorferi* gene *vsIE* involves cassette-specific, segmental gene conversation," *Infect. Immun.*, 66:3698-3704, 1998a.
- 10 Zhang and Norris, "Kinetics and *in vivo* induction of genetic variation of *vsIE* in *Borrelia burgdorferi*, *Infect. Immun.*, 66:3689-3697, 1998b.

## CLAIMS

We claim:

- 5 1. An isolated nucleic acid comprising a nucleotide sequence that encodes a *vls* peptide of *Borrelia garinii* or *Borrelia afzelii*.
2. The nucleic acid of claim 1 further defined as comprising a nucleotide sequence that encodes at least 16 contiguous amino acids of SEQ ID NO:6, SEQ ID  
10 NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.
- 15 3. The nucleic acid of claim 1 further defined as comprising a nucleotide sequence that encodes at least 20 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID  
20 NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.
4. The nucleic acid of claim 1 further defined as comprising a nucleotide sequence that encodes at least 35 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID  
25 NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID  
30 NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

5. The nucleic acid of claim 1 further defined as comprising a nucleotide sequence that encodes at least 50 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

10

6. The nucleic acid of claim 1 further defined as comprising a nucleotide sequence that encodes a peptide comprising SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

20

7. The nucleic acid of claim 1 further defined as an RNA segment.

8. The nucleic acid of claim 1 further defined as comprising at least 50 contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96.

9. The nucleic acid of claim 1 further defined as comprising at least 75 contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96.
10. The isolated nucleic acid of claim 1 further defined as comprising at least 103 contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96.
11. The isolated nucleic acid of claim 1 further defined as comprising at least 110 contiguous nucleotides of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96.

12. The isolated nucleic acid of claim 1 further defined as comprising the nucleotide sequence of SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:37, SEQ ID NO:39, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:92, SEQ ID NO:94, and SEQ ID NO:96.
13. An isolated nucleic acid obtained by the amplification of a *Borrelia* nucleic acid with a primer selected from the group consisting of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:107.
14. A recombinant host cell comprising a heterologous nucleic acid comprising a nucleotide sequence that encodes a *vls* peptide of *Borrelia garinii* or *Borrelia afzelii*.
15. The recombinant host cell of claim 14, further defined as an *E. coli* cell.
16. A method of using an isolated nucleic acid that encodes a *vls* polypeptide of *Borrelia garinii* or *Borrelia afzelii*, comprising the steps of:
- (a) preparing a recombinant vector in which the isolated nucleic acid is positioned under the control of a promoter;
  - (b) introducing said recombinant vector into a host cell;
  - (c) culturing said host cell under conditions effective to allow expression of the polypeptide; and
  - (d) collecting said expressed polypeptide.

17. An isolated polypeptide comprising a Vls polypeptide of *Borrelia garinii* or *Borrelia afzelii*.
18. The isolated polypeptide of claim 17 further defined as comprising at least 16  
5 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID  
NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID  
NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID  
NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID  
NO:87, and SEQ ID NO:97.
- 10
19. The isolated polypeptide of claim 17 further defined as comprising at least 20  
contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID  
NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID  
NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID  
15 NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID  
NO:87, and SEQ ID NO:97.
20. The isolated polypeptide of claim 17 further defined as comprising at least 35  
contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID  
20 NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID  
NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID  
NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID  
NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID  
NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID  
25 NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID  
NO:93, SEQ ID NO:95, and SEQ ID NO:97.
21. The isolated polypeptide of claim 17 further defined as comprising at least 50  
contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID  
30 NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID  
NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID  
NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID  
NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID

NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

- 5 22. The isolated polypeptide of claim 17 further defined as comprising the amino acid sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.
- 10
- 15 23. An isolated polypeptide which specifically binds with antibodies raised against a Vls polypeptide of *Borrelia garinii* or *Borrelia afzelii*.
24. The polypeptide of claim 23 further defined as having at least 75% homology to a *Borrelia garinii* or *Borrelia afzelii* Vls polypeptide.
- 20
25. The polypeptide of claim 23 further defined as having at least 75% homology to a polypeptide comprising the amino acid sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.
- 25
- 30
26. The polypeptide of claim 23 further defined as having at least 85% homology to a polypeptide comprising the amino acid sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18,

SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36,  
SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56,  
SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67,  
SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77,  
5 SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87,  
SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

27. The polypeptide of claim 23 further defined as having at least 90% homology  
to a polypeptide comprising the amino acid sequence of SEQ ID NO:6, SEQ ID NO:8,  
10 SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18,  
SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36,  
SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56,  
SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67,  
SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77,  
15 SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87,  
SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

28. A protein composition comprising an isolated polypeptide which specifically  
binds with antibodies raised against a Vls polypeptide of *Borrelia garinii* or *Borrelia*  
20 *afzelii*.

29. The composition of claim 28 further defined as comprising a physiologically  
acceptable excipient.

25 30. The composition of claim 28, wherein the isolated polypeptide further  
comprises an anchoring moiety.

31. The composition of claim 30, wherein the anchoring moiety is biotin.

30 32. A purified antibody that binds immunologically to a Vls polypeptide of  
*Borrelia garinii* or *Borrelia afzelii*.



33. The purified antibody of claim 32, wherein the Vls polypeptide is further defined as comprising at least 16 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

34. The purified antibody of claim 32, wherein the Vls polypeptide is further defined as comprising at least 20 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

35. The purified antibody of claim 32, wherein the Vls polypeptide is further defined as comprising at least 35 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

36. The purified antibody of claim 32, wherein the Vls polypeptide is further defined as comprising at least 50 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

37. The purified antibody of claim 32, wherein the Vls polypeptide is further defined as comprising the amino acid sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

38. The antibody of claim 32 wherein the antibody is linked to a detectable label.

39. A method of generating an immune response, comprising administering to a mammal an immunologically effective amount of a Vls polypeptide of *Borrelia garinii* or *Borrelia afzelii*.

40. The method of claim 39, wherein the Vls polypeptide comprises at least 16 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

41. The method of claim 39, wherein the Vls polypeptide comprises at least 20 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

42. The method of claim 39, wherein the Vls polypeptide comprises at least 35 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID

NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

43. The method of claim 39, wherein the Vls polypeptide comprises at least 50 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

44. The method of claim 39, wherein the Vls polypeptide comprises SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

45. A method of assaying for *Borrelia* infection comprising:

- (a) obtaining a sample from a subject;
- (b) obtaining an antibody that binds immunologically to a Vls polypeptide of *Borrelia afzelii* or *Borrelia garinii* or a Vls polypeptide that binds immunologically to such an antibody;

- (c) admixing the sample and the antibody or the Vls polypeptide; and  
(d) determining whether immunologic binding occurs between the antibody and a polypeptide in the sample or between the Vls polypeptide and an antibody in the sample;  
5 wherein immunologic binding is indicative of *Borrelia* infection.

46. The method of claim 45, wherein the *Borrelia* infection is further defined as Lyme disease.

10 47. The method of claim 45, wherein the sample is blood.

48. The method of claim 45, wherein the sample is urine.

15 49. The method of claim 45, wherein the assay is an immunoassay.

50. The method of claim 49, wherein the immunoassay is an enzyme immunoassay.

20 51. The method of claim 50, wherein the enzyme immunoassay comprises an ELISA.

52. The method of claim 49, wherein the immunoassay assay comprises western blotting.

25 53. The method of claim 49, wherein the immunoassay comprises immunoprecipitation.

54. The method of claim 49, wherein the immunoassay comprises a radioimmunoassay.

30 55. The method of claim 45, wherein the subject is an animal.

56. The method of claim 55, wherein the animal is a human.

57. The method of claim 55, wherein the animal is a dog, deer, horse, cow or mouse.

5 58. The method of claim 45, wherein the Vls polypeptide comprises at least 16 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID  
10 NO:87, and SEQ ID NO:97.

59. The method of claim 45, wherein the Vls polypeptide comprises at least 20 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID  
15 NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

60. The method of claim 45, wherein the Vls polypeptide comprises at least 35 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID  
25 NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

61. The method of claim 45, wherein the Vls polypeptide comprises at least 50 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID  
30

NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

5

62. The method of claim 45, wherein the Vls polypeptide comprises SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

15

63. The method of claim 45, wherein the Vls polypeptide further comprises an anchoring moiety.

20

64. The method of claim 63, wherein the anchoring moiety is biotin.

65. The method of claim 45 further defined as comprising:  
(a) obtaining an antibody that binds immunologically to a Vls polypeptide of *Borrelia garinii* or *Borrelia afzelii*;  
(b) admixing the sample and the antibody; and  
(c) determining whether there is immunologic binding between the antibody and a polypeptide in the sample.

25

66. The method of claim 45 further defined as comprising:  
(a) obtaining a polypeptide that binds immunologically to an antibody that binds immunologically to a Vls polypeptide of *Borrelia garinii* or *Borrelia afzelii*;  
(b) admixing the sample and the polypeptide; and

30

- (c) determining whether the polypeptide binds immunologically to an antibody in the sample.

67. The method of claim 66, wherein determining immunological binding  
5 comprises an ELISA.

68. A method of assaying for *Borrelia* infection comprising:
- (a) obtaining a sample from a subject;
  - (b) obtaining a Vls polypeptide of *Borrelia afzelii* or *Borrelia garinii*;
  - 10 (c) admixing the sample and the the polypeptide; and
  - (d) determining whether immunologic binding occurs between the polypeptide and an antibody in the sample;

wherein immunologic binding is indicative of *Borrelia* infection.

15 69. The method of claim 68, wherein the *Borrelia* infection is further defined as Lyme disease.

70. The method of claim 68, wherein the sample is blood.

20 71. The method of claim 68, wherein the sample is urine.

72. The method of claim 68, wherein the assay is an immunoassay.

73. The method of claim 72, wherein the immunoassay is an enzyme  
25 immunoassay.

74. The method of claim 73, wherein the enzyme immunoassay comprises an ELISA.

30 75. The method of claim 72, wherein the immunoassay assay comprises western blotting.

76. The method of claim 72, wherein the immunoassay comprises immunoprecipitation.

77. The method of claim 72, wherein the immunoassay comprises a  
5 radioimmunoassay.

78. The method of claim 68, wherein the subject is an animal.

79. The method of claim 78, wherein the animal is a human.

10

80. The method of claim 78, wherein the animal is a dog, deer, horse, cow or mouse.

81. The method of claim 68, wherein the Vls polypeptide comprises at least 16  
15 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

20

82. The method of claim 68, wherein the Vls polypeptide comprises at least 20  
contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID  
25 NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, and SEQ ID NO:97.

83. The method of claim 68, wherein the Vls polypeptide comprises at least 35  
contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID  
30 NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID



NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

5      84.      The method of claim 68, wherein the Vls polypeptide comprises at least 50 contiguous amino acids of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID  
10      NO:61, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

15      85.      The method of claim 68, wherein the Vls polypeptide comprises SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:63, SEQ ID  
20      NO:65, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:95, and SEQ ID NO:97.

25      86.      The method of claim 68, wherein the Vls polypeptide further comprises an anchoring moiety.

87.      The method of claim 86, wherein the anchoring moiety is biotin.

30      88.      A kit for use in diagnosing Lyme disease in a subject comprising:

- (a) a first container including an antibody that binds immunologically to a Vls polypeptide of *Borrelia afzelii* or *Borrelia garinii* or a Vls polypeptide that binds immunologically to such an antibody.

89. The kit of claim 88, wherein the antibody is a monoclonal antibody.
90. The kit of claim 88, wherein the antibody is a polyclonal antibody.
- 5 91. The kit of claim 88, wherein the antibody is labeled.
92. The kit of claim 88, wherein the polypeptide is labeled.
- 10 93. The kit of claim 88, wherein the Vls polypeptide further comprises an anchoring moiety.
94. The composition of claim 93, wherein the anchoring moiety is biotin.

15

10/539956




FIG. 1A




FIG. 1B




FIG. 2A

[illegible]

TABLE	NAME	AGE
1	JOHN	25
2	MARY	22
3	JOHN	25
4	MARY	22
5	JOHN	25
6	MARY	22
7	JOHN	25
8	MARY	22
9	JOHN	25
10	MARY	22
11	JOHN	25
12	MARY	22
13	JOHN	25
14	MARY	22
15	JOHN	25
16	MARY	22
17	JOHN	25
18	MARY	22
19	JOHN	25
20	MARY	22
21	JOHN	25
22	MARY	22
23	JOHN	25
24	MARY	22
25	JOHN	25
26	MARY	22
27	JOHN	25
28	MARY	22
29	JOHN	25
30	MARY	22
31	JOHN	25
32	MARY	22
33	JOHN	25
34	MARY	22
35	JOHN	25
36	MARY	22
37	JOHN	25
38	MARY	22
39	JOHN	25
40	MARY	22
41	JOHN	25
42	MARY	22
43	JOHN	25
44	MARY	22
45	JOHN	25
46	MARY	22
47	JOHN	25
48	MARY	22
49	JOHN	25
50	MARY	22
51	JOHN	25
52	MARY	22
53	JOHN	25
54	MARY	22
55	JOHN	25
56	MARY	22
57	JOHN	25
58	MARY	22
59	JOHN	25
60	MARY	22
61	JOHN	25
62	MARY	22
63	JOHN	25
64	MARY	22
65	JOHN	25
66	MARY	22
67	JOHN	25
68	MARY	22
69	JOHN	25
70	MARY	22
71	JOHN	25
72	MARY	22
73	JOHN	25
74	MARY	22
75	JOHN	25
76	MARY	22
77	JOHN	25
78	MARY	22
79	JOHN	25
80	MARY	22
81	JOHN	25
82	MARY	22
83	JOHN	25
84	MARY	22
85	JOHN	25
86	MARY	22
87	JOHN	25
88	MARY	22
89	JOHN	25
90	MARY	22
91	JOHN	25
92	MARY	22
93	JOHN	25
94	MARY	22
95	JOHN	25
96	MARY	22
97	JOHN	25
98	MARY	22
99	JOHN	25
100	MARY	22

FIG. 2B




FIG. 3

FIG. 4





FIG. 4A








FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11






FIG. 12




FIG. 13

## SEQUENCE LISTING

&lt;110&gt; NORRIS, STEVEN J.

&lt;120&gt; VMP-LIKE SEQUENCES OF PATHOGENIC BORRELIA SPECIES AND STRAINS

&lt;130&gt; UTFH:264WO

&lt;140&gt; UNKNOWN

&lt;141&gt; 2003-12-22

&lt;150&gt; 60/435,077

&lt;151&gt; 2002-12-20

&lt;160&gt; 107

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1227

&lt;212&gt; DNA

&lt;213&gt; Borrelia burgdorferi

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (75)..(1142)

&lt;400&gt; 1

```

acctacactt gttaaaactc tcttttttgag ttaagatgat aacttatact tttcattata 60
aggagacgat gaat atg aaa aaa att tca agt gca agt tta tta aca act 110
          Met Lys Lys Ile Ser Ser Ala Ser Leu Leu Thr Thr
              1              5              10

ttc ttt gtt ttt att aat tgt aaa agc caa gtt gct gat aag gac gac 158
Phe Phe Val Phe Ile Asn Cys Lys Ser Gln Val Ala Asp Lys Asp Asp
          15              20              25

cca aca aac aaa ttt tac caa tct gtc ata caa tta ggt aac gga ttt 206
Pro Thr Asn Lys Phe Tyr Gln Ser Val Ile Gln Leu Gly Asn Gly Phe
          30              35              40

ctt gat gta ttc aca tct ttt ggt ggg tta gta gca gag gct ttt gga 254
Leu Asp Val Phe Thr Ser Phe Gly Gly Leu Val Ala Glu Ala Phe Gly
          45              50              55              60

ttt aaa tca gat cca aaa aaa tct gat gta aaa acc tat ttt act act 302
Phe Lys Ser Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Thr Thr
          65              70              75

gta gct gcc aaa ttg gaa aaa aca aaa acc gat ctt aat agt ttg cct 350
Val Ala Ala Lys Leu Glu Lys Thr Lys Thr Asp Leu Asn Ser Leu Pro
          80              85              90

aag gaa aaa agc gat ata agt agt acg acg ggg aaa cca gat agt aca 398
Lys Glu Lys Ser Asp Ile Ser Ser Thr Thr Gly Lys Pro Asp Ser Thr
          95              100              105

ggg tct gtt gga act gcc gtt gag ggg gct att aag gaa gtt agc gag 446
Gly Ser Val Gly Thr Ala Val Glu Gly Ala Ile Lys Glu Val Ser Glu
          110              115              120

ttg ttg gat aag ctg gta aaa gct gta aag aca gct gag ggg gct tca 494
Leu Leu Asp Lys Leu Val Lys Ala Val Lys Thr Ala Glu Gly Ala Ser

```

125	130	135	140	
agt ggt act gct gca att gga gaa gtt gtg gct gat gct gat gct gca				542
Ser Gly Thr Ala Ala Ile Gly Glu Val Val Ala Asp Ala Asp Ala Ala	145	150	155	
aag gtt gct gat aag gcg agt gtg aag ggg att gct aag ggg ata aag				590
Lys Val Ala Asp Lys Ala Ser Val Lys Gly Ile Ala Lys Gly Ile Lys	160	165	170	
gag att gtt gaa gct gct ggg ggg agt gaa aag ctg aaa gct gtt gct				638
Glu Ile Val Glu Ala Ala Gly Gly Ser Glu Lys Leu Lys Ala Val Ala	175	180	185	
gct gct aaa ggg gag aat aat aaa ggg gca ggg aag ttg ttt ggg aag				686
Ala Ala Lys Gly Glu Asn Asn Lys Gly Ala Gly Lys Lys Leu Phe Gly Lys	190	195	200	
gct ggt gct gct gct cat ggg gac agt gag gct gct agc aag gcg gct				734
Ala Gly Ala Ala Ala His Gly Asp Ser Glu Ala Ala Ser Lys Ala Ala	205	210	215	220
ggt gct gtt agt gct gtt agt ggg gag cag ata tta agt gcg att gtt				782
Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Ser Ala Ile Val	225	230	235	
acg gct gct gat gcg gct gag cag gat gga aag aag cct gag gag gct				830
Thr Ala Ala Asp Ala Ala Glu Gln Asp Gly Lys Lys Pro Glu Glu Ala	240	245	250	
aaa aat ccg att gct gct gct att ggg gat aaa gat ggg ggt gcg gag				878
Lys Asn Pro Ile Ala Ala Ala Ile Gly Asp Lys Asp Gly Gly Ala Glu	255	260	265	
ttt ggt cag gat gag atg aag aag gat gat cag att gct gct gct att				926
Phe Gly Gln Asp Glu Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Ile	270	275	280	
gct ttg agg ggg atg gct aag gat gga aag ttt gct gtg aag gat ggt				974
Ala Leu Arg Gly Met Ala Lys Asp Gly Lys Phe Ala Val Lys Asp Gly	285	290	295	300
gag aaa gag aag gct gag ggg gct att aag gga gct gct gag tct gca				1022
Glu Lys Glu Lys Ala Glu Gly Ala Ile Lys Gly Ala Ala Glu Ser Ala	305	310	315	
gtt cgc aaa gtt tta ggg gct att act ggg cta ata gga gac gcc gtg				1070
Val Arg Lys Val Leu Gly Ala Ile Thr Gly Leu Ile Gly Asp Ala Val	320	325	330	
agt tcc ggg cta agg aaa gtc ggt gat tca gtg aag gct gct agt aaa				1118
Ser Ser Gly Leu Arg Lys Val Gly Asp Ser Val Lys Ala Ala Ser Lys	335	340	345	
gaa aca cct cct gcc ttg aat aag tgattttaatt aagtgtatgg acacgactat				1172
Glu Thr Pro Pro Ala Leu Asn Lys	350	355		
gccctcatga ttgaggaaat agtcgagaga tatatatatact aaaagataat aaata				1227

<210> 2  
 <211> 356  
 <212> PRT

<213> *Borrelia burgdorferi*

&lt;400&gt; 2

```

Met Lys Lys Ile Ser Ser Ala Ser Leu Leu Thr Thr Phe Phe Val Phe
 1             5             10             15

Ile Asn Cys Lys Ser Gln Val Ala Asp Lys Asp Asp Pro Thr Asn Lys
      20             25             30

Phe Tyr Gln Ser Val Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe
      35             40             45

Thr Ser Phe Gly Gly Leu Val Ala Glu Ala Phe Gly Phe Lys Ser Asp
      50             55             60

Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Thr Thr Val Ala Ala Lys
      65             70             75             80

Leu Glu Lys Thr Lys Thr Asp Leu Asn Ser Leu Pro Lys Glu Lys Ser
      85             90             95

Asp Ile Ser Ser Thr Thr Gly Lys Pro Asp Ser Thr Gly Ser Val Gly
      100            105            110

Thr Ala Val Glu Gly Ala Ile Lys Glu Val Ser Glu Leu Leu Asp Lys
      115            120            125

Leu Val Lys Ala Val Lys Thr Ala Glu Gly Ala Ser Ser Gly Thr Ala
      130            135            140

Ala Ile Gly Glu Val Val Ala Asp Ala Asp Ala Ala Lys Val Ala Asp
      145            150            155            160

Lys Ala Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Glu Ile Val Glu
      165            170            175

Ala Ala Gly Gly Ser Glu Lys Leu Lys Ala Val Ala Ala Ala Lys Gly
      180            185            190

Glu Asn Asn Lys Gly Ala Gly Lys Leu Phe Gly Lys Ala Gly Ala Ala
      195            200            205

Ala His Gly Asp Ser Glu Ala Ala Ser Lys Ala Ala Gly Ala Val Ser
      210            215            220

Ala Val Ser Gly Glu Gln Ile Leu Ser Ala Ile Val Thr Ala Ala Asp
      225            230            235            240

Ala Ala Glu Gln Asp Gly Lys Lys Pro Glu Glu Ala Lys Asn Pro Ile
      245            250            255

Ala Ala Ala Ile Gly Asp Lys Asp Gly Gly Ala Glu Phe Gly Gln Asp
      260            265            270

Glu Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Ile Ala Leu Arg Gly
      275            280            285

Met Ala Lys Asp Gly Lys Phe Ala Val Lys Asp Gly Glu Lys Glu Lys
      290            295            300

Ala Glu Gly Ala Ile Lys Gly Ala Ala Glu Ser Ala Val Arg Lys Val
      305            310            315            320

Leu Gly Ala Ile Thr Gly Leu Ile Gly Asp Ala Val Ser Ser Gly Leu

```

```

325          330          335
Arg Lys Val Gly Asp Ser Val Lys Ala Ala Ser Lys Glu Thr Pro Pro
340          345          350

Ala Leu Asn Lys
355

<210> 3
<211> 1141
<212> DNA
<213> Borrelia hermsii

<220>
<221> CDS
<222> (1) .. (1062)

<400> 3
atg aga aaa aga ata agt gca ata ata atg act tta ttt atg gta tta 48
Met Arg Lys Arg Ile Ser Ala Ile Ile Met Thr Leu Phe Met Val Leu
1          5          10          15

gta agc tgt aat agc ggt ggg gtt gcg gaa gat cct aaa act gtg tat 96
Val Ser Cys Asn Ser Gly Gly Val Ala Glu Asp Pro Lys Thr Val Tyr
20          25          30

tta aca tct ata gct aat tta ggg aaa gga ttt tta gat gtt ttt gtg 144
Leu Thr Ser Ile Ala Asn Leu Gly Lys Gly Phe Leu Asp Val Phe Val
35          40          45

act ttt gga gat atg gtt act gga gct ttt ggt att aag gca gat act 192
Thr Phe Gly Asp Met Val Thr Gly Ala Phe Gly Ile Lys Ala Asp Thr
50          55          60

aag aaa agt gat ata ggg aag tat ttt act gat att gag agc act atg 240
Lys Lys Ser Asp Ile Gly Lys Tyr Phe Thr Asp Ile Glu Ser Thr Met
65          70          75          80

aca tca gtt aaa aag aag ttg caa gat gaa gtt gct aag aat ggt aac 288
Thr Ser Val Lys Lys Lys Leu Gln Asp Glu Val Ala Lys Asn Gly Asn
85          90          95

tat cca aag gta aag aca gct gtt gac gaa ttt gtt gca atc tta gga 336
Tyr Pro Lys Val Lys Thr Ala Val Asp Glu Phe Val Ala Ile Leu Gly
100          105          110

aag atc gag aaa gga gca aaa gaa gca tct aaa ggg gct act ggt gat 384
Lys Ile Glu Lys Gly Ala Lys Glu Ala Ser Lys Gly Ala Thr Gly Asp
115          120          125

gtt att att ggg aat act gtt aag aat ggt gat gct gta cct gga gaa 432
Val Ile Ile Gly Asn Thr Val Lys Asn Gly Asp Ala Val Pro Gly Glu
130          135          140

gca aca agt gtc aat tct ctt gtt aaa gga att aaa gaa ata gtt ggg 480
Ala Thr Ser Val Asn Ser Leu Val Lys Gly Ile Lys Glu Ile Val Gly
145          150          155          160

gta gtc ttg aag gaa ggt aag gca gat gct gat gct act aaa gat gat 528
Val Val Leu Lys Glu Gly Lys Ala Asp Ala Asp Ala Thr Lys Asp Asp
165          170          175

agt aag aaa gat att ggt aaa tta ttt acc gca acc act gat gcg aat 576

```

Ser Lys Lys Asp Ile Gly Lys Leu Phe Thr Ala Thr Thr Asp Ala Asn  
 180 185 190  
 aga gct gat aat gcg gca gct caa gca gct gca gcg tca ata gga gca 624  
 Arg Ala Asp Asn Ala Ala Ala Gln Ala Ala Ala Ser Ile Gly Ala  
 195 200 205  
 gtg aca ggt gct gat atc ttg caa gct ata gta caa tct aag gaa aat 672  
 Val Thr Gly Ala Asp Ile Leu Gln Ala Ile Val Gln Ser Lys Glu Asn  
 210 215 220  
 cct gtt gca aat agt act gat gga att gaa aaa gca aca gat gca gct 720  
 Pro Val Ala Asn Ser Thr Asp Gly Ile Glu Lys Ala Thr Asp Ala Ala  
 225 230 235 240  
 gag att gca gtt gct cca gct aaa gat aat aaa aaa gag att aaa gat 768  
 Glu Ile Ala Val Ala Pro Ala Lys Asp Asn Lys Lys Glu Ile Lys Asp  
 245 250 255  
 gga gca aaa aaa gac gca gtt att gct gca ggc att gca ctg cga gca 816  
 Gly Ala Lys Lys Asp Ala Val Ile Ala Ala Gly Ile Ala Leu Arg Ala  
 260 265 270  
 atg gct aag aat ggt aca ttt tct att aaa aac aat gaa gat gcg gct 864  
 Met Ala Lys Asn Gly Thr Phe Ser Ile Lys Asn Asn Glu Asp Ala Ala  
 275 280 285  
 gta acg acg ata aat agt gca gca gca agc gca gtg aac aag att tta 912  
 Val Thr Thr Ile Asn Ser Ala Ala Ala Ser Ala Val Asn Lys Ile Leu  
 290 295 300  
 agc act cta ata ata gca ata agg aat aca gtt gat agt ggt tta aaa 960  
 Ser Thr Leu Ile Ile Ala Ile Arg Asn Thr Val Asp Ser Gly Leu Lys  
 305 310 315 320  
 aca ata aat gag gct ctt gct aca gtt aaa caa gaa gat aaa tct gta 1008  
 Thr Ile Asn Glu Ala Leu Ala Thr Val Lys Gln Glu Asp Lys Ser Val  
 325 330 335  
 gaa gca act aat act gca gaa gca aca act agt ggt cag caa gcg aaa 1056  
 Glu Ala Thr Asn Thr Ala Glu Ala Thr Thr Ser Gly Gln Gln Ala Lys  
 340 345 350  
 aac tag ttaagggttaa atataaagga taaagttatt gtaagggaaa agcttttctt 1112  
 Asn  
 gtttttaatg caggaatgta gtttctctg 1141

&lt;210&gt; 4

&lt;211&gt; 353

&lt;212&gt; PRT

&lt;213&gt; Borrelia hermsii

&lt;400&gt; 4

Met Arg Lys Arg Ile Ser Ala Ile Ile Met Thr Leu Phe Met Val Leu  
 1 5 10 15  
 Val Ser Cys Asn Ser Gly Gly Val Ala Glu Asp Pro Lys Thr Val Tyr  
 20 25 30  
 Leu Thr Ser Ile Ala Asn Leu Gly Lys Gly Phe Leu Asp Val Phe Val  
 35 40 45  
 Thr Phe Gly Asp Met Val Thr Gly Ala Phe Gly Ile Lys Ala Asp Thr  
 50 55 60  
 Lys Lys Ser Asp Ile Gly Lys Tyr Phe Thr Asp Ile Glu Ser Thr Met

```

      65              70              75              80
Thr Ser Val Lys Lys Lys Leu Gln Asp Glu Val Ala Lys Asn Gly Asn
      85              90              95
Tyr Pro Lys Val Lys Thr Ala Val Asp Glu Phe Val Ala Ile Leu Gly
      100             105             110
Lys Ile Glu Lys Gly Ala Lys Glu Ala Ser Lys Gly Ala Thr Gly Asp
      115             120             125
Val Ile Ile Gly Asn Thr Val Lys Asn Gly Asp Ala Val Pro Gly Glu
      130             135             140
Ala Thr Ser Val Asn Ser Leu Val Lys Gly Ile Lys Glu Ile Val Gly
      145             150             155             160
Val Val Leu Lys Glu Gly Lys Ala Asp Ala Asp Ala Thr Lys Asp Asp
      165             170             175
Ser Lys Lys Asp Ile Gly Lys Leu Phe Thr Ala Thr Thr Asp Ala Asn
      180             185             190
Arg Ala Asp Asn Ala Ala Ala Gln Ala Ala Ala Ser Ile Gly Ala
      195             200             205
Val Thr Gly Ala Asp Ile Leu Gln Ala Ile Val Gln Ser Lys Glu Asn
      210             215             220
Pro Val Ala Asn Ser Thr Asp Gly Ile Glu Lys Ala Thr Asp Ala Ala
      225             230             235             240
Glu Ile Ala Val Ala Pro Ala Lys Asp Asn Lys Lys Glu Ile Lys Asp
      245             250             255
Gly Ala Lys Lys Asp Ala Val Ile Ala Ala Gly Ile Ala Leu Arg Ala
      260             265             270
Met Ala Lys Asn Gly Thr Phe Ser Ile Lys Asn Asn Glu Asp Ala Ala
      275             280             285
Val Thr Thr Ile Asn Ser Ala Ala Ala Ser Ala Val Asn Lys Ile Leu
      290             295             300
Ser Thr Leu Ile Ile Ala Ile Arg Asn Thr Val Asp Ser Gly Leu Lys
      305             310             315             320
Thr Ile Asn Glu Ala Leu Ala Thr Val Lys Gln Glu Asp Lys Ser Val
      325             330             335
Glu Ala Thr Asn Thr Ala Glu Ala Thr Thr Ser Gly Gln Gln Ala Lys
      340             345             350
Asn

```

<210> 5  
 <211> 416  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(414)

```

<400> 5
aag ggg att gcg aag ggg ata aag ggg att gtt gcg gct gct ggg aag 48
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala Gly Lys
1 5 10 15

gct ttt ggc aag gat ggt gat gcg ctg aca ggt gtt gca aaa gct gct 96
Ala Phe Gly Lys Asp Gly Asp Ala Leu Thr Gly Val Ala Lys Ala Ala
20 25 30

gag aat gat gct aac aag gat gcg ggg aag ttg ttt gct ggt aag aat 144
Glu Asn Asp Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
35 40 45

ggt aat gct ggt gct gct gac att gcg aag gcg gct gct gct gtt act 192
Gly Asn Ala Gly Ala Ala Asp Ile Ala Lys Ala Ala Ala Ala Val Thr
50 55 60

```



```

gcg gtt agt ggg gag cag ata cta aaa gct att gtt gag gcg gct ggt      240
Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala Ala Gly
 65                               70                               75                               80

gat gcg gat cag gcg ggt gta aag gct gat gcg gct aag aat ccg att      288
Asp Ala Asp Gln Ala Gly Val Lys Ala Asp Ala Ala Lys Asn Pro Ile
                               85                               90                               95

gca gct gcg att ggg act gct gat gat ggt gct gcg ttt ggt aag gat      336
Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Ala Phe Gly Lys Asp
                               100                              105                              110

gag atg aag aag aga aat gat aag att gtt gca gct att gtt ttg agg      384
Glu Met Lys Lys Arg Asn Asp Lys Ile Val Ala Ala Ile Val Leu Arg
                               115                              120                              125

ggg gtg cct aag gat gga aag ttt gct gct aa                          416
Gly Val Pro Lys Asp Gly Lys Phe Ala Ala
                               130                              135

```

<210> 6  
 <211> 138  
 <212> PRT  
 <213> Borrelia afzelii

```

<400> 6
Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala Gly Lys
 1                               5                               10                               15

Ala Phe Gly Lys Asp Gly Asp Ala Leu Thr Gly Val Ala Lys Ala Ala
 20                              25                              30

Glu Asn Asp Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn
 35                              40                              45

Gly Asn Ala Gly Ala Ala Asp Ile Ala Lys Ala Ala Ala Val Thr
 50                              55                              60

Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala Ala Gly
 65                               70                               75                               80

Asp Ala Asp Gln Ala Gly Val Lys Ala Asp Ala Ala Lys Asn Pro Ile
 85                               90                               95

Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Ala Phe Gly Lys Asp
100                              105                              110

Glu Met Lys Lys Arg Asn Asp Lys Ile Val Ala Ala Ile Val Leu Arg
115                              120                              125

Gly Val Pro Lys Asp Gly Lys Phe Ala Ala
130                              135

```

<210> 7  
 <211> 413  
 <212> DNA  
 <213> Borrelia afzelii

<220>  
 <221> CDS  
 <222> (1)..(411)

&lt;400&gt; 7

aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag 48  
 Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys  
 1 5 10 15

gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt gca aaa gtt gct 96  
 Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val Ala  
 20 25 30

gat gat gat aac aag gat gcg ggg aag ttg ttt gct ggt aag aat ggt 144  
 Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn Gly  
 35 40 45

ggt gct ggt gct gct gat gcg att ggg aag gcg gct gct gct gtt act 192  
 Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Ala Val Thr  
 50 55 60

gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gat gct gct ggt 240  
 Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly  
 65 70 75 80

gct gca gct aat cag gcg ggt aaa aag gct gcg gat gct aag aat ccg 288  
 Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn Pro  
 85 90 95

att gcg gct gcg att ggg act gct gat gat ggg gcg gag ttt aag gat 336  
 Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys Asp  
 100 105 110

gat atg aag aag agt gat aat att gct gcg gct att gtt ttg agg ggg 384  
 Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg Gly  
 115 120 125

gtg cct aag gat gga aag ttt gct gct aa 413  
 Val Pro Lys Asp Gly Lys Phe Ala Ala  
 130 135

&lt;210&gt; 8

&lt;211&gt; 137

&lt;212&gt; PRT

<213> *Borrelia afzelii*

&lt;400&gt; 8

Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys  
 1 5 10 15

Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val Ala  
 20 25 30

Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn Gly  
 35 40 45

Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Val Thr  
 50 55 60

Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly  
 65 70 75 80

Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn Pro  
 85 90 95

Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys Asp  
 100 105 110

Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg Gly  
 115 120 125

Val Pro Lys Asp Gly Lys Phe Ala Ala  
 130 135

<210> 9

<211> 428

<212> DNA

<213> *Borrelia afzelii*

<220>

<221> CDS

<222> (1) .. (426)

<400> 9

aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag 48  
 Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys  
 1 5 10 15

gct ttt ggt aag gag ggt gat gcg ctg aag gat gtt gca aaa gtt gct 96  
 Ala Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala  
 20 25 30

gat gag aat ggg gat aac aag gat gcg ggg aag ttg ttt gct ggt gag 144  
 Asp Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu  
 35 40 45

aat ggt aat gct ggt ggt gct gct gat gct gac att gcg aag gcg gct 192  
 Asn Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala  
 50 55 60

gct gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt 240  
 Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
 65 70 75 80

gag gcg gct ggt gct gcg gat cag gcg ggt gta aag gct gag gag gct 288  
 Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala Glu Glu Ala  
 85 90 95

aag aat ccg att gca gct gcg att ggg act gat gat gct ggt gcg gcg 336  
 Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly Ala Ala  
 100 105 110

gag ttt ggt gag aat gat atg aag aag aat gat aat att gct gcg gct 384  
 Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile Ala Ala Ala  
 115 120 125

att gtt ttg agg ggg gtg cct aag gat gga aag ttt gct gct aa 428  
 Ile Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala  
 130 135 140

<210> 10

<211> 142

<212> PRT

<213> *Borrelia afzelii*

<400> 10

Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys  
 1 5 10 15

Ala Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala  
                   20                  25                  30  
 Asp Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu  
                   35                  40                  45  
 Asn Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala  
                   50                  55                  60  
 Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
                   65                  70                  75                  80  
 Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala Glu Glu Ala  
                   85                  90                  95  
 Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly Ala Ala  
                   100                  105                  110  
 Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile Ala Ala Ala  
                   115                  120                  125  
 Ile Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala  
                   130                  135                  140

<210> 11  
 <211> 426  
 <212> DNA  
 <213> Borrelia afzelii

<220>  
 <221> CDS  
 <222> (3)..(425)

<400> 11  
 ag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag 47  
   Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys  
     1                  5                  10                  15  
 gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt aaa aca gtt gct 95  
 Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr Val Ala  
                   20                  25                  30  
 gct gag aat gag gct aac aag gat gcg ggg aag ttg ttt gct ggt aag 143  
 Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys  
                   35                  40                  45  
 aat ggt aat gct gat gct gct gat gct gct gac att gcg aag gcg gct 191  
 Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys Ala Ala  
                   50                  55                  60  
 ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa gct att gtt 239  
 Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
                   65                  70                  75  
 gat ggt gct ggt gat gca gct aat cag gcg ggt aaa aag gct gct gag 287  
 Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Glu  
                   80                  85                  90                  95  
 gct aag aat ccg att gcg gct gcg att ggg act aat gaa gct ggg gcg 335  
 Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala  
                   100                  105                  110  
 gag ttt ggt gat gat atg aag aag aga aat gat aag att gct gcg gct 383

Glu Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala  
 115 120 125

att gtt ttg agg ggg gtg cct aag gat gga aag ttt gct gct a 426  
 Ile Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala  
 130 135 140

<210> 12

<211> 141

<212> PRT

<213> *Borrelia afzelii*

<400> 12

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala  
 1 5 10 15

Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr Val Ala Ala  
 20 25 30

Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn  
 35 40 45

Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys Ala Ala Gly  
 50 55 60

Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp  
 65 70 75 80

Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Glu Ala  
 85 90 95

Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala Glu  
 100 105 110

Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala Ile  
 115 120 125

Val Leu Arg Gly Val Pro Lys Asp Gly Lys Phe Ala Ala  
 130 135 140

<210> 13

<211> 396

<212> DNA

<213> *Borrelia garinii*

<220>

<221> CDS

<222> (2)..(394)

<400> 13

g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa 49  
 Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu  
 1 5 10 15

ggg aag ttg aat gct gct ggt gct gag ggt acg act aac gcg gat gct 97  
 Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala Asp Ala  
 20 25 30

ggg aag ttg ttt gtg aag aat gct ggt aat gtg ggt ggt gaa gca ggt 145  
 Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu Ala Gly  
 35 40 45

```

gat gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag 193
Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln
   50               55               60

ata tta aaa gcg att gtt gat gct gct aag gat ggt ggt gag aag cag 241
Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu Lys Gln
   65               70               75               80

ggt aag aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg 289
Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly
           85               90               95

ggt gcg ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag 337
Gly Ala Gly Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys
           100               105               110

gat gat cag att gct gct gct atg gtt ctg agg gga atg gct aag gat 385
Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp
           115               120               125

ggg cag ttt gc
Gly Gln Phe
   130

```

<210> 14  
 <211> 131  
 <212> PRT  
 <213> *Borrelia garinii*

```

<400> 14
Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
  1               5               10               15

Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala Asp Ala
  20               25               30

Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu Ala Gly
  35               40               45

Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln
  50               55               60

Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu Lys Gln
  65               70               75               80

Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly
           85               90               95

Gly Ala Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys
           100               105               110

Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp
           115               120               125

Gly Gln Phe
   130

```

<210> 15  
 <211> 390  
 <212> DNA  
 <213> *Borrelia garinii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (2) .. (388)

&lt;400&gt; 15

```

g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa 49
  Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
    1             5             10             15

ggg aag ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct 97
Gly Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala
      20             25             30

ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat 145
Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp
      35             40             45

gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 193
Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile
      50             55             60

tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag 241
Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys
      65             70             75             80

aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt gcg 289
Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala
      85             90             95

ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag gat gat 337
Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp
      100            105            110

cag att gct act gct atg gtt ctg agg gga atg gct aag gat ggg cag 385
Gln Ile Ala Thr Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln
      115            120            125

ttt gc
Phe
                                     390

```

&lt;210&gt; 16

&lt;211&gt; 129

&lt;212&gt; PRT

<213> *Borrelia garinii*

&lt;400&gt; 16

```

Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu
  1             5             10             15

Gly Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala
  20             25             30

Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp
  35             40             45

Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile
  50             55             60

Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys
  65             70             75             80

Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala
  85             90             95

```

Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp  
 100 105 110

Gln Ile Ala Thr Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln  
 115 120 125

Phe

<210> 17

<211> 390

<212> DNA

<213> *Borrelia garinii*

<220>

<221> CDS

<222> (2)..(388)

<400> 17

g ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa 49  
 Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu  
 1 5 10 15

ggg agg ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct 97  
 Gly Arg Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala  
 20 25 30

ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat 145  
 Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp  
 35 40 45

gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 193  
 Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
 50 55 60

tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag 241  
 Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys  
 65 70 75 80

aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt gcg 289  
 Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala  
 85 90 95

ggt gat aat gat gct gct gcg gcg ttt gct act atg aag aag gat gat 337  
 Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp  
 100 105 110

cag att gct gct gct atg gtt ctg agg gga atg gct aag gat ggg cag 385  
 Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln  
 115 120 125

ttt gc 390  
 Phe

<210> 18

<211> 129

<212> PRT

<213> *Borrelia garinii*

<400> 18

Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu



1                      5                      10                      15  
 Gly Arg Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala  
                     20                      25                      30  
 Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp  
                     35                      40                      45  
 Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
                     50                      55                      60  
 Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys  
                     65                      70                      75                      80  
 Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala  
                     85                      90                      95  
 Gly Asp Asn Asp Ala Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp  
                     100                      105                      110  
 Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln  
                     115                      120                      125  
 Phe

<210> 19  
 <211> 339  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (2)..(337)

<400> 19  
 g ggg ata aag ggg att gtt gat gct gct ggt gaa act aac aag gat gct 49  
   Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Glu Thr Asn Lys Asp Ala  
     1                      5                      10                      15  
 ggg aag ttg ttt gtg aag aag aat aat gag ggt ggt gaa gca aat gat 97  
   Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp  
                     20                      25                      30  
 gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 145  
   Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
                     35                      40                      45  
 tta aaa gcg att gtt gat gct gct gag ggt ggt gag aag cag ggt aag 193  
   Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys  
                     50                      55                      60  
 aag gct gcg gat gct aca aat ccg att gag gcg gct att ggg ggt aca 241  
   Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Thr  
                     65                      70                      75                      80  
 aat gat aat gat gct gcg gcg ttt gct act atg aag aag gat gat cag 289  
   Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp Gln  
                     85                      90                      95  
 att gct gct gct atg gtt ctg agg gga atg gct aag gat ggg cag ttt 337  
   Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe  
                     100                      105                      110

gc

339

<210> 20  
 <211> 112  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 20  
 Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Glu Thr Asn Lys Asp Ala  
   1                  5                  10                  15  
 Gly Lys Leu Phe Val Lys Lys Asn Asn Glu Gly Gly Glu Ala Asn Asp  
                   20                  25                  30  
 Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
           35                  40                  45  
 Leu Lys Ala Ile Val Asp Ala Ala Glu Gly Gly Glu Lys Gln Gly Lys  
       50                  55                  60  
 Lys Ala Ala Asp Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Thr  
       65                  70                  75                  80  
 Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met Lys Lys Asp Asp Gln  
                   85                  90                  95  
 Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe  
           100                  105                  110

<210> 21  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
       Primer

<400> 21  
 ccagcaaaca acttccccgc c

21

<210> 22  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
       Primer

<400> 22  
 atccttaaac tccgccccat catc

24

<210> 23  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 23  
gagtgtgtg gagagtgtg ttgatgag 28

<210> 24  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 24  
ggggataaag gggattgtg atgctgc 27

<210> 25  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 25  
gcaaactgcc catccttagc cattcc 26

<210> 26  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 26  
aaggggattg cgaaggggat aaagg 25

<210> 27  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 27  
ttagcagcaa actttccatc cttagcc 27

<210> 28  
<211> 5897  
<212> DNA  
<213> Borrelia garinii

&lt;400&gt; 28

```

cggaaatcaa gccacctaata acaacttccc aaaagtttct caaaaaatat tatattcagc 60
agtaaattct ataagtcatt aattattttaa tactattcaa cagtaaattc tataagtcac 120
taattattta atactattca gcagtaaatt ctataagtca ttaattattt aatactattc 180
agcagtaaat tctataagtc attaattatt taatactatt cagcagtaaa ttctataagt 240
cattaattat ttaatactat tcagcagtaa attctataag tcattaatta tttaatacta 300
ttcagcagta aattctataa gtcattaatt caattaggta acggattcct agatgtattc 360
acctcttttg gtggattagt tgcagatgca ttggggttta aagctgatcc aaaaaaatct 420
gatgtaaaaa cttattttga atctctagct aaaaaattag aagaaacaaa agatggttta 480
actaagttgt ccaaaggtaa tgacgggtgat actggaaagg ctgggtgatgc tgggtggggc 540
gggtgggtggc ctagtgctgc aggtggcgct ggtgggattg agggcgctat aacagagatt 600
agcaaattgt tagatgatag ggcaaaagct gctgcggaag ctgcaagtgc tgctactggc 660
aatgcagcaa ttggggatgt tgtaattggt aatgggtggag cagcaaaagg tgggtgatgcg 720
gagagtgtta atggggata taaggggattg aaggggattg ttgatgctgc tgagaaggct 780
gatgcgaagg aagggaagtt ggatgtggct ggtgatgctg gtggggctgg tgggtggcgct 840
gggtgctgcag gtggcgctgg tgggattgag ggcgctataa cagagattag caaatggtta 900
gatgatattg caaaagctgc tgcggttgct gcaagtgcct caagtgcctc tactggtaac 960
gcagcaattg gggatgttgt taatggtaac gatggagcag caaaagggtg tgatgcggcg 1020
agtgttaatt ggattgctaa ggggataaag gggattgttg atgctgctga gaaggctgat 1080
gcgaaggaa ggaagtggga tgtggctggg gatgctggg agggtaacaa ggtgctggg 1140
aagctgtttg tgaagaagaa tgctgggtgat gaggggtggg aagcaaatga tgctgggaag 1200
gctgctgctg cggttgctgc tgtagtgctg gagcagatat taaaagcgat tgttgatgct 1260
gctgaggggtg atgataagca gggtaagaag gctgcggatg ctacaaatcc gattgagggc 1320
gctattgggg gtgcggatgc ggggtgctaa gctgagggct ttaataagat gaagaaggat 1380
gatcagattg ctgctgctat tgaagggact ggaatggcta aggatgggca gtttgccttg 1440
aaggatgatg ctgctgctca tgcaagtgc gctactggta gtacaacgat tggagatgct 1500
gctgcggaag ctgcaagtgc agatgggtgat gctgaggaag ttaatgggat tgctaagggg 1620
gttaagagtg gtgaggcaaa tgctgagaag gctgatgcga aggaagggaa gttggatgtg 1680
ataaagggga ttggtgatgc taacgtgaat gttgggaagt tgtttgtgaa gaataatgg 1740
gctggtgctg ctggtacgac tgatgctggg aaagctgctg ctgctgctgc tgctgcttag 1800
aatgaggggtg gtgagcgaag gattgttgat gctgctaaag atgggtgataa gcaggggggt 1860
ggggagcaga tattaaaagc aaatccgatt gaggcggcta ttgggggtac aaatgataat 1920
actgatgtaa aggatgctac tatgaagaag gatgatcaga ttgctgctgc tatggttctg 1980
gatgctgcgg cgtttgctac gcaaggtgg tgctgaaaat cggctatttg ggtgctgat 2040
aggggaatgg ctaaggtggg tgctgaaaat atgaagaagg atgatcagat tgctgctgct 2100
aaaacggggg ttgctgcgga cggcgtttta taaggagggt cagtttgctt tgacgaataa tgctgctgct 2160
gctgatgctg ggggaatggc taaggatggg atggcaaaag ctgctgcggg tgctgcaagt 2220
catgaaggga ctgtaagaa gcaatgcagc aattggggat gttgttaaga gtaatggtg 2280
gctgctactg aaaggtgggt atgctgctgc aagttggatg tggctggtgc agcagcagca 2340
aaaggtgggt aggtgatgc gaaggagggt gttgtttgtg aagaagaatg gtgatgatg 2400
actaacaagg atgctgggaa ggaaggctgc tgctgcgggt gctgctgcta gtggggagca gatattaaaa 2460
ggtgatgctg ggaaggctgc agatgggtgat aagacggggg ttactgatgt aaaggatgct 2520
gcgattgttg atgctgctaa tattgggggg gctatggttc tgaggggaat ggctaaggat 2580
acaaatccga aagatgaaga aggatgatca taatgatcat gataatcata aggggactgt taagaatgct 2640
aagatgaaga aggatgatca taatgatcat gataatcata aggggactgt taagaatgct 2680
gggcagtttg ctttgaagaa taatgatcat gataatcata aggggactgt taagaatgct 2720
ggtgatattg caaaggcgcg tgaggaagct caagtgcctg tactggtaac 2780
gcagcgattg gggatgttgt taagaatagt ggggcagcag caaaagggtg tgagggcgcg 2840
agtgttaatg ggattgctaa ggggataaag gggattgttg atgctgctgg aaaggctgat 2900
gcgaaggaa ggaagtggga tgcactgggt gctgagggta cgactaacgt gaatgctggg 3060
aagttgtttg tgaagagggc ggctgatgat ggtggtgatg cagatgatgc tgggaaggct 3120
gctgctgcgg ttgctgcaag aggtggtgat ggttaatcag cgattggaga tgttgttaac 3180
gggtgatgtg caaaagcaaa tgctgagaag gctgaggaag ttaatgggat tgctaagggg 3240
ataaagggga ttggtgatgc tgctgagaag gctgatgcga aggaagggaa gttgaatgct 3300
gctggtgctg agggtagcac taacgcggat gctgggaagt tgtttgtgaa gaatgctgg 3360
aatgtgggtg gtgaagcagg tgatgctggg aaggctgctg ctgctggttc atggtggtga gaagcagggt 3420
ggggagcaga tattaaaagc aaatccgatt gctgctaagg tgggggttac aaatgataat 3480
aagaaggctg cggatgctac tactatgaag gacgcggcta aaggatgatc agattgctgc tgctatggtt 3540
gatgctgctg cggcgtttgc tgggcaattt aaggatgatc agattgctgc tgctatggtt 3600
ctgaggggaa tggctaagga tgggcaattt aaggatgatc agattgctgc tgctatggtt 3660
gggactgtta agaattgctg tgatataata aaggctgctg atgctgctgc tgctcatgaa 3720
agtgctgcta ctggtagtgc agcaattggg gatgttgtta atggtaatgg agcaacagca 3780

```

```

aaagggtggtg atgcgaagag tgttaaatggg attgctaagg ggataaaggg gattgttgat 3840
gctgctgaga aggctgatgc gaaggaaggg aagttggatg tggctggtga tgctggtgaa 3900
actaacaagg atgctgggaa gttgtttgtg aagaacaatg gtaatgaggg tgggtgatgca 3960
gatgatgctg ggaaggctgc tgctgcgggt gctgctgtta gtggggagca gatattaaaa 4020
gcgattgttg atgctgctaa ggggtggatg aagacgggta agaataatgt gaaggatgct 4080
gaaaatccga ttgaggcggc tattgggagt agtgcggtat ctgatgctgc ggcgtttaat 4140
aaggagggga tgaagaagga tgatcagatt gctgctgcta tggttctgag gggaatggct 4200
aaggatgggc agtttgcttt gacgaatgat gctgctgctc atgaaggac tgtaagaat 4260
gctgttggga gtgcaacaaa taagaccgtt gttgcttttg ctaacttggg tcgaaagacc 4320
gtgcaagctg ggttgaagaa ggttggggat gttgttaaga atagtgaggg aaaagatggg 4380
gatgcggcga gtgttaatgg gattgctaag gggataaagg ggattgttga tgctgctgag 4440
aaggctgatg cgaaggaagg gaagttggat gtggctggtg ctgctggtga aactaacaag 4500
gatgctggga agttgtttgt gaagaagaat aatgaggggt gtgaagcaaa tgatgctggg 4560
aaggctgctg ctgctgttgc tgctgttagt ggggagcaga tattaaaagc gattgttgat 4620
gctgctaagg atggtgatga taagcagggt aggaaggctg aggatgctac aaatccgatt 4680
gacgcggcta ttgggggtgc aggtgcgggt gctaattgct ctgcggcgtt taataatatg 4740
aagaaggatg atcagattgc tgctgctatg gttctgaggg gaatggctaa ggatgggcag 4800
tttgctttga cgaataatgc tcatactaat cataagggga ctgttaagaa tgctgttgat 4860
atgacaaaag ctgctgcggg tgctgcaagt gctgcaagt ctgctactgg taatgcagca 4920
attggggatg ttgttaatgg taatgatgga gcagcaaaaag gtggtgatgc ggcgagtgtt 4980
aatgggattg ctaaggggat aaaggggatt gttgatgctg ctgagaaggc tgatgcgaag 5040
gaagggaagt tgaatgtggc tgggtgctgct ggtgctgagg gtaacgaggc tgctgggaag 5100
ctgtttgtga agaagaatgc tggatgatcat ggtggtgaag caggtgatgc tgggagggct 5160
gctgctgcgg ttgctgctgt tagtggggag cagatattaa aagcgattgt tgatgctgct 5220
aaggatggtg gtgataagca gggtaagaag gctgaggatg ctgaaaatcc gattgacgcg 5280
gctattggga gtacgggtgc ggatgataat gctgctgagg cgtttgctac tatgaagaag 5340
gatgatcaga ttgctgctgc tatggttctg aggggaatgg ctaaggatgg gcagtttgct 5400
ttgaaggatg ctgctcatga taatcataag gggactgtta agaatgctgt tgatataata 5460
aaggctactg cggttgctgc aagtgcctgct actggtagta caacgattgg ggatgttggt 5520
aagaatggtg aggcaaaaagg tggtgaggcg aagagtgtta atgggattgc taaggggata 5580
aaggggattg ttgatgctgc tggaaaggct gatgcgaagg aaggggaagt gaatgtggct 5640
ggtgctgctg gtgagggtaa cgaggctgct gggaagctgt ttgtgtaaat tactatagga 5700
ttagaactag tgtacgatat gagtcctttg gttattttgc agctgctaata gaatttgaaa 5760
taagtgaagt taaaattgct gatgttaatg gaacacattt tattgctaca aaagagaaag 5820
aatattata tgattcactt gatttaaggg ctctgaggaa aatatttgaa ataacttcaa 5880
agcgaatggt taagctt                                     5897

```

<210> 29  
 <211> 300  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(300)

```

<400> 29
gtc att aat tat tta ata cta ttc agc agt aaa ttc tat aag tca tta 48
Val Ile Asn Tyr Leu Ile Leu Phe Ser Ser Lys Phe Tyr Lys Ser Leu
1 5 10 15

att caa tta ggt aac gga ttc tta gat gta ttc acc tct ttt ggt gga 96
Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe Thr Ser Phe Gly Gly
20 25 30

tta gtt gca gat gca ttg ggg ttt aaa gct gat cca aaa aaa tct gat 144
Leu Val Ala Asp Ala Leu Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp
35 40 45

gta aaa act tat ttt gaa tct cta gct aaa aaa tta gaa gaa aca aaa 192
Val Lys Thr Tyr Phe Glu Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys
50 55 60

```

gat ggt tta act aag ttg tcc aaa ggt aat gac ggt gat act gga aag 240  
 Asp Gly Leu Thr Lys Leu Ser Lys Gly Asn Asp Gly Asp Thr Gly Lys  
 65 70 75 80

gct ggt gat gct ggt ggg gct ggt ggt ggc gct agt gct gca ggt ggc 288  
 Ala Gly Asp Ala Gly Gly Ala Gly Gly Gly Ala Ser Ala Ala Gly Gly  
 85 90 95

gct ggt ggg att 300  
 Ala Gly Gly Ile  
 100

<210> 30  
 <211> 100  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 30  
 Val Ile Asn Tyr Leu Ile Leu Phe Ser Ser Lys Phe Tyr Lys Ser Leu  
 1 5 10 15

Ile Gln Leu Gly Asn Gly Phe Leu Asp Val Phe Thr Ser Phe Gly Gly  
 20 25 30

Leu Val Ala Asp Ala Leu Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp  
 35 40 45

Val Lys Thr Tyr Phe Glu Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys  
 50 55 60

Asp Gly Leu Thr Lys Leu Ser Lys Gly Asn Asp Gly Asp Thr Gly Lys  
 65 70 75 80

Ala Gly Asp Ala Gly Gly Ala Gly Gly Gly Ala Ser Ala Ala Gly Gly  
 85 90 95

Ala Gly Gly Ile  
 100

<210> 31  
 <211> 102  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(102)

<400> 31  
 ggg ttt aaa gct gat cca aaa aaa tct gat gta aaa act tat ttt gaa 48  
 Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Glu  
 1 5 10 15

tct cta gct aaa aaa tta gaa gaa aca aaa gat ggt tta act aag ttg 96  
 Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys Asp Gly Leu Thr Lys Leu  
 20 25 30

tcc aaa 102  
 Ser Lys

<210> 32  
 <211> 34  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 32  
 Gly Phe Lys Ala Asp Pro Lys Lys Ser Asp Val Lys Thr Tyr Phe Glu  
     1                    5                    10                    15  
 Ser Leu Ala Lys Lys Leu Glu Glu Thr Lys Asp Gly Leu Thr Lys Leu  
                     20                    25                    30  
 Ser Lys

<210> 33  
 <211> 288  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(288)

<400> 33  
 gag ggc gct ata aca gag att agc aaa tgg tta gat gat atg gca aaa 48  
 Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys  
     1                    5                    10                    15  
 gct gct gcg gaa gct gca agt gct gct act ggt aat gca gca att ggg 96  
 Ala Ala Ala Glu Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly  
                     20                    25                    30  
 gat gtt gtt aat ggt aat ggt gga gca gca aaa ggt ggt gat gcg gag 144  
 Asp Val Val Asn Gly Asn Gly Gly Ala Ala Lys Gly Gly Asp Ala Glu  
                     35                    40                    45  
 agt gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct 192  
 Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala  
                     50                    55                    60  
 gag aag gct gat gcg aag gaa ggg aag ttg gat gtg gct ggt gat gct 240  
 Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala  
     65                    70                    75                    80  
 ggt ggg gct ggt ggt ggc gct ggt gct gca ggt ggc gct ggt ggg att 288  
 Gly Gly Ala Gly Gly Gly Ala Gly Ala Gly Gly Ala Gly Gly Ile  
                     85                    90                    95

<210> 34  
 <211> 96  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 34  
 Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys  
     1                    5                    10                    15  
 Ala Ala Ala Glu Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly  
                     20                    25                    30  
 Asp Val Val Asn Gly Asn Gly Gly Ala Ala Lys Gly Gly Asp Ala Glu

35 40 45

Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala  
50 55 60

Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala  
65 70 75 80

Gly Gly Ala Gly Gly Gly Ala Gly Ala Ala Gly Gly Ala Gly Gly Ile  
85 90 95

<210> 35  
<211> 594  
<212> DNA  
<213> *Borrelia garinii*

<220>  
<221> CDS  
<222> (1)..(594)

<400> 35

gag ggc gct ata aca gag att agc aaa tgg tta gat gat atg gca aaa	48
Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys	
1 5 10 15	
gct gct gcg gtt gct gca agt gct gca agt gct gct act ggt aat gca	96
Ala Ala Ala Val Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala	
20 25 30	
gca att ggg gat gtt gtt aat ggt aat gat gga gca gca aaa ggt ggt	144
Ala Ile Gly Asp Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly	
35 40 45	
gat gcg gcg agt gtt aat ggg att gct aag ggg ata aag ggg att gtt	192
Asp Ala Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val	
50 55 60	
gat gct gct gag aag gct gat gcg aag gaa ggg aag ttg gat gtg gct	240
Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala	
65 70 75 80	
ggt gat gct ggt gag ggt aac aag gat gct ggg aag ctg ttt gtg aag	288
Gly Asp Ala Gly Glu Gly Asn Lys Asp Ala Gly Lys Leu Phe Val Lys	
85 90 95	
aag aat gct ggt gat gag ggt ggt gaa gca aat gat gct ggg aag gct	336
Lys Asn Ala Gly Asp Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala	
100 105 110	
gct gct gcg gtt gct gct gtt agt ggg gag cag ata tta aaa gcg att	384
Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile	
115 120 125	
gtt gat gct gct gag ggt gat gat aag cag ggt aag aag gct gcg gat	432
Val Asp Ala Ala Glu Gly Asp Asp Lys Gln Gly Lys Lys Ala Ala Asp	
130 135 140	
gct aca aat ccg att gag gcg gct att ggg ggt gcg gat gcg ggt gct	480
Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala Asp Ala Gly Ala	
145 150 155 160	
aat gct gag gcg ttt aat aag atg aag aag gat gat cag att gct gct	528
Asn Ala Glu Ala Phe Asn Lys Met Lys Lys Asp Asp Gln Ile Ala Ala	



	165		170		175	
gct atg gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg aag						576
Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys						
	180		185		190	

gat gat gct gct gct cat						594
Asp Asp Ala Ala Ala His						
	195					

<210> 36  
 <211> 198  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 36																			
Glu Gly Ala Ile Thr Glu Ile Ser Lys Trp Leu Asp Asp Met Ala Lys																			
1				5				10								15			
Ala Ala Ala Val Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala																			
			20				25									30			
Ala Ile Gly Asp Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly																			
			35				40									45			
Asp Ala Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val																			
			50				55									60			
Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala																			
			65				70									75			80
Gly Asp Ala Gly Glu Gly Asn Lys Asp Ala Gly Lys Leu Phe Val Lys																			
							85									90			95
Lys Asn Ala Gly Asp Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala																			
			100													105			110
Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile																			
			115													120			125
Val Asp Ala Ala Glu Gly Asp Asp Lys Gln Gly Lys Lys Ala Ala Asp																			
			130													135			140
Ala Thr Asn Pro Ile Glu Ala Ala Ile Gly Gly Ala Asp Ala Gly Ala																			
							150									155			160
Asn Ala Glu Ala Phe Asn Lys Met Lys Lys Asp Asp Gln Ile Ala Ala																			
							165									170			175
Ala Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys																			
							180									185			190
Asp Asp Ala Ala Ala His																			
							195												

<210> 37  
 <211> 573  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS

&lt;222&gt; (1) .. (573)

&lt;400&gt; 37

gaa ggg act gtt aag aat gct gtt gat atg gca aag gcc gct gcg gaa 48  
 Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Glu  
 1 5 10 15

gct gca agt gct gca agt gct gct act ggt agt aca acg att gga gat 96  
 Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp  
 20 25 30

gtt gtt aag agt ggt gag gca aaa gat ggt gat gcg gcg agt gtt aat 144  
 Val Val Lys Ser Gly Glu Ala Lys Asp Gly Asp Ala Ala Ser Val Asn  
 35 40 45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct 192  
 Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala  
 50 55 60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt acg act 240  
 Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Thr Thr  
 65 70 75 80

aac gtg aat gtt ggg aag ttg ttt gtg aag aat aat ggt aat gag ggt 288  
 Asn Val Asn Val Gly Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly  
 85 90 95

ggt gat gca agt gat gct ggg aaa gct gct gct gcg gtt gct gct gtt 336  
 Gly Asp Ala Ser Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val  
 100 105 110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt 384  
 Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
 115 120 125

gat aag cag ggg gtt act gat gta aag gat gct aca aat ccg att gag 432  
 Asp Lys Gln Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Glu  
 130 135 140

gcg gct att ggg ggt aca aat gat aat gat gct gcg gcg ttt gct act 480  
 Ala Ala Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr  
 145 150 155 160

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg 528  
 Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met  
 165 170 175

gct aag gat ggg cag ttt gct ttg aag gat gat gct gct aag gat 573  
 Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp Asp Ala Ala Lys Asp  
 180 185 190

&lt;210&gt; 38

&lt;211&gt; 191

&lt;212&gt; PRT

&lt;213&gt; Borrelia garinii

&lt;400&gt; 38

Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Ala Glu  
 1 5 10 15

Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp  
 20 25 30

Val Val Lys Ser Gly Glu Ala Lys Asp Gly Asp Ala Ala Ser Val Asn  
           35                          40                          45  
 Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala  
           50                          55                          60  
 Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Thr Thr  
           65                          70                          75                          80  
 Asn Val Asn Val Gly Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly  
                           85                          90                          95  
 Gly Asp Ala Ser Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val  
                           100                          105                          110  
 Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
           115                          120                          125  
 Asp Lys Gln Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Glu  
           130                          135                          140  
 Ala Ala Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr  
           145                          150                          155                          160  
 Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met  
                           165                          170                          175  
 Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp Asp Ala Ala Lys Asp  
           180                          185                          190

<210> 39  
 <211> 189  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1) .. (189)

<400> 39  
 ggt gat aaa acg ggg gtt gct gcg gat gct gaa aat ccg att gac gcg 48  
 Gly Asp Lys Thr Gly Val Ala Ala Asp Ala Glu Asn Pro Ile Asp Ala  
           1                          5                          10                          15  
 gct att ggg ggt gcg gat gct gat gct gcg gcg ttt aat aag gag ggg 96  
 Ala Ile Gly Gly Ala Asp Ala Asp Ala Ala Ala Phe Asn Lys Glu Gly  
                           20                          25                          30  
 atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg 144  
 Met Lys Lys Asp Asp Gln Ile Ala Ala Met Val Leu Arg Gly Met  
           35                          40                          45  
 gct aag gat ggg cag ttt gct ttg acg aat aat gct gct gct cat 189  
 Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn Ala Ala Ala His  
           50                          55                          60

<210> 40  
 <211> 63  
 <212> PRT  
 <213> *Borrelia garinii*

&lt;400&gt; 40

Gly Asp Lys Thr Gly Val Ala Ala Asp Ala Glu Asn Pro Ile Asp Ala  
 1 5 10 15

Ala Ile Gly Gly Ala Asp Ala Asp Ala Ala Ala Phe Asn Lys Glu Gly  
 20 25 30

Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met  
 35 40 45

Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn Ala Ala Ala His  
 50 55 60

&lt;210&gt; 41

&lt;211&gt; 576

&lt;212&gt; DNA

<213> *Borrelia garinii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1) .. (576)

&lt;400&gt; 41

gaa ggg act gtt aag aat gct gtt gat atg gca aaa gct gct gcg gtt 48  
 Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val  
 1 5 10 15

gct gca agt gct gct act ggc aat gca gca att ggg gat gtt gtt aag 96  
 Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys  
 20 25 30

agt aat ggt gga gca gca gca aaa ggt ggt gat gcg gcg agt gtt aat 144  
 Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn  
 35 40 45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct 192  
 Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala  
 50 55 60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt gaa act 240  
 Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr  
 65 70 75 80

aac aag gat gct ggg aag ttg ttt gtg aag aag aat ggt gat gat ggt 288  
 Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly  
 85 90 95

ggt gat gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt 336  
 Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val  
 100 105 110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt 384  
 Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
 115 120 125

gat aag acg ggg gtt act gat gta aag gat gct aca aat ccg att gac 432  
 Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp  
 130 135 140

gcg gct att ggg ggg agt gcg gat gct aat gct gag gcg ttt gat aag 480  
 Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys  
 145 150 155 160

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg 528  
 Met Lys Lys Asp Asp Gln Ile Ala Ala Met Val Leu Arg Gly Met  
                   165                                  170                  175

gct aag gat ggg cag ttt gct ttg aag aat aat gat cat gat aat cat 576  
 Ala Lys Asp Gly Gln Phe Ala Leu Lys Asn Asn Asp His Asp Asn His  
                   180                                  185                  190

<210> 42

<211> 192

<212> PRT

<213> *Borrelia garinii*

<400> 42

Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val  
   1                  5                                  10                                  15

Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys  
                   20                                  25                                  30

Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn  
                   35                                  40                                  45

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala  
                   50                                  55                                  60

Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr  
   65                                  70                                  75                                  80

Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly  
                   85                                  90                                  95

Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Val  
                   100                                  105                                  110

Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
                   115                                  120                                  125

Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp  
                   130                                  135                                  140

Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys  
   145                                  150                                  155                                  160

Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met  
                   165                                  170                                  175

Ala Lys Asp Gly Gln Phe Ala Leu Lys Asn Asn Asp His Asp Asn His  
                   180                                  185                                  190

<210> 43

<211> 336

<212> DNA

<213> *Borrelia garinii*

<220>

<221> CDS

<222> (1)..(336)

<400> 43

aag ggg act gtt aag aat gct gtt gat atg gca aag gcc gct gag gaa 48

```

Lys Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Glu Glu
 1          5          10          15
gct gca agt gct gca agt gct gct act ggt aat gca gcg att ggg gat 96
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp
          20          25          30
gtt gtt aag aat agt ggg gca gca gca aaa ggt ggt gag gcg gcg agt 144
Val Val Lys Asn Ser Gly Ala Ala Ala Lys Gly Gly Glu Ala Ala Ser
          35          40          45
gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gga 192
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
          50          55          60
aag gct gat gcg aag gaa ggg aag ttg gat gct act ggt gct gag ggt 240
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Ala Thr Gly Ala Glu Gly
          65          70          75          80
acg act aac gtg aat gct ggg aag ttg ttt gtg aag agg gcg gct gat 288
Thr Thr Asn Val Asn Ala Gly Lys Leu Phe Val Lys Arg Ala Ala Asp
          85          90          95
gat ggt ggt gat gca gat gat gct ggg aag gct gct gct gcg gtt gct 336
Asp Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Val Ala
          100          105          110

```

<210> 44  
 <211> 112  
 <212> PRT  
 <213> *Borrelia garinii*

```

<400> 44
Lys Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Glu Glu
 1          5          10          15
Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp
          20          25          30
Val Val Lys Asn Ser Gly Ala Ala Ala Lys Gly Gly Glu Ala Ala Ser
          35          40          45
Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
          50          55          60
Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Ala Thr Gly Ala Glu Gly
          65          70          75          80
Thr Thr Asn Val Asn Ala Gly Lys Leu Phe Val Lys Arg Ala Ala Asp
          85          90          95
Asp Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Val Ala
          100          105          110

```

<210> 45  
 <211> 522  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(522)

<400> 45  
gca agt gct gct act ggt aat gca gcg att gga gat gtt gtt aat ggt 48  
Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly  
1 5 10 15

gat gtg gca aaa gca aaa ggt ggt gat gcg gcg agt gtt aat ggg att 96  
Asp Val Ala Lys Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile  
20 25 30

gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct gat gcg 144  
Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala  
35 40 45

aag gaa ggg aag ttg aat gct gct ggt gct gag ggt acg act aac gcg 192  
Lys Glu Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala  
50 55 60

gat gct ggg aag ttg ttt gtg aag aat gct ggt aat gtg ggt ggt gaa 240  
Asp Ala Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu  
65 70 75 80

gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg 288  
Ala Gly Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly  
85 90 95

gag cag ata tta aaa gcg att gtt gat gct gct aag gat ggt ggt gag 336  
Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu  
100 105 110

aag cag ggt aag aag gct gcg gat gct aca aat ccg att gac gcg gct 384  
Lys Gln Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Asp Ala Ala  
115 120 125

att ggg ggt aca aat gat aat gat gct gct gcg gcg ttt gct act atg 432  
Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met  
130 135 140

aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg gct 480  
Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala  
145 150 155 160

aag gat ggg caa ttt gct ttg aag gat gct gct gct gct cat 522  
Lys Asp Gly Gln Phe Ala Leu Lys Asp Ala Ala Ala Ala His  
165 170

<210> 46  
<211> 174  
<212> PRT  
<213> Borrelia garinii

<400> 46  
Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly  
1 5 10 15

Asp Val Ala Lys Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile  
20 25 30

Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala  
35 40 45

Lys Glu Gly Lys Leu Asn Ala Ala Gly Ala Glu Gly Thr Thr Asn Ala  
50 55 60

Asp Ala Gly Lys Leu Phe Val Lys Asn Ala Gly Asn Val Gly Gly Glu  
 65 70 75 80  
 Ala Gly Asp Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly  
 85 90 95  
 Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly Gly Glu  
 100 105 110  
 Lys Gln Gly Lys Lys Ala Ala Asp Ala Thr Asn Pro Ile Asp Ala Ala  
 115 120 125  
 Ile Gly Gly Thr Asn Asp Asn Asp Ala Ala Ala Phe Ala Thr Met  
 130 135 140  
 Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala  
 145 150 155 160  
 Lys Asp Gly Gln Phe Ala Leu Lys Asp Ala Ala Ala Ala His  
 165 170

<210> 47  
 <211> 585  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(585)

<400> 47  
 gaa ggg act gtt aag aat gct gtt gat ata ata aag gct gct gcg gaa 48  
 Glu Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Ala Ala Glu  
 1 5 10 15  
 gct gca agt gct gca agt gct gct act ggt agt gca gca att ggg gat 96  
 Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Ala Ala Ile Gly Asp  
 20 25 30  
 gtt gtt aat ggt aat gga gca aca gca aaa ggt ggt gat gcg aag agt 144  
 Val Val Asn Gly Asn Gly Ala Thr Ala Lys Gly Gly Asp Ala Lys Ser  
 35 40 45  
 gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag 192  
 Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu  
 50 55 60  
 aag gct gat gcg aag gaa ggg aag ttg gat gtg gct ggt gat gct ggt 240  
 Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala Gly  
 65 70 75 80  
 gaa act aac aag gat gct ggg aag ttg ttt gtg aag aac aat ggt aat 288  
 Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Asn Asn Gly Asn  
 85 90 95  
 gag ggt ggt gat gca gat gat gct ggg aag gct gct gct gcg gtt gct 336  
 Glu Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Ala Val Ala  
 100 105 110  
 gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aag 384  
 Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys  
 115 120 125



```

ggt ggt gat aag acg ggt aag aat aat gtg aag gat gct gaa aat ccg      432
Gly Gly Asp Lys Thr Gly Lys Asn Asn Val Lys Asp Ala Glu Asn Pro
130                               135                               140

att gag gcg gct att ggg agt agt gcg gat gct gat gct gcg gcg ttt      480
Ile Glu Ala Ala Ile Gly Ser Ser Ala Asp Ala Asp Ala Ala Ala Phe
145                               150                               155                               160

aat aag gag ggg atg aag aag gat gat cag att gct gct gct atg gtt      528
Asn Lys Glu Gly Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val
165                               170                               175

ctg agg gga atg gct aag gat ggg cag ttt gct ttg acg aat gat gct      576
Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asp Ala
180                               185                               190

gct gct cat
Ala Ala His
195

```

585

```

<210> 48
<211> 195
<212> PRT
<213> Borrelia garinii

```

```

<400> 48
Glu Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Ala Ala Glu
1      5      10      15

Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Ser Ala Ala Ile Gly Asp
20      25      30

Val Val Asn Gly Asn Gly Ala Thr Ala Lys Gly Gly Asp Ala Lys Ser
35      40      45

Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu
50      55      60

Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Asp Ala Gly
65      70      75      80

Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Asn Asn Gly Asn
85      90      95

Glu Gly Gly Asp Ala Asp Asp Ala Gly Lys Ala Ala Ala Ala Val Ala
100     105     110

Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys
115     120     125

Gly Gly Asp Lys Thr Gly Lys Asn Asn Val Lys Asp Ala Glu Asn Pro
130     135     140

Ile Glu Ala Ala Ile Gly Ser Ser Ala Asp Ala Asp Ala Ala Ala Phe
145     150     155     160

Asn Lys Glu Gly Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val
165     170     175

Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asp Ala
180     185     190

```

Ala Ala His  
195

<210> 49  
<211> 591  
<212> DNA  
<213> *Borrelia garinii*

<220>  
<221> CDS  
<222> (1)..(591)

<400> 49

gaa ggg act gtt aag aat gct gtt ggg agt gca aca aat aag acc gtt	48
Glu Gly Thr Val Lys Asn Ala Val Gly Ser Ala Thr Asn Lys Thr Val	
1 5 10 15	
gtt gct ttg gct aac ttg gtt cga aag acc gtg caa gct ggg ttg aag	96
Val Ala Leu Ala Asn Leu Val Arg Lys Thr Val Gln Ala Gly Leu Lys	
20 25 30	
aag gtt ggg gat gtt gtt aag aat agt gag gca aaa gat ggt gat gcg	144
Lys Val Gly Asp Val Val Lys Asn Ser Glu Ala Lys Asp Gly Asp Ala	
35 40 45	
gcg agt gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct	192
Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala	
50 55 60	
gct gag aag gct gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct	240
Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala	
65 70 75 80	
gct ggt gaa act aac aag gat gct ggg aag ttg ttt gtg aag aag aat	288
Ala Gly Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn	
85 90 95	
aat gag ggt ggt gaa gca aat gat gct ggg aag gct gct gct gcg gtt	336
Asn Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala Ala Ala Ala Val	
100 105 110	
gct gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct gct	384
Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala	
115 120 125	
aag gat ggt gat gat aag cag ggt aag aag gct gag gat gct aca aat	432
Lys Asp Gly Asp Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Thr Asn	
130 135 140	
ccg att gac gcg gct att ggg ggt gca ggt gcg ggt gct aat gct gct	480
Pro Ile Asp Ala Ala Ile Gly Gly Ala Gly Ala Gly Ala Asn Ala Ala	
145 150 155 160	
gcg gcg ttt aat aat atg aag aag gat gat cag att gct gct gct atg	528
Ala Ala Phe Asn Asn Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met	
165 170 175	
gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg acg aat aat	576
Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn	
180 185 190	
gct cat act aat cat	591
Ala His Thr Asn His	

195

<210> 50  
 <211> 197  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 50  
 Glu Gly Thr Val Lys Asn Ala Val Gly Ser Ala Thr Asn Lys Thr Val  
 1 5 10 15  
 Val Ala Leu Ala Asn Leu Val Arg Lys Thr Val Gln Ala Gly Leu Lys  
 20 25 30  
 Lys Val Gly Asp Val Val Lys Asn Ser Glu Ala Lys Asp Gly Asp Ala  
 35 40 45  
 Ala Ser Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala  
 50 55 60  
 Ala Glu Lys Ala Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala  
 65 70 75 80  
 Ala Gly Glu Thr Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn  
 85 90 95  
 Asn Glu Gly Gly Glu Ala Asn Asp Ala Gly Lys Ala Ala Ala Val  
 100 105 110  
 Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala  
 115 120 125  
 Lys Asp Gly Asp Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Thr Asn  
 130 135 140  
 Pro Ile Asp Ala Ala Ile Gly Gly Ala Gly Ala Gly Ala Asn Ala Ala  
 145 150 155 160  
 Ala Ala Phe Asn Asn Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met  
 165 170 175  
 Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Thr Asn Asn  
 180 185 190  
 Ala His Thr Asn His  
 195

<210> 51  
 <211> 594  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(594)

<400> 51  
 aag ggg act gtt aag aat gct gtt gat atg aca aaa gct gct gcg gtt 48  
 Lys Gly Thr Val Lys Asn Ala Val Asp Met Thr Lys Ala Ala Ala Val  
 1 5 10 15  
 gct gca agt gct gca agt gct gct act ggt aat gca gca att ggg gat 96

Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp  
 20 25 30  
 gtt gtt aat ggt aat gat gga gca gca aaa ggt ggt gat gcg gcg agt 144  
 Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser  
 35 40 45  
 gtt aat ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag 192  
 Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu  
 50 55 60  
 aag gct gat gcg aag gaa ggg aag ttg aat gtg gct ggt gct gct ggt 240  
 Lys Ala Asp Ala Lys Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly  
 65 70 75 80  
 gct gag ggt aac gag gct gct ggg aag ctg ttt gtg aag aag aat gct 288  
 Ala Glu Gly Asn Glu Ala Ala Gly Lys Leu Phe Val Lys Lys Asn Ala  
 85 90 95  
 ggt gat cat ggt ggt gaa gca ggt gat gct ggg agg gct gct gct gcg 336  
 Gly Asp His Gly Gly Glu Ala Gly Asp Ala Gly Arg Ala Ala Ala Ala  
 100 105 110  
 gtt gct gct gtt agt ggg gag cag ata tta aaa gcg att gtt gat gct 384  
 Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala  
 115 120 125  
 gct aag gat ggt ggt gat aag cag ggt aag aag gct gag gat gct gaa 432  
 Ala Lys Asp Gly Gly Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Glu  
 130 135 140  
 aat ccg att gac gcg gct att ggg agt acg ggt gcg gat gat aat gct 480  
 Asn Pro Ile Asp Ala Ala Ile Gly Ser Thr Gly Ala Asp Asp Asn Ala  
 145 150 155 160  
 gct gag gcg ttt gct act atg aag aag gat gat cag att gct gct gct 528  
 Ala Glu Ala Phe Ala Thr Met Lys Lys Asp Asp Gln Ile Ala Ala Ala  
 165 170 175  
 atg gtt ctg agg gga atg gct aag gat ggg cag ttt gct ttg aag gat 576  
 Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp  
 180 185 190  
 gct gct cat gat aat cat 594  
 Ala Ala His Asp Asn His  
 195

&lt;210&gt; 52

&lt;211&gt; 198

&lt;212&gt; PRT

&lt;213&gt; Borrelia garinii

&lt;400&gt; 52

Lys Gly Thr Val Lys Asn Ala Val Asp Met Thr Lys Ala Ala Ala Val  
 1 5 10 15

Ala Ala Ser Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp  
 20 25 30

Val Val Asn Gly Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser  
 35 40 45

Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu

50                      55                      60  
 Lys Ala Asp Ala Lys Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly  
 65                      70                      75                      80  
 Ala Glu Gly Asn Glu Ala Ala Gly Lys Leu Phe Val Lys Lys Asn Ala  
 85                      90                      95  
 Gly Asp His Gly Gly Glu Ala Gly Asp Ala Gly Arg Ala Ala Ala Ala  
 100                      105                      110  
 Val Ala Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala  
 115                      120                      125  
 Ala Lys Asp Gly Gly Asp Lys Gln Gly Lys Lys Ala Glu Asp Ala Glu  
 130                      135                      140  
 Asn Pro Ile Asp Ala Ala Ile Gly Ser Thr Gly Ala Asp Asp Asn Ala  
 145                      150                      155                      160  
 Ala Glu Ala Phe Ala Thr Met Lys Lys Asp Asp Gln Ile Ala Ala Ala  
 165                      170                      175  
 Met Val Leu Arg Gly Met Ala Lys Asp Gly Gln Phe Ala Leu Lys Asp  
 180                      185                      190  
 Ala Ala His Asp Asn His  
 195

<210> 53  
 <211> 261  
 <212> DNA  
 <213> Borrelia garinii

<220>  
 <221> CDS  
 <222> (1) .. (261)

<400> 53  
 aag ggg act gtt aag aat gct gtt gat ata ata aag gct act gcg gtt 48  
 Lys Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Thr Ala Val  
 1                      5                      10                      15  
 gct gca agt gct gct act ggt agt aca acg att ggg gat gtt gtt aag 96  
 Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp Val Val Lys  
 20                      25                      30  
 aat ggt gag gca aaa ggt ggt gag gcg aag agt gtt aat ggg att gct 144  
 Asn Gly Glu Ala Lys Gly Gly Glu Ala Lys Ser Val Asn Gly Ile Ala  
 35                      40                      45  
 aag ggg ata aag ggg att gtt gat gct gct gga aag gct gat gcg aag 192  
 Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala Asp Ala Lys  
 50                      55                      60  
 gaa ggg aag ttg aat gtg gct ggt gct gct ggt gag ggt aac gag gct 240  
 Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly Glu Gly Asn Glu Ala  
 65                      70                      75                      80  
 gct ggg aag ctg ttt gtg taa 261  
 Ala Gly Lys Leu Phe Val  
 85

<210> 54  
 <211> 86  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 54  
 Lys Gly Thr Val Lys Asn Ala Val Asp Ile Ile Lys Ala Thr Ala Val  
 1 5 10 15  
 Ala Ala Ser Ala Ala Thr Gly Ser Thr Thr Ile Gly Asp Val Val Lys  
 20 25 30  
 Asn Gly Glu Ala Lys Gly Gly Glu Ala Lys Ser Val Asn Gly Ile Ala  
 35 40 45  
 Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala Asp Ala Lys  
 50 55 60  
 Glu Gly Lys Leu Asn Val Ala Gly Ala Ala Gly Glu Gly Asn Glu Ala  
 65 70 75 80  
 Ala Gly Lys Leu Phe Val  
 85

<210> 55  
 <211> 213  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(213)

<400> 55  
 gta aat tac tat agg att aga act agt gta cga tat gag tcc ttt ggt 48  
 Val Asn Tyr Tyr Arg Ile Arg Thr Ser Val Arg Tyr Glu Ser Phe Gly  
 1 5 10 15  
 tat ttt gca gct gct aat gaa ttt gaa ata agt gaa gtt aaa att gcg 96  
 Tyr Phe Ala Ala Ala Asn Glu Phe Glu Ile Ser Glu Val Lys Ile Ala  
 20 25 30  
 gat gtt aat gga aca cat ttt att gct aca aaa gag aaa gaa ata tta 144  
 Asp Val Asn Gly Thr His Phe Ile Ala Thr Lys Glu Lys Glu Ile Leu  
 35 40 45  
 tat gat tca ctt gat tta agg gct cgt gga aaa ata ttt gaa ata act 192  
 Tyr Asp Ser Leu Asp Leu Arg Ala Arg Gly Lys Ile Phe Glu Ile Thr  
 50 55 60  
 tca aag cga atg ttt aag ctt 213  
 Ser Lys Arg Met Phe Lys Leu  
 65 70

<210> 56  
 <211> 71  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 56  
 Val Asn Tyr Tyr Arg Ile Arg Thr Ser Val Arg Tyr Glu Ser Phe Gly  
 1 5 10 15  
 Tyr Phe Ala Ala Ala Asn Glu Phe Glu Ile Ser Glu Val Lys Ile Ala  
 20 25 30

Asp Val Asn Gly Thr His Phe Ile Ala Thr Lys Glu Lys Glu Ile Leu  
 35 40 45

Tyr Asp Ser Leu Asp Leu Arg Ala Arg Gly Lys Ile Phe Glu Ile Thr  
 50 55 60

Ser Lys Arg Met Phe Lys Leu  
 65 70

<210> 57

<211> 8762

<212> DNA

<213> *Borrelia afzelii*

<400> 57

```

gagagtgtctg ttgatgggggt tagcaagtgg ttagaagaga tgataaaagc tgctaaggag 60
gctgctacaa aggggtggtac tgggtggtgg agcgaaaaga ttggggatgt tgggtgctgct 120
aataatcagg gtgctgtagc tgataaggac agtggttaagg ggattgcgaa ggggataaaag 180
gggattgttg atgctgctgg gaaggctttt ggtaaggatg gtaatgcgct gacagggtgta 240
aaagaagttg ctgatgaggg tggggctaac gaggatgcgg ggaagttggt tgctggtaaat 300
gctggtaatg ctgctgctgc tgacattgcg aaggcggctg gtgctgttac tgcgggttagt 360
ggggagcaga tactgaaagc tattgttgat ggtgctggtg gtgctgctca agatggtaaa 420
aaggctgcgg aggctaagaa tccgattgca gctgcgattg gggctgatgc tgctgggtgcg 480
gatttttggtg atgatatgaa gaagagtgat aagattgctg cggctattgt tttgaggggg 540
gtggctaaga gtggaagtt tgctgttgct aatgctgcta agaaggagag tgtgaagagt 600
gctgtggaga gtgctgttga tgagggttagc aagtgggttag aagagatgat aaaagctgct 660
ggtggggctg ctaagggtgg tactgggtgg aataacgaaa agattgggga ttctgataat 720
aataagggtg ctgtagctga taaggacagt gtttaaggga ttgcgaaggg gataaagggg 780
attgttgatg ctgctgggaa ggcttttggg aaggatggtg atgcgctgaa ggatgttgca 840
aaagttgctg atgatgcggc tggggctaac gcgaatgcag ggaagttggt tgctggtaaat 900
gctgctgggtg gtgcccgtga tgctgatgat gctaacattg cgaaggcggc tgggtgctggt 960
agtgcggtta gtggggagca gatactgaaa gctattgttg atgctgctgg tgctgctgct 1020
aatcaggatg gtaagaaggc tgcggatgct aagaatccga ttgcagctgc gattgggact 1080
aatgatgatg gggcggagtt taaggatgga atgaagaaga gtgataatat tgctgcagct 1140
attgttttga ggggggtggc taagggtgga aagtttgctg ttgctaatagc tgctaattgat 1200
agtaaggcga gtgtgaagag tgctgtggag agtgctgttg atgaggttag caagtgggta 1260
gaagagatga taacagctgc tggtgaggct gctacaaagg gtggtgatgc tgggtggtggt 1320
gctgataaga ttggggatgt tgggtgctgct aataatgggt ctgtagctga tgcgagcagt 1380
gttaaggaga ttgcgaaggg gataaagggg attgttgatg ctgctgggaa ggcttttggc 1440
aaggatggta atgcgctgaa ggatgttgca gaagttgctg atgataagaa ggaggcgggg 1500
aagttgtttg ctggtaatgc tgggtggtgct gtgctgctg atgctgcgat tgggaaggcg 1560
gctggtgctg ttactgcggg tagtggggag cagatactga aagctattgt tgatgctgct 1620
ggtggtgcgg atcaggcggg taagaaggct gatgcggcta agaatccgat tgcagctgcg 1680
attggggctg atgctgctgg tgctgggtgcg gatttttggt atgatatgaa gaagagaaat 1740
gataagattg ttgctggctat tgttttgagg ggggtggcta aggatggaaa gtttgctgct 1800
gctgctaata atgataatag taaggcgagt gtgaagagt ctgtggagag tgctgttgat 1860
gaggttagca agtgggttaga agagatgata acagctgctg atggggctgc taaagggtgg 1920
actggtggta atagcgaaaa gattggggat gctggtgata ataataatgg tgctgtagct 1980
gatgagaaca gtgttaagga gattgcaaag gggataaagg ggattgttgc ggctgctggg 2040
aaggcttttg gcaaggatgg caaggatggt gatgcgctga aggatgttga aacagttgct 2100
gctgagaatg aggctaacaa ggatgcgggg aagttgtttg ctggtgctaa tggtaatgct 2160
ggtgctgctg ttggtgacat tgcgaaggcg gctgctgctg ttactgcggg tagtggggag 2220
cagatactaa aagctattgt tgatgctgct ggtgatgcgg atcaggcggg taaggaggtc 2280
gctgaggcta agaatccgat tgcagctgcg attggggcta atgctgctga taatgcggcg 2340
gcgttttggt aggatgagat gaagaagagt gataagattg ctgcagctat tgttttgagg 2400
ggggtggcta aggatggaaa gtttgctggt gctaattgcta atgatgataa gaaggcgagt 2460
gtgaagagtg ctgtggagag tgctgtggat gaggttagca agtggttaga agagatgata 2520
acagctgcta aggaagctgc tacaaagggt ggtactgggt gtaataacga aaagattgga 2580
gattctgatg ctaataatgg tgcaaggct gtgttaatgg gattgcgaat 2640
gggataaagg ggattgttga tgctgctggg aaggcttttg gcaaggaggg tagtgcgctg 2700
aaggatgtta aaacagttgc tgctgagaat gaggctaaca aggatgcggg gaagttgttt 2760
gctggtaaga atggtaatgc tgatgctgct gatgctgctg acattgcgaa ggcggctggt 2820

```

gctgttagtg	cggtttagtg	ggagcagata	ctgaaagcta	ttgttgatgg	tgctgggtgat	2880
gcagctaatac	agggcgggtaa	aaaggctgct	gaggctaaga	atccgattgc	ggctgcgatt	2940
gggactaatg	aagctggggc	ggagtttggg	gatgatatga	agaagagaaa	tgataagatt	3000
gctgcggtcta	ttgttttgag	gggggtggct	aaggatggaa	agtttgctgt	tgctaattgct	3060
gctgctgata	atagtaaggc	gagtgtgaag	agtgtgttg	atgaggttag	caagtgggta	3120
gaagagatga	taaaggctgc	tggtgaggct	gctacaaagg	gtggtgatgc	tggtgggtgg	3180
gctgataaga	ttggggatgc	tggtgataag	gggtgctgtg	ctgatgctgag	cagtgttaag	3240
gagattgcga	atgggataaa	ggggattggt	gatgctgctg	ggaaggcctt	tggcaaggag	3300
ggtagtgctg	tgaaggatgt	taaaacagtt	gctgctgaga	atgaggctaa	caaggatgct	3360
gggaagtgtg	ttgctggtaa	tgctggtaat	gggtgctgtg	atgacattgc	gaaggcggct	3420
gctgctgtta	ctgctggttag	tggggagcag	atactgaaag	ctattgttga	tgctgctggg	3480
gataaggcta	atcaggatgg	taaaaaggct	gctgctgata	agaatccgat	tgctgctgctg	3540
attggggctg	ctgatgctgg	tgctgctggc	gctgttaata	agaatgatata	gaagaagagt	3600
gataagattg	ctgcagctat	tggtttgagg	gggtggtgta	aggatggaaa	gtttgctgct	3660
gctgctgctg	atgctaataa	tagtaaggcg	agcgtgaaga	gtgctgttgg	tgagggttagc	3720
aagtgggttag	aagagatgat	aaaagctgct	gctgaggtctg	caaaagtggg	tggtactggg	3780
ggtagcgaaa	agattgggga	tgctgataat	aataagggtg	ctgtagctga	tgctgctggg	3840
gttaattggga	ttgcgaatgg	gataaagggg	attgttgatg	ctgctgggaa	ggcttttggg	3900
aaggatgggtg	cgctggcagg	tggtgcagct	gctgctgaga	atgatgataa	gaaggatgctg	3960
gggaagtgtg	ttgctggtaa	gaatgggtgg	ctggtgctg	ctgatgctgat	tgggaaggcg	4020
gctgctgctg	ttactggcgt	tagtggggag	cagatctga	aagctattgt	tgatgctgct	4080
gggtgctgcag	ctaatacaggc	gggtaaaaag	gctgctggatg	ctaagaatcc	gattgctggg	4140
gcgattggga	ctgctgatga	tggggcggag	tttaaggatg	atatgaagaa	gagtataaat	4200
attgctgctg	ctattgtttt	gaggggggtg	gctaaggatg	gaaagtgtgc	tggtgctaat	4260
gctgatgata	ataaggcgag	tgtaagagt	gctgtggaga	gtgctgttga	tgagggttagc	4320
aagtgggttag	aagagatgat	aacagctgct	ggtaggctg	caaaagtggg	tgctgggtgg	4380
gggtgctgata	agattgggga	tgctgctaata	aaggatgggtg	cgaaggctga	tgagagcagt	4440
gttaattggaa	ttgcaaaggg	gataaagggg	attgttgatg	ctgctgggaa	ggcttttggc	4500
aaggaggggta	gtgcgctgaa	ggatgttgca	aaagttgctg	atgatgataa	caaggatgctg	4560
gggaagtgtg	ttgctggtaa	tgctgggtgg	gggtgctggg	ctgatattgc	gaaggcggct	4620
gctgctgtta	ctgctggttag	tggggagcag	atactgaaag	ctattgttga	tgctgctggg	4680
gctgcggatc	agggcgggtg	agctgctggg	gctgctaaaga	atccgattgc	ggctgctgatt	4740
ggggctgatg	ctggctgctg	ggaggagttt	gagagtgaga	tgaagaagag	tgataagatt	4800
gctgctgctg	ttgttttgag	gggggtggct	aagggtggaa	agtttgctgt	tgctgctaat	4860
gatgctgcaa	atgtgaagag	tgctgtggag	agtgtgttg	gtgagggttag	cgcatgggta	4920
gaagagatga	taacagctgc	tagtgaggct	gctacaaagg	gtggtactgg	tggtactggg	4980
gggtgatagt	aaaagattgg	ggattctgat	gctaataatg	gtgctgtagc	tgatgctgagc	5040
agtgttaagg	agattgcgaa	ggggataaag	gggattgttg	atgctgctgg	gaaggccttt	5100
ggtaaggatg	gtaatgcgct	gaaggatggt	gcagaagttg	ctgatgatga	ggctaacgctg	5160
gatgcgggga	agttgtttgc	tggtaatgct	ggtaatgctg	ctgctgctga	cgttgcgaa	5220
gcggctgggtg	ctgttactgc	gggttagtggt	gagcagatac	tgaaagctat	tggtgatgct	5280
gctgggtgctg	cggatcaggc	gggtgcaaag	gctgatgcgg	ctaagaatcc	gattgctgct	5340
gcgattggga	ctaatagaagc	tggggcggcg	tttaaggatg	gaatgaagaa	gagaaatgat	5400
aatattgctg	cggctattgt	tttgaggggg	gtggctaaga	gtggaaagt	tgctgttgc	5460
gctgctgatg	ctggtaaggc	gagagtgtga	agagtgtgtg	ggagagtgtg	gttgatgagg	5520
ttagcaagtg	gttagaagag	atgataacag	ctgctagtga	ggctgcaaaa	gttggtgctg	5580
gtggtgatga	taagattggg	gattctgcta	ataatgggtg	tgtagctgat	gcgggcagtg	5640
ttaagggaat	tgcaaggggg	ataaagggga	ttgttgatgc	tgctgggaag	gcttttggta	5700
aggaggggtga	tgctgctgaag	gatgttgcaa	aagttgctga	tgagaatggg	gataacaagg	5760
atgcggggaa	gttggttgc	gggtgagaatg	gtaagtctgg	tggtgctgct	gatgctgaca	5820
ttgcgaaggc	ggctgctgct	gttactgcgg	ttagtgggga	gcagatactg	aaagctattg	5880
ttgaggctgc	tggtgctggg	gatgcagcta	atcaggcggg	taagaaggct	gatgaggcta	5940
agaatccgat	tgctgctgctg	attgggactg	atgatgctgg	ggcggcgttt	ggtcaggatg	6000
atatgaagaa	gagaaatgat	aatattgctg	cggctattgt	tttgaggggg	gtggctaagg	6060
tggaagagtt	tgctgttgc	aatgctgcta	atgataagaa	ggcagagtgtg	aagagtgtg	6120
tggaagagtt	tggtgatgag	gttagcaagt	ggtagaaga	gataataaca	gctactggga	6180
aggcttttgg	taaggatggg	aatgcgctgg	cagggtgtgc	aaaagtgtgc	gatgatgagg	6240
ctaacgcgga	tgctggggaag	ttgtttgctg	gtgagaatgg	taatgctggg	gctgctgcca	6300
ttgggaaggc	ggctgctgct	gttactgcgg	ttagtgggga	gcagatactg	aaagctattg	6360
ttgatgctgc	tggtgggtgcg	gctcaggtgg	gtgctgggtg	tggtgcggct	acgaatccga	6420
ttgcagctgc	gattggggctg	gctgggtgatg	gtgcggattt	tggttaaggat	gagatgaaga	6480
agagaatga	taagattgct	gcggctattg	ttttgagggg	gggtggctaag	gatggaaagt	6540
ttgctgctgc	tgctaataatg	agtaaggcga	gtgtgaagag	tgctgtggag	agtgtgttg	6600
atgaggttag	caagtgggta	gaagagatga	taacagctgc	tgatgctgct	gctgctaaag	6660



```

ttggcgatgc tgggtgggtggt gctgataaga ttgggggatgt tgggtgctgct aataaggggtg 6720
cgaaggctga tgcgagcagt gttaaggaga ttgcgaaggg gataaagggg attgttgatg 6780
ctgctgggaa ggcttttggg ggtgatgcgc tgaaggatgt taaagctgct ggtgatgata 6840
acaaggaggc aggggaagttg tttgctgggtg ctaatggtaa tgctgggtgct aatgctgctg 6900
ctgctgatga cattgcgaag gcggctgggtg ctgttagtgc ggttagtggg gagcagatac 6960
tgaaagctat tgttgaggcg gctgggtgctg cggatcaggc ggggtgtaaag gctgaggagg 7020
ctaagaatcc gattgcagct gcgattggga ctgatgatgc tgggtgcggcg gagtttgggtg 7080
agaatgatat gaagaagaat gataaatattg ctgcggctat tgttttgagg ggggtggcta 7140
agagtggaaa gtttgcctgct aatgctaattg atgctggtaa gaaggagagt gtgaagagtg 7200
ctgtggatga ggctagcaag tgggttagaag agatgataac agctgctggg gaggctgcta 7260
caaaggggtg tactggtgaa gctagcgaaa agattgggga tgttgggtgat aataatcatg 7320
gtgctgtagc gatgaagaag agtggttaagg ggattgcgaa ggggataaag gggattgttg 7380
attgctgctgg gaaggctttt ggtaaggatg gtgcgctgaa ggatgttgca gctgctgctg 7440
gtgatgaggc taacaaggat gcggggaaagt tgtttgctgg tcaggatggg ggtgggtgctg 7500
atgggtgacat tgcgaaggcg gctgctgctg ttactgcggg tagtggggag cagatactga 7560
aagctattgt tgaggctgct ggtgataagg ctaatcaggg ggggtgtaaag gctgctgggtg 7620
cggctacgaa tccgattgca gctgcgattg ggactgatga tgataatgcg gcggcggttg 7680
ataaggatga gatgaagaag agtaatgata agattgctgc ggctattgtt ttgagggggg 7740
tggctaagga tggaaagtgt gctgctaattg ctaatgataa tagtaaggcg agtgtgaaga 7800
gtgctgtgga tgaggttagc aagtggtagc aagagatgat aacagctgct agtcatgctg 7860
ctacaaaggg tggtagctgt gaagctagcg aaaagattgg ggattctgat gctaataagg 7920
gtgctgggtg tggggcgggcg tttggtgaga atgatatgaa gaagagaaat gataaatattg 7980
ctgcagctat tgttttgagg ggggtggcta aggatggaaa gtttgcctgtt aaggaggatt 8040
attgaactca gctttatagg ggaacagcaa ttcgctagaa aatgattaaa aagcttaact 8100
tcgactgggt cttgccttaa ttttattcct ttgttattat ttatcaatta aattcacttc 8160
ggtttgcttt taaattaatt ctggtatact atgtatacta gatacacaaa ttaaggagaa 8220
gtgaaatgga aaaaatagaa aaatttaaaa acaaattgtc acataaaact caacataaac 8280
taatcgtatt agtatcaaca ctttgcctata taaacaataa aaataaaaaa tattcacaaa 8340
gcaacatcct ttattatttt aatgaaaatt taaaaagaaa tgggcaaacc cctattaaaa 8400
taaaaacatt acaaaaactat ctttataaac tggaaaaaga atttgaagta acaactaatt 8460
attataaaca cttggggggtt aattgtggaa ccgaaattta ctataaactt aaatatcaaa 8520
aacaaaaatg ctatcataaa ataaaccaat attttaaaaa gaaaaaagaa attaaattta 8580
acttaagagt aagtgcattt ttaataaaaa aacactcaaa aaaagggagt gtagaattaa 8640
aggaatgtaa taataataat aataataaag agaaagaaac atcccaaaaa attgaaaattt 8700
tacaacaaca agtctatgcc aaaaaatgta aatttttgac aaactactat actaaaattt 8760
ta

```

&lt;210&gt; 58

&lt;211&gt; 606

&lt;212&gt; DNA

&lt;213&gt; Borrelia afzelii

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(606)

&lt;400&gt; 58

```

gag agt gct gtt gat ggg gtt agc aag tgg tta gaa gag atg ata aaa 48
Glu Ser Ala Val Asp Gly Val Ser Lys Trp Leu Glu Glu Met Ile Lys
1 5 10 15

```

```

gct gct aag gag gct gct aca aag ggt ggt act ggt ggt ggt agc gaa 96
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Gly Ser Glu
20 25 30

```

```

aag att ggg gat gtt ggt gct gct aat aat cag ggt gct gta gct gat 144
Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gln Gly Ala Val Ala Asp
35 40 45

```

```

aag gac agt gtt aag ggg att gcg aag ggg ata aag ggg att gtt gat 192
Lys Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp
50 55 60

```

```

gct gct ggg aag gct ttt ggt aag gat ggt aat gcg ctg aca ggt gta 240

```

Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Thr Gly Val  
65 70 75 80

aaa gaa gtt gct gat gag gct ggg gct aac gag gat gcg ggg aag ttg 288  
Lys Glu Val Ala Asp Glu Ala Gly Ala Asn Glu Asp Ala Gly Lys Leu  
85 90 95

ttt gct ggt aat gct ggt aat gct gct gct gct gac att gcg aag gcg 336  
Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Asp Ile Ala Lys Ala  
100 105 110

gct ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att 384  
Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile  
115 120 125

gtt gat ggt gct ggt ggt gcg gct caa gat ggt aaa aag gct gcg gag 432  
Val Asp Gly Ala Gly Gly Ala Ala Gln Asp Gly Lys Lys Ala Ala Glu  
130 135 140

gct aag aat ccg att gca gct gcg att ggg gct gat gct gct ggt gcg 480  
Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala  
145 150 155 160

gat ttt ggt gat gat atg aag aag agt gat aag att gct gcg gct att 528  
Asp Phe Gly Asp Asp Met Lys Lys Ser Asp Lys Ile Ala Ala Ala Ile  
165 170 175

gtt ttg agg ggg gtg gct aag agt gga aag ttt gct gtt gct aat gct 576  
Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Asn Ala  
180 185 190

gct aag aag gag agt gtg aag agt gct gtg 606  
Ala Lys Lys Glu Ser Val Lys Ser Ala Val  
195 200

<210> 59  
<211> 202  
<212> PRT  
<213> Borrelia afzelii

<400> 59  
Glu Ser Ala Val Asp Gly Val Ser Lys Trp Leu Glu Glu Met Ile Lys  
1 5 10 15

Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Gly Ser Glu  
20 25 30

Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gln Gly Ala Val Ala Asp  
35 40 45

Lys Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp  
50 55 60

Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Thr Gly Val  
65 70 75 80

Lys Glu Val Ala Asp Glu Ala Gly Ala Asn Glu Asp Ala Gly Lys Leu  
85 90 95

Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Asp Ile Ala Lys Ala  
100 105 110

Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile

115		120		125
Val Asp Gly Ala Gly Gly	Ala Ala Gln Asp Gly Lys Lys Ala Ala Glu			
130	135	140		
Ala Lys Asn Pro Ile Ala	Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala			
145	150	155		160
Asp Phe Gly Asp Asp Met	Lys Lys Ser Asp Lys Ile Ala Ala Ala Ile			
165	170	175		
Val Leu Arg Gly Val Ala	Lys Ser Gly Lys Phe Ala Val Ala Asn Ala			
180	185	190		
Ala Lys Lys Glu Ser Val	Lys Ser Ala Val			
195	200			

<210> 60  
 <211> 621  
 <212> DNA  
 <213> Borrelia afzelii

<220>  
 <221> CDS  
 <222> (1) .. (621)

<400> 60	
gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aaa	48
Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys	
1 5 10 15	
gct gct ggt ggg gct gct aag ggt ggt act ggt ggt aat aac gaa aag	96
Ala Ala Gly Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Asn Glu Lys	
20 25 30	
att ggg gat tct gat aat aat aag ggt gct gta gct gat aag gac agt	144
Ile Gly Asp Ser Asp Asn Asn Lys Gly Ala Val Ala Asp Lys Asp Ser	
35 40 45	
gtt aag ggg att gcg aag ggg ata aag ggg att gtt gat gct gct ggg	192
Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly	
50 55 60	
aag gct ttt ggt aag gat ggt aat gcg ctg aag gat gtt gca aaa gtt	240
Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val Ala Lys Val	
65 70 75 80	
gct gat gat gcg gct ggg gct aac gcg aat gca ggg aag ttg ttt gct	288
Ala Asp Asp Ala Ala Gly Ala Asn Ala Asn Ala Gly Lys Leu Phe Ala	
85 90 95	
ggt aat gct gct ggt ggt gcc gct gat gct gat gat gct aac att gcg	336
Gly Asn Ala Ala Gly Gly Ala Ala Asp Ala Asp Asp Ala Asn Ile Ala	
100 105 110	
aag gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa	384
Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys	
115 120 125	
gct att gtt gat gct gct ggt gct gct gct aat cag gat ggt aag aag	432
Ala Ile Val Asp Ala Ala Gly Ala Ala Ala Asn Gln Asp Gly Lys Lys	
130 135 140	

```

gct gcg gat gct aag aat ccg att gca gct gcg att ggg act aat gat 480
Ala Ala Asp Ala Lys Asn Pro Ile Ala Ala Ile Gly Thr Asn Asp
145          150          155          160

gat ggg gcg gag ttt aag gat gga atg aag aag agt gat aat att gct 528
Asp Gly Ala Glu Phe Lys Asp Gly Met Lys Lys Ser Asp Asn Ile Ala
          165          170          175

gca gct att gtt ttg agg ggg gtg gct aag ggt gga aag ttt gct gtt 576
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val
          180          185          190

gct aat gct gct aat gat agt aag gcg agt gtg aag agt gct gtg 621
Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val
          195          200          205

```

&lt;210&gt; 61

&lt;211&gt; 207

&lt;212&gt; PRT

&lt;213&gt; Borrelia afzelii

&lt;400&gt; 61

```

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys
 1          5          10          15

Ala Ala Gly Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Asn Glu Lys
 20          25          30

Ile Gly Asp Ser Asp Asn Asn Lys Gly Ala Val Ala Asp Lys Asp Ser
 35          40          45

Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
 50          55          60

Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val Ala Lys Val
 65          70          75          80

Ala Asp Asp Ala Ala Gly Ala Asn Ala Asn Ala Gly Lys Leu Phe Ala
 85          90          95

Gly Asn Ala Ala Gly Gly Ala Ala Asp Ala Asp Asp Ala Asn Ile Ala
100          105          110

Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys
115          120          125

Ala Ile Val Asp Ala Ala Gly Ala Ala Ala Asn Gln Asp Gly Lys Lys
130          135          140

Ala Ala Asp Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Asp
145          150          155          160

Asp Gly Ala Glu Phe Lys Asp Gly Met Lys Lys Ser Asp Asn Ile Ala
165          170          175

Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val
180          185          190

Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val
195          200          205

```

&lt;210&gt; 62

<211> 618  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(618)

<400> 62  
 gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
           1                  5                  10                  15

gct gct ggt gag gct gct aca aag ggt ggt gat gct ggt ggt ggt gct 96  
 Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala  
                   20                  25                  30

gat aag att ggg gat gtt ggt gct gct aat aat ggt gct gta gct gat 144  
 Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gly Ala Val Ala Asp  
           35                  40                  45

gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att gtt gat 192  
 Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Asp  
           50                  55                  60

gct gct ggg aag gct ttt ggc aag gat ggt aat gcg ctg aag gat gtt 240  
 Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val  
           65                  70                  75                  80

gca gaa gtt gct gat gat aag aag gag gcg ggg aag ttg ttt gct ggt 288  
 Ala Glu Val Ala Asp Asp Lys Lys Glu Ala Gly Lys Leu Phe Ala Gly  
                   85                  90                  95

aat gct ggt ggt gct gtt gct gat gct gct gcg att ggg aag gcg gct 336  
 Asn Ala Gly Gly Ala Val Ala Asp Ala Ala Ala Ile Gly Lys Ala Ala  
           100                  105                  110

ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt 384  
 Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
           115                  120                  125

gat gct gct ggt ggt gcg gat cag gcg ggt aag aag gct gat gcg gct 432  
 Asp Ala Ala Gly Gly Ala Asp Gln Ala Gly Lys Lys Ala Asp Ala Ala  
           130                  135                  140

aag aat ccg att gca gct gcg att ggg gct gat gct gct ggt gct ggt 480  
 Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala Gly  
           145                  150                  155                  160

gcg gat ttt ggt aat gat atg aag aag aga aat gat aag att gtt gcg 528  
 Ala Asp Phe Gly Asn Asp Met Lys Lys Arg Asn Asp Lys Ile Val Ala  
                   165                  170                  175

gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct gct 576  
 Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala  
                   180                  185                  190

gct aat gat gat aat agt aag gcg agt gtg aag agt gct gtg 618  
 Ala Asn Asp Asp Asn Ser Lys Ala Ser Val Lys Ser Ala Val  
           195                  200                  205

<210> 63  
 <211> 206

&lt;212&gt; PRT

<213> *Borrelia afzelii*

&lt;400&gt; 63

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15

Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala  
 20 25 30

Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Asn Gly Ala Val Ala Asp  
 35 40 45

Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Asp  
 50 55 60

Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp Val  
 65 70 75 80

Ala Glu Val Ala Asp Asp Lys Lys Glu Ala Gly Lys Leu Phe Ala Gly  
 85 90 95

Asn Ala Gly Gly Ala Val Ala Asp Ala Ala Ala Ile Gly Lys Ala Ala  
 100 105 110

Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
 115 120 125

Asp Ala Ala Gly Gly Ala Asp Gln Ala Gly Lys Lys Ala Asp Ala Ala  
 130 135 140

Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asp Ala Ala Gly Ala Gly  
 145 150 155 160

Ala Asp Phe Gly Asn Asp Met Lys Lys Arg Asn Asp Lys Ile Val Ala  
 165 170 175

Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala  
 180 185 190

Ala Asn Asp Asp Asn Ser Lys Ala Ser Val Lys Ser Ala Val  
 195 200 205

&lt;210&gt; 64

&lt;211&gt; 630

&lt;212&gt; DNA

<213> *Borrelia afzelii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(630)

&lt;400&gt; 64

gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15

gct gct gat ggg gct gct aaa ggt ggt act ggt ggt aat agc gaa aag 96  
 Ala Ala Asp Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Ser Glu Lys  
 20 25 30

att ggg gat gct ggt gat aat aat aat ggt gct gta gct gat gag aac 144  
 Ile Gly Asp Ala Gly Asp Asn Asn Asn Gly Ala Val Ala Asp Glu Asn

35	40	45	
agt gtt aag gag att gca aag ggg ata aag ggg att gtt gcg gct gct			192
Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala			
50	55	60	
ggg aag gct ttt ggc aag gat ggc aag gat ggt gat gcg ctg aag gat			240
Gly Lys Ala Phe Gly Lys Asp Gly Lys Asp Gly Asp Ala Leu Lys Asp			
65	70	75	80
gtt gaa aca gtt gct gct gag aat gag gct aac aag gat gcg ggg aag			288
Val Glu Thr Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys			
85	90	95	
ttg ttt gct ggt gct aat ggt aat gct ggt gct gct gtt ggt gac att			336
Leu Phe Ala Gly Ala Asn Gly Asn Ala Gly Ala Ala Val Gly Asp Ile			
100	105	110	
gcg aag gcg gct gct gct gtt act gcg gtt agt ggg gag cag ata cta			384
Ala Lys Ala Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu			
115	120	125	
aaa gct att gtt gat gct gct ggt gat gcg gat cag gcg ggt aag aag			432
Lys Ala Ile Val Asp Ala Ala Gly Asp Ala Asp Gln Ala Gly Lys Lys			
130	135	140	
gct gct gag gct aag aat ccg att gca gct gcg att ggg gct aat gct			480
Ala Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asn Ala			
145	150	155	160
gct gat aat gcg gcg gcg ttt ggt aag gat gag atg aag aag agt gat			528
Ala Asp Asn Ala Ala Ala Phe Gly Lys Asp Glu Met Lys Lys Ser Asp			
165	170	175	
aag att gct gca gct att gtt ttg agg ggg gtg gct aag gat gga aag			576
Lys Ile Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys			
180	185	190	
ttt gct gtt gct aat gct aat gat gat aag aag gcg agt gtg aag agt			624
Phe Ala Val Ala Asn Ala Asn Asp Asp Lys Lys Ala Ser Val Lys Ser			
195	200	205	
gct gtg			630
Ala Val			
210			

<210> 65  
 <211> 210  
 <212> PRT  
 <213> Borrelia afzelii

<400> 65  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15  
 Ala Ala Asp Gly Ala Ala Lys Gly Gly Thr Gly Gly Asn Ser Glu Lys  
 20 25 30  
 Ile Gly Asp Ala Gly Asp Asn Asn Asn Gly Ala Val Ala Asp Glu Asn  
 35 40 45  
 Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val Ala Ala Ala  
 50 55 60

Gly Lys Ala Phe Gly Lys Asp Gly Lys Asp Gly Asp Ala Leu Lys Asp  
 65 70 75 80  
 Val Glu Thr Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys  
 85 90 95  
 Leu Phe Ala Gly Ala Asn Gly Asn Ala Gly Ala Ala Val Gly Asp Ile  
 100 105 110  
 Ala Lys Ala Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu  
 115 120 125  
 Lys Ala Ile Val Asp Ala Ala Gly Asp Ala Asp Gln Ala Gly Lys Lys  
 130 135 140  
 Ala Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Asn Ala  
 145 150 155 160  
 Ala Asp Asn Ala Ala Ala Phe Gly Lys Asp Glu Met Lys Lys Ser Asp  
 165 170 175  
 Lys Ile Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys  
 180 185 190  
 Phe Ala Val Ala Asn Ala Asn Asp Asp Lys Lys Ala Ser Val Lys Ser  
 195 200 205  
 Ala Val  
 210

<210> 66  
 <211> 612  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(612)

<400> 66  
 gag agt gct gtg gat gag gtt agc aag tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15  
 gct gct aag gag gct gct aca aag ggt ggt act ggt ggt aat aac gaa 96  
 Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Asn Asn Glu  
 20 25 30  
 aag att gga gat tct gat gct aat aat ggt gcg aag gct gat gcg agc 144  
 Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Lys Ala Asp Ala Ser  
 35 40 45  
 agt gtt aat ggg att gcg aat ggg ata aag ggg att gtt gat gct gct 192  
 Ser Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala  
 50 55 60  
 ggg aag gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt aaa aca 240  
 Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr  
 65 70 75 80  
 gtt gct gct gag aat gag gct aac aag gat gcg ggg aag ttg ttt gct 288  
 Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala



85										90					95					
ggt aag aat ggt aat gct gat gct gct gat gct gct gac att gcg aag	336																			
Gly Lys Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys																				
100 105 110																				
gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa gct	384																			
Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala																				
115 120 125																				
att gtt gat ggt gct ggt gat gca gct aat cag gcg ggt aaa aag gct	432																			
Ile Val Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala																				
130 135 140																				
gct gag gct aag aat ccg att gcg gct gcg att ggg act aat gaa gct	480																			
Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala																				
145 150 155 160																				
ggg gcg gag ttt ggt gat gat atg aag aag aga aat gat aag att gct	528																			
Gly Ala Glu Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala																				
165 170 175																				
gcg gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gtt	576																			
Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val																				
180 185 190																				
gct aat gct gct gct gat aat agt aag gcg agt gtg	612																			
Ala Asn Ala Ala Ala Asp Asn Ser Lys Ala Ser Val																				
195 200																				

&lt;210&gt; 67

&lt;211&gt; 204

&lt;212&gt; PRT

&lt;213&gt; Borrelia afzelii

&lt;400&gt; 67

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
Ala Ala Lys Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Asn Asn Glu	
20 25 30	
Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Lys Ala Asp Ala Ser	
35 40 45	
Ser Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala	
50 55 60	
Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr	
65 70 75 80	
Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala	
85 90 95	
Gly Lys Asn Gly Asn Ala Asp Ala Ala Asp Ala Ala Asp Ile Ala Lys	
100 105 110	
Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys Ala	
115 120 125	
Ile Val Asp Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala	
130 135 140	

Ala Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala  
145 150 155 160

Gly Ala Glu Phe Gly Asp Asp Met Lys Lys Arg Asn Asp Lys Ile Ala  
165 170 175

Ala Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val  
180 185 190

Ala Asn Ala Ala Ala Asp Asn Ser Lys Ala Ser Val  
195 200

<210> 68

<211> 609

<212> DNA

<213> Borrelia afzelii

<220>

<221> CDS

<222> (1)..(609)

<400> 68

aag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aag 48  
Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys  
1 5 10 15

gct gct ggt gag gct gct aca aag ggt ggt gat gct ggt ggt ggt gct 96  
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala  
20 25 30

gat aag att ggg gat gct ggt gat aag ggt gct gta gct gat gcg agc 144  
Asp Lys Ile Gly Asp Ala Gly Asp Lys Gly Ala Val Ala Asp Ala Ser  
35 40 45

agt gtt aag gag att gcg aat ggg ata aag ggg att gtt gat gct gct 192  
Ser Val Lys Glu Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala  
50 55 60

ggg aag gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt aaa aca 240  
Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr  
65 70 75 80

ggt gct gct gag aat gag gct aac aag gat gcg ggg aag ttg ttt gct 288  
Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala  
85 90 95

ggt aat gct ggt aat ggt gct gct gat gac att gcg aag gcg gct gct 336  
Gly Asn Ala Gly Asn Gly Ala Ala Asp Asp Ile Ala Lys Ala Ala Ala  
100 105 110

gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gat 384  
Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp  
115 120 125

gct gct ggt gat aag gct aat cag gat ggt aaa aag gct gcg gat gct 432  
Ala Ala Gly Asp Lys Ala Asn Gln Asp Gly Lys Lys Ala Ala Asp Ala  
130 135 140

aag aat ccg att gcg gct gcg att ggg gct gct gat gct ggt gct gcg 480  
Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Asp Ala Gly Ala Ala  
145 150 155 160

gcg gcg ttt aat gag aat gat atg aag aag agt gat aag att gct gca 528

Ala Ala Phe Asn Glu Asn Asp Met Lys Lys Ser Asp Lys Ile Ala Ala  
 165 170 175

gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct gct 576  
 Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala  
 180 185 190

gat gct gat gct aat aat agt aag gcg agc gtg 609  
 Asp Ala Asp Ala Asn Asn Ser Lys Ala Ser Val  
 195 200

<210> 69  
 <211> 203  
 <212> PRT  
 <213> *Borrelia afzelii*

<400> 69  
 Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys  
 1 5 10 15

Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Asp Ala Gly Gly Gly Ala  
 20 25 30

Asp Lys Ile Gly Asp Ala Gly Asp Lys Gly Ala Val Ala Asp Ala Ser  
 35 40 45

Ser Val Lys Glu Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala  
 50 55 60

Gly Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Lys Thr  
 65 70 75 80

Val Ala Ala Glu Asn Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala  
 85 90 95

Gly Asn Ala Gly Asn Gly Ala Ala Asp Asp Ile Ala Lys Ala Ala Ala  
 100 105 110

Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp  
 115 120 125

Ala Ala Gly Asp Lys Ala Asn Gln Asp Gly Lys Lys Ala Ala Asp Ala  
 130 135 140

Lys Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Asp Ala Gly Ala Ala  
 145 150 155 160

Ala Ala Phe Asn Glu Asn Asp Met Lys Lys Ser Asp Lys Ile Ala Ala  
 165 170 175

Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala  
 180 185 190

Asp Ala Asp Ala Asn Asn Ser Lys Ala Ser Val  
 195 200

<210> 70  
 <211> 600  
 <212> DNA  
 <213> *Borrelia afzelii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(600)

&lt;400&gt; 70

aag	agt	gct	gtt	ggg	gag	gtt	agc	aag	tgg	tta	gaa	gag	atg	ata	aaa	48
Lys	Ser	Ala	Val	Gly	Glu	Val	Ser	Lys	Trp	Leu	Glu	Glu	Met	Ile	Lys	
1				5					10					15		

gct	gct	ggg	gag	gct	gca	aaa	gtt	ggg	ggg	act	ggg	ggg	agc	gaa	aag	96
Ala	Ala	Gly	Glu	Ala	Ala	Lys	Val	Gly	Gly	Thr	Gly	Gly	Ser	Glu	Lys	
		20						25					30			

att	ggg	gat	gct	gat	aat	aat	aag	ggg	gct	gta	gct	gat	gcg	agc	agt	144
Ile	Gly	Asp	Ala	Asp	Asn	Asn	Lys	Gly	Ala	Val	Ala	Asp	Ala	Ser	Ser	
		35					40						45			

gtt	aat	ggg	att	gcg	aat	ggg	ata	aag	ggg	att	gtt	gat	gct	gct	ggg	192
Val	Asn	Gly	Ile	Ala	Asn	Gly	Ile	Lys	Gly	Ile	Val	Asp	Ala	Ala	Gly	
	50					55					60					

aag	gct	ttt	ggg	aag	gat	ggg	gcg	ctg	gca	ggg	gtt	gca	gct	gct	gct	240
Lys	Ala	Phe	Gly	Lys	Asp	Gly	Ala	Leu	Ala	Gly	Val	Ala	Ala	Ala	Ala	
65					70					75					80	

gag	aat	gat	gat	aag	aag	gat	gcg	ggg	aag	ttg	ttt	gct	ggg	aag	aat	288
Glu	Asn	Asp	Asp	Lys	Lys	Asp	Ala	Gly	Lys	Leu	Phe	Ala	Gly	Lys	Asn	
				85					90					95		

ggg	ggg	gct	ggg	gct	gct	gat	gcg	att	ggg	aag	gcg	gct	gct	gct	gtt	336
Gly	Gly	Ala	Gly	Ala	Ala	Asp	Ala	Ile	Gly	Lys	Ala	Ala	Ala	Ala	Val	
			100					105						110		

act	gcg	gtt	agt	ggg	gag	cag	ata	ctg	aaa	gct	att	gtt	gat	gct	gct	384
Thr	Ala	Val	Ser	Gly	Glu	Gln	Ile	Leu	Lys	Ala	Ile	Val	Asp	Ala	Ala	
		115					120					125				

ggg	gct	gca	gct	aat	cag	gcg	ggg	aaa	aag	gct	gcg	gat	gct	aag	aat	432
Gly	Ala	Ala	Ala	Asn	Gln	Ala	Gly	Lys	Lys	Ala	Ala	Asp	Ala	Lys	Asn	
		130				135						140				

ccg	att	gcg	gct	gcg	att	ggg	act	gct	gat	gat	ggg	gcg	gag	ttt	aag	480
Pro	Ile	Ala	Ala	Ala	Ile	Gly	Thr	Ala	Asp	Asp	Gly	Ala	Glu	Phe	Lys	
145					150					155					160	

gat	gat	atg	aag	aag	agt	gat	aat	att	gct	gcg	gct	att	gtt	ttg	agg	528
Asp	Asp	Met	Lys	Lys	Ser	Asp	Asn	Ile	Ala	Ala	Ala	Ile	Val	Leu	Arg	
				165					170					175		

ggg	gtg	gct	aag	gat	gga	aag	ttt	gct	gtt	gct	aat	gct	gat	gat	aat	576
Gly	Val	Ala	Lys	Asp	Gly	Lys	Phe	Ala	Val	Ala	Asn	Ala	Asp	Asp	Asn	
			180					185					190			

aag	gcg	agt	gtg	aag	agt	gct	gtg									600
Lys	Ala	Ser	Val	Lys	Ser	Ala	Val									
			195				200									

&lt;210&gt; 71

&lt;211&gt; 200

&lt;212&gt; PRT

<213> *Borrelia afzelii*

&lt;400&gt; 71

Lys Ser Ala Val Gly Glu Val Ser Lys Trp Leu Glu Glu Met Ile Lys  
 1 5 10 15  
 Ala Ala Gly Glu Ala Ala Lys Val Gly Gly Thr Gly Gly Ser Glu Lys  
 20 25 30  
 Ile Gly Asp Ala Asp Asn Asn Lys Gly Ala Val Ala Asp Ala Ser Ser  
 35 40 45  
 Val Asn Gly Ile Ala Asn Gly Ile Lys Gly Ile Val Asp Ala Ala Gly  
 50 55 60  
 Lys Ala Phe Gly Lys Asp Gly Ala Leu Ala Gly Val Ala Ala Ala Ala  
 65 70 75 80  
 Glu Asn Asp Asp Lys Lys Asp Ala Gly Lys Leu Phe Ala Gly Lys Asn  
 85 90 95  
 Gly Gly Ala Gly Ala Ala Asp Ala Ile Gly Lys Ala Ala Ala Ala Val  
 100 105 110  
 Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala  
 115 120 125  
 Gly Ala Ala Ala Asn Gln Ala Gly Lys Lys Ala Ala Asp Ala Lys Asn  
 130 135 140  
 Pro Ile Ala Ala Ala Ile Gly Thr Ala Asp Asp Gly Ala Glu Phe Lys  
 145 150 155 160  
 Asp Asp Met Lys Lys Ser Asp Asn Ile Ala Ala Ala Ile Val Leu Arg  
 165 170 175  
 Gly Val Ala Lys Asp Gly Lys Phe Ala Val Ala Asn Ala Asp Asp Asn  
 180 185 190  
 Lys Ala Ser Val Lys Ser Ala Val  
 195 200

<210> 72  
 <211> 592  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1) .. (591)

<400> 72  
 gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15  
 gct gct ggt gag gct gca aaa gtt ggt gct ggt ggt ggt gct gat aag 96  
 Ala Ala Gly Glu Ala Ala Lys Val Gly Ala Gly Gly Gly Ala Asp Lys  
 20 25 30  
 att ggg gat gct gct aat aat cag ggt gcg aag gct gat gag agc agt 144  
 Ile Gly Asp Ala Ala Asn Asn Gln Gly Ala Lys Ala Asp Glu Ser Ser  
 35 40 45  
 gtt aat gga att gca aag ggg ata aag ggg att gtt gat gct gct ggg 192  
 Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly

50	55	60	
aag gct ttt ggc aag gag ggt agt gcg ctg aag gat gtt gca aaa gtt			240
Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val			
65	70	75	80
gct gat gat gat aac aag gat gcg ggg aag ttg ttt gct ggt aat gct			288
Ala Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Asn Ala			
	85	90	95
ggt ggt ggt gct ggt gct gat att gcg aag gcg gct gct gct gtt act			336
Gly Gly Gly Ala Gly Ala Asp Ile Ala Lys Ala Ala Ala Val Thr			
	100	105	110
gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gat gct gct ggt			384
Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly			
	115	120	125
gct gcg gat cag gcg ggt gca gct gct ggt gcg gct aag aat ccg att			432
Ala Ala Asp Gln Ala Gly Ala Ala Ala Gly Ala Ala Lys Asn Pro Ile			
	130	135	140
gcg gct gcg att ggg gct gat gct ggt gct gcg gag gag ttt aag gat			480
Ala Ala Ala Ile Gly Ala Asp Ala Gly Ala Ala Glu Glu Phe Lys Asp			
	145	150	155
gag atg aag aag agt gat aag att gct gcg gct att gtt ttg agg ggg			528
Glu Met Lys Lys Ser Asp Lys Ile Ala Ala Ala Ile Val Leu Arg Gly			
	165	170	175
gtg gct aag ggt gga aag ttt gct gtt gct gct aat gat gct gca aat			576
Val Ala Lys Gly Gly Lys Phe Ala Val Ala Ala Asn Asp Ala Ala Asn			
	180	185	190
gtg aag agt gct gtg g			592
Val Lys Ser Ala Val			
	195		

&lt;210&gt; 73

&lt;211&gt; 197

&lt;212&gt; PRT

&lt;213&gt; Borrelia afzelii

&lt;400&gt; 73

Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr
1 5 10 15

Ala Ala Gly Glu Ala Ala Lys Val Gly Ala Gly Gly Gly Ala Asp Lys
20 25 30

Ile Gly Asp Ala Ala Asn Asn Gln Gly Ala Lys Ala Asp Glu Ser Ser
35 40 45

Val Asn Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly
50 55 60

Lys Ala Phe Gly Lys Glu Gly Ser Ala Leu Lys Asp Val Ala Lys Val
65 70 75 80

Ala Asp Asp Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Asn Ala
85 90 95

Gly Gly Gly Ala Gly Ala Asp Ile Ala Lys Ala Ala Ala Val Thr
---

100 105 110  
 Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Gly  
 115 120 125  
 Ala Ala Asp Gln Ala Gly Ala Ala Ala Gly Ala Ala Lys Asn Pro Ile  
 130 135 140  
 Ala Ala Ala Ile Gly Ala Asp Ala Gly Ala Ala Glu Glu Phe Lys Asp  
 145 150 155 160  
 Glu Met Lys Lys Ser Asp Lys Ile Ala Ala Ile Val Leu Arg Gly  
 165 170 175  
 Val Ala Lys Gly Gly Lys Phe Ala Val Ala Ala Asn Asp Ala Ala Asn  
 180 185 190  
 Val Lys Ser Ala Val  
 195

<210> 74  
 <211> 597  
 <212> DNA  
 <213> Borrelia afzelii

<220>  
 <221> CDS  
 <222> (1)..(597)

<400> 74  
 gag agt gct gtt ggt gag gtt agc gca tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Gly Glu Val Ser Ala Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15  
 gct gct agt gag gct gct aca aag ggt ggt act ggt ggt act ggt ggt 96  
 Ala Ala Ser Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Thr Gly Gly  
 20 25 30  
 gat agt gaa aag att ggg gat tct gat gct aat aat ggt gct gta gct 144  
 Asp Ser Glu Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Val Ala  
 35 40 45  
 gat gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att gtt 192  
 Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val  
 50 55 60  
 gat gct gct ggg aag gct ttt ggt aag gat ggt aat gcg ctg aag gat 240  
 Asp Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp  
 65 70 75 80  
 gtt gca gaa gtt gct gat gat gag gct aac gcg gat gcg ggg aag ttg 288  
 Val Ala Glu Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu  
 85 90 95  
 ttt gct ggt aat gct ggt aat gct gct gct gct gac gtt gcg aag gcg 336  
 Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Ala Asp Val Ala Lys Ala  
 100 105 110  
 gct ggt gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att 384  
 Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile  
 115 120 125  
 gtt gat gct gct ggt gct gcg gat cag gcg ggt gca aag gct gat gcg 432

Val Asp Ala Ala Gly Ala Ala Asp Gln Ala Gly Ala Lys Ala Asp Ala  
 130 135 140

gct aag aat ccg att gca gct gcg att ggg act aat gaa gct ggg gcg 480  
 Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala  
 145 150 155 160

gcg ttt aag gat gga atg aag aag aga aat gat aat att gct gcg gct 528  
 Ala Phe Lys Asp Gly Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala  
 165 170 175

att gtt ttg agg ggg gtg gct aag agt gga aag ttt gct gtt gct gct 576  
 Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Ala  
 180 185 190

gct gat gct ggt aag gcg aga 597  
 Ala Asp Ala Gly Lys Ala Arg  
 195

<210> 75  
 <211> 199  
 <212> PRT  
 <213> Borrelia afzelii

<400> 75  
 Glu Ser Ala Val Gly Glu Val Ser Ala Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15

Ala Ala Ser Glu Ala Ala Thr Lys Gly Gly Thr Gly Gly Thr Gly Gly  
 20 25 30

Asp Ser Glu Lys Ile Gly Asp Ser Asp Ala Asn Asn Gly Ala Val Ala  
 35 40 45

Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile Val  
 50 55 60

Asp Ala Ala Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Lys Asp  
 65 70 75 80

Val Ala Glu Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu  
 85 90 95

Phe Ala Gly Asn Ala Gly Asn Ala Ala Ala Ala Asp Val Ala Lys Ala  
 100 105 110

Ala Gly Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile  
 115 120 125

Val Asp Ala Ala Gly Ala Ala Asp Gln Ala Gly Ala Lys Ala Asp Ala  
 130 135 140

Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asn Glu Ala Gly Ala  
 145 150 155 160

Ala Phe Lys Asp Gly Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala  
 165 170 175

Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala Val Ala Ala  
 180 185 190

Ala Asp Ala Gly Lys Ala Arg  
 195



<210> 76  
 <211> 621  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(621)

<400> 76  
 gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
 1 5 10 15  
 gct gct agt gag gct gca aaa gtt ggt gct ggt ggt gat gat aag att 96  
 Ala Ala Ser Glu Ala Ala Lys Val Gly Ala Gly Gly Asp Asp Lys Ile  
 20 25 30  
 ggg gat tct gct aat aat ggt gct gta gct gat gcg ggc agt gtt aag 144  
 Gly Asp Ser Ala Asn Asn Gly Ala Val Ala Asp Ala Gly Ser Val Lys  
 35 40 45  
 gga att gcg aag ggg ata aag ggg att gtt gat gct gct ggg aag gct 192  
 Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala  
 50 55 60  
 ttt ggt aag gag ggt gat gcg ctg aag gat gtt gca aaa gtt gct gat 240  
 Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala Asp  
 65 70 75 80  
 gag aat ggg gat aac aag gat gcg ggg aag ttg ttt gct ggt gag aat 288  
 Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu Asn  
 85 90 95  
 ggt aat gct ggt ggt gct gct gat gct gac att gcg aag gcg gct gct 336  
 Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala Ala  
 100 105 110  
 gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gag 384  
 Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu  
 115 120 125  
 gct gct ggt gct ggt gat gca gct aat cag gcg ggt aag aag gct gat 432  
 Ala Ala Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Asp  
 130 135 140  
 gag gct aag aat ccg att gcg gct gcg att ggg act gat gat gct ggg 480  
 Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly  
 145 150 155 160  
 gcg gcg ttt ggt cag gat gat atg aag aag aga aat gat aat att gct 528  
 Ala Ala Phe Gly Gln Asp Asp Met Lys Lys Arg Asn Asp Asn Ile Ala  
 165 170 175  
 gcg gct att gtt ttg agg ggg gtg gct aag ggt gga aag ttt gct gtt 576  
 Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val  
 180 185 190  
 gct aat gct gct aat gat agt aag gcg agt gtg aag agt gct gtg 621  
 Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val  
 195 200 205

<210> 77  
 <211> 207  
 <212> PRT  
 <213> *Borrelia afzelii*

<400> 77  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr  
   1                  5                  10                  15  
 Ala Ala Ser Glu Ala Ala Lys Val Gly Ala Gly Gly Asp Asp Lys Ile  
                   20                  25                  30  
 Gly Asp Ser Ala Asn Asn Gly Ala Val Ala Asp Ala Gly Ser Val Lys  
                   35                  40                  45  
 Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Gly Lys Ala  
                   50                  55                  60  
 Phe Gly Lys Glu Gly Asp Ala Leu Lys Asp Val Ala Lys Val Ala Asp  
   65                  70                  75                  80  
 Glu Asn Gly Asp Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly Glu Asn  
                   85                  90                  95  
 Gly Asn Ala Gly Gly Ala Ala Asp Ala Asp Ile Ala Lys Ala Ala Ala  
                   100                  105                  110  
 Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu  
                   115                  120                  125  
 Ala Ala Gly Ala Gly Asp Ala Ala Asn Gln Ala Gly Lys Lys Ala Asp  
                   130                  135                  140  
 Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala Gly  
   145                  150                  155                  160  
 Ala Ala Phe Gly Gln Asp Asp Met Lys Lys Arg Asn Asp Asn Ile Ala  
                   165                  170                  175  
 Ala Ala Ile Val Leu Arg Gly Val Ala Lys Gly Gly Lys Phe Ala Val  
                   180                  185                  190  
 Ala Asn Ala Ala Asn Asp Ser Lys Ala Ser Val Lys Ser Ala Val  
                   195                  200                  205

<210> 78  
 <211> 459  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(459)

<400> 78  
 gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag ata ata aca 48  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Ile Ile Thr  
   1                  5                  10                  15  
 gct act ggg aag gct ttt ggt aag gat ggt aat gcg ctg gca ggt gtt 96  
 Ala Thr Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Ala Gly Val  
                   20                  25                  30

gca aaa gtt gct gat gat gag gct aac gcg gat gcg ggg aag ttg ttt 144  
 Ala Lys Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu Phe  
                   35                                  40                                  45

gct ggt gag aat ggt aat gct ggt gct gct gcg att ggg aag gcg gct 192  
 Ala Gly Glu Asn Gly Asn Ala Gly Ala Ala Ala Ile Gly Lys Ala Ala  
                   50                                  55                                  60

gct gct gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt 240  
 Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
                   65                                  70                                  75                                  80

gat gct gct ggt ggt gcg gct cag gtg ggt gct ggt gct ggt gcg gct 288  
 Asp Ala Ala Gly Gly Ala Ala Gln Val Gly Ala Gly Ala Gly Ala Ala  
                                   85                                  90                                  95

acg aat ccg att gca gct gcg att ggg gct gct ggt gat ggt gcg gat 336  
 Thr Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Gly Asp Gly Ala Asp  
                                   100                                  105                                  110

ttt ggt aag gat gag atg aag aag aga aat gat aag att gct gcg gct 384  
 Phe Gly Lys Asp Glu Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala  
                   115                                  120                                  125

att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct gct gct 432  
 Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala Ala  
                   130                                  135                                  140

aat gat agt aag gcg agt gtg aag agt 459  
 Asn Asp Ser Lys Ala Ser Val Lys Ser  
                   145                                  150

<210> 79  
 <211> 153  
 <212> PRT  
 <213> *Borrelia afzelii*

<400> 79  
 Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Ile Ile Thr  
           1                                  5                                  10                                  15

Ala Thr Gly Lys Ala Phe Gly Lys Asp Gly Asn Ala Leu Ala Gly Val  
                   20                                  25                                  30

Ala Lys Val Ala Asp Asp Glu Ala Asn Ala Asp Ala Gly Lys Leu Phe  
                   35                                  40                                  45

Ala Gly Glu Asn Gly Asn Ala Gly Ala Ala Ala Ile Gly Lys Ala Ala  
                   50                                  55                                  60

Ala Ala Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val  
                   65                                  70                                  75                                  80

Asp Ala Ala Gly Gly Ala Ala Gln Val Gly Ala Gly Ala Gly Ala Ala  
                                   85                                  90                                  95

Thr Asn Pro Ile Ala Ala Ala Ile Gly Ala Ala Gly Asp Gly Ala Asp  
                   100                                  105                                  110

Phe Gly Lys Asp Glu Met Lys Lys Arg Asn Asp Lys Ile Ala Ala Ala  
                   115                                  120                                  125

Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Ala Ala  
 130 135 140

Asn Asp Ser Lys Ala Ser Val Lys Ser  
 145 150

<210> 80  
 <211> 612  
 <212> DNA  
 <213> *Borrelia afzelii*

<220>  
 <221> CDS  
 <222> (1)..(612)

<400> 80  
 gct gtg gag agt gct gtt gat gag gtt agc aag tgg tta gaa gag atg 48  
 Ala Val Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met  
 1 5 10 15

ata aca gct gct gat gct gct gct gct aaa gtt ggc gat gct ggt ggt 96  
 Ile Thr Ala Ala Asp Ala Ala Ala Ala Lys Val Gly Asp Ala Gly Gly  
 20 25 30

ggt gct gat aag att ggg gat gtt ggt gct gct aat aag ggt gcg aag 144  
 Gly Ala Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Lys Gly Ala Lys  
 35 40 45

gct gat gcg agc agt gtt aag gag att gcg aag ggg ata aag ggg att 192  
 Ala Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile  
 50 55 60

gtt gat gct gct ggg aag gct ttt ggt ggt gat gcg ctg aag gat gtt 240  
 Val Asp Ala Ala Gly Lys Ala Phe Gly Gly Asp Ala Leu Lys Asp Val  
 65 70 75 80

aaa gct gct ggt gat gat aac aag gag gca ggg aag ttg ttt gct ggt 288  
 Lys Ala Ala Gly Asp Asp Asn Lys Glu Ala Gly Lys Leu Phe Ala Gly  
 85 90 95

gct aat ggt aat gct ggt gct aat gct gct gct gct gat gac att gcg 336  
 Ala Asn Gly Asn Ala Gly Ala Asn Ala Ala Ala Asp Asp Ile Ala  
 100 105 110

aag gcg gct ggt gct gtt agt gcg gtt agt ggg gag cag ata ctg aaa 384  
 Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys  
 115 120 125

gct att gtt gag gcg gct ggt gct gcg gat cag gcg ggt gta aag gct 432  
 Ala Ile Val Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala  
 130 135 140

gag gag gct aag aat ccg att gca gct gcg att ggg act gat gat gct 480  
 Glu Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala  
 145 150 155 160

ggt gcg gcg gag ttt ggt gag aat gat atg aag aag aat gat aat att 528  
 Gly Ala Ala Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile  
 165 170 175

gct gcg gct att gtt ttg agg ggg gtg gct aag agt gga aag ttt gct 576  
 Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala  
 180 185 190

gct aat gct aat gat gct ggt aag aag gag agt gtg  
 Ala Asn Ala Asn Asp Ala Gly Lys Lys Glu Ser Val  
 195 200

612

<210> 81  
 <211> 204  
 <212> PRT  
 <213> Borrelia afzelii

<400> 81  
 Ala Val Glu Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met  
 1 5 10 15  
 Ile Thr Ala Ala Asp Ala Ala Ala Lys Val Gly Asp Ala Gly Gly  
 20 25 30  
 Gly Ala Asp Lys Ile Gly Asp Val Gly Ala Ala Asn Lys Gly Ala Lys  
 35 40 45  
 Ala Asp Ala Ser Ser Val Lys Glu Ile Ala Lys Gly Ile Lys Gly Ile  
 50 55 60  
 Val Asp Ala Ala Gly Lys Ala Phe Gly Gly Asp Ala Leu Lys Asp Val  
 65 70 75 80  
 Lys Ala Ala Gly Asp Asp Asn Lys Glu Ala Gly Lys Leu Phe Ala Gly  
 85 90 95  
 Ala Asn Gly Asn Ala Gly Ala Asn Ala Ala Ala Asp Asp Ile Ala  
 100 105 110  
 Lys Ala Ala Gly Ala Val Ser Ala Val Ser Gly Glu Gln Ile Leu Lys  
 115 120 125  
 Ala Ile Val Glu Ala Ala Gly Ala Ala Asp Gln Ala Gly Val Lys Ala  
 130 135 140  
 Glu Glu Ala Lys Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Ala  
 145 150 155 160  
 Gly Ala Ala Glu Phe Gly Glu Asn Asp Met Lys Lys Asn Asp Asn Ile  
 165 170 175  
 Ala Ala Ala Ile Val Leu Arg Gly Val Ala Lys Ser Gly Lys Phe Ala  
 180 185 190  
 Ala Asn Ala Asn Asp Ala Gly Lys Lys Glu Ser Val  
 195 200

<210> 82  
 <211> 603  
 <212> DNA  
 <213> Borrelia afzelii

<220>  
 <221> CDS  
 <222> (1)..(603)

<400> 82  
 aag agt gct gtg gat gag gct agc aag tgg tta gaa gag atg ata aca  
 Lys Ser Ala Val Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr 48

1	5	10	15	
gct gct ggt gag gct gct aca aag ggt ggt act ggt gaa gct agc gaa				96
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu				
20		25	30	
aag att ggg gat gtt ggt gat aat aat cat ggt gct gta gct gat gcg				144
Lys Ile Gly Asp Val Gly Asp Asn Asn His Gly Ala Val Ala Asp Ala				
35	40	45		
gac agt gtt aag ggg att gcg aag ggg ata aag ggg att gtt gat gct				192
Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala				
50	55	60		
gct ggg aag gct ttt ggt aag gat ggt gcg ctg aag gat gtt gca gct				240
Ala Gly Lys Ala Phe Gly Lys Asp Gly Ala Leu Lys Asp Val Ala Ala				
65	70	75	80	
gct gct ggt gat gag gct aac aag gat gcg ggg aag ttg ttt gct ggt				288
Ala Ala Gly Asp Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly				
85	90	95		
cag gat ggt ggt ggt gct gat ggt gac att gcg aag gcg gct gct gct				336
Gln Asp Gly Gly Gly Ala Asp Gly Asp Ile Ala Lys Ala Ala Ala Ala				
100	105	110		
gtt act gcg gtt agt ggg gag cag ata ctg aaa gct att gtt gag gct				384
Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala				
115	120	125		
gct ggt gat aag gct aat cag gtg ggt gta aag gct gct ggt gcg gct				432
Ala Gly Asp Lys Ala Asn Gln Val Gly Val Lys Ala Ala Gly Ala Ala				
130	135	140		
acg aat ccg att gca gct gcg att ggg act gat gat gat aat gcg gcg				480
Thr Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Asp Asn Ala Ala				
145	150	155	160	
gcg ttt gat aag gat gag atg aag aag agt aat gat aag att gct gcg				528
Ala Phe Asp Lys Asp Glu Met Lys Lys Ser Asn Asp Lys Ile Ala Ala				
165	170	175		
gct att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gct aat				576
Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Asn				
180	185	190		
gct aat gat aat agt aag gcg agt gtg				603
Ala Asn Asp Asn Ser Lys Ala Ser Val				
195	200			
<210> 83				
<211> 201				
<212> PRT				
<213> Borrelia afzelii				
<400> 83				
Lys Ser Ala Val Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr				
1	5	10	15	
Ala Ala Gly Glu Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu				
20	25	30		
Lys Ile Gly Asp Val Gly Asp Asn Asn His Gly Ala Val Ala Asp Ala				

35 40 45

Asp Ser Val Lys Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala  
50 55 60

Ala Gly Lys Ala Phe Gly Lys Asp Gly Ala Leu Lys Asp Val Ala Ala  
65 70 75 80

Ala Ala Gly Asp Glu Ala Asn Lys Asp Ala Gly Lys Leu Phe Ala Gly  
85 90 95

Gln Asp Gly Gly Gly Ala Asp Gly Asp Ile Ala Lys Ala Ala Ala  
100 105 110

Val Thr Ala Val Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Glu Ala  
115 120 125

Ala Gly Asp Lys Ala Asn Gln Val Gly Val Lys Ala Ala Gly Ala Ala  
130 135 140

Thr Asn Pro Ile Ala Ala Ala Ile Gly Thr Asp Asp Asp Asn Ala Ala  
145 150 155 160

Ala Phe Asp Lys Asp Glu Met Lys Lys Ser Asn Asp Lys Ile Ala Ala  
165 170 175

Ala Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Ala Asn  
180 185 190

Ala Asn Asp Asn Ser Lys Ala Ser Val  
195 200

<210> 84  
<211> 249  
<212> DNA  
<213> *Borrelia afzelii*

<220>  
<221> CDS  
<222> (1)..(249)

<400> 84

aag agt gct gtg gat gag gtt agc aag tgg tta gaa gag atg ata aca	48
Lys Ser Ala Val Asp Glu Val Ser Lys Trp Leu Glu Glu Met Ile Thr	
1 5 10 15	
gct gct agt gat gct gct aca aag ggt ggt act ggt gaa gct agc gaa	96
Ala Ala Ser Asp Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu	
20 25 30	
aag att ggg gat tct gat gct aat aag ggt gct ggt gct ggg gcg gcg	144
Lys Ile Gly Asp Ser Asp Ala Asn Lys Gly Ala Gly Ala Gly Ala Ala	
35 40 45	
ttt ggt gag aat gat atg aag aag aga aat gat aat att gct gca gct	192
Phe Gly Glu Asn Asp Met Lys Lys Arg Asn Asp Asn Ile Ala Ala Ala	
50 55 60	
att gtt ttg agg ggg gtg gct aag gat gga aag ttt gct gtt aag gag	240
Ile Val Leu Arg Gly Val Ala Lys Asp Gly Lys Phe Ala Val Lys Glu	
65 70 75 80	
gat tat tga	249

Asp Tyr

```
<210> 85
<211> 82
<212> PRT
<213> Borrelia afzelii
```

<400>	85																
Lys	Ser	Ala	Val	Asp	Glu	Val	Ser	Lys	Trp	Leu	Glu	Glu	Met	Ile	Thr		
1				5					10					15			
Ala	Ala	Ser	Asp	Ala	Ala	Thr	Lys	Gly	Gly	Thr	Gly	Glu	Ala	Ser	Glu		
			20					25					30				
Lys	Ile	Gly	Asp	Ser	Asp	Ala	Asn	Lys	Gly	Ala	Gly	Ala	Gly	Ala	Ala		
		35					40					45					
Phe	Gly	Glu	Asn	Asp	Met	Lys	Lys	Arg	Asn	Asp	Asn	Ile	Ala	Ala	Ala		
50						55					60						
Ile	Val	Leu	Arg	Gly	Val	Ala	Lys	Asp	Gly	Lys	Phe	Ala	Val	Lys	Glu		
65					70					75					80		
Asp	Tyr																

```
<210> 86
<211> 537
<212> DNA
<213> Borrelia afzelii
```

```
<220>
<221> CDS
<222> (1) .. (537)
```

<400>	86																	48
atg	gaa	aaa	ata	gaa	aaa	ttt	aaa	aac	aaa	tgt	caa	cat	aaa	cta	caa			
Met	Glu	Lys	Ile	Glu	Lys	Phe	Lys	Asn	Lys	Cys	Gln	His	Lys	Leu	Gln			
1				5					10					15				
cat	aaa	cta	atc	gta	tta	gta	tca	aca	ctt	tgc	tat	ata	aac	aat	aaa		96	
His	Lys	Leu	Ile	Val	Leu	Val	Ser	Thr	Leu	Cys	Tyr	Ile	Asn	Asn	Lys			
			20					25					30					
aat	aaa	aaa	tat	tca	caa	agc	aac	atc	ctt	tat	tat	ttt	aat	gaa	aat		144	
Asn	Lys	Lys	Tyr	Ser	Gln	Ser	Asn	Ile	Leu	Tyr	Tyr	Phe	Asn	Glu	Asn			
		35					40					45						
tta	aaa	aga	aat	ggg	caa	acc	cct	att	aaa	ata	aaa	aca	tta	caa	aac		192	
Leu	Lys	Arg	Asn	Gly	Gln	Thr	Pro	Ile	Lys	Ile	Lys	Thr	Leu	Gln	Asn			
	50					55					60							
tat	ctt	tat	aaa	ctg	gaa	aaa	gaa	ttt	gaa	gta	aca	act	aat	tat	tat		240	
Tyr	Leu	Tyr	Lys	Leu	Glu	Lys	Glu	Phe	Glu	Val	Thr	Thr	Asn	Tyr	Tyr			
65					70					75					80			
aaa	cac	ttg	ggg	gtt	aat	tgt	gga	acc	gaa	att	tac	tat	aaa	ctt	aaa		288	
Lys	His	Leu	Gly	Val	Asn	Cys	Gly	Thr	Glu	Ile	Tyr	Tyr	Lys	Leu	Lys			
				85					90					95				
tat	caa	aaa	caa	aaa	tgc	tat	cat	aaa	ata	aac	caa	tat	ttt	aaa	aag		336	
Tyr	Gln	Lys	Gln	Lys	Cys	Tyr	His	Lys	Ile	Asn	Gln	Tyr	Phe	Lys	Lys			
			100					105					110					
aaa	aaa	gaa	att	aaa	ttt	aac	tta	aga	gta	agt	gca	ttt	ttt	aat	aaa		384	
Lys	Lys	Glu	Ile	Lys	Phe	Asn	Leu	Arg	Val	Ser	Ala	Phe	Phe	Asn	Lys			
		115					120					125						



```

aaa cac tca aaa aaa ggg agt gta gaa tta aag gaa tgt aat aat aat 432
Lys His Ser Lys Lys Gly Ser Val Glu Leu Lys Glu Cys Asn Asn Asn
130 135 140

aat aat aat aaa gag aaa gaa aca tcc caa aaa att gaa att tta caa 480
Asn Asn Asn Lys Glu Lys Glu Thr Ser Gln Lys Ile Glu Ile Leu Gln
145 150 155 160

aca aaa gtc tat gcc aaa aaa tgt aaa ttt ttg aca aac tac tat act 528
Thr Lys Val Tyr Ala Lys Lys Cys Lys Phe Leu Thr Asn Tyr Tyr Thr
165 170 175

aaa att tta 537
Lys Ile Leu

```

```

<210> 87
<211> 179
<212> PRT
<213> Borrelia afzelii

```

```

<400> 87
Met Glu Lys Ile Glu Lys Phe Lys Asn Lys Cys Gln His Lys Leu Gln
1 5 10 15

His Lys Leu Ile Val Leu Val Ser Thr Leu Cys Tyr Ile Asn Asn Lys
20 25 30

Asn Lys Lys Tyr Ser Gln Ser Asn Ile Leu Tyr Tyr Phe Asn Glu Asn
35 40 45

Leu Lys Arg Asn Gly Gln Thr Pro Ile Lys Ile Lys Thr Leu Gln Asn
50 55 60

Tyr Leu Tyr Lys Leu Glu Lys Glu Phe Glu Val Thr Thr Asn Tyr Tyr
65 70 75 80

Lys His Leu Gly Val Asn Cys Gly Thr Glu Ile Tyr Tyr Lys Leu Lys
85 90 95

Tyr Gln Lys Gln Lys Cys Tyr His Lys Ile Asn Gln Tyr Phe Lys Lys
100 105 110

Lys Lys Glu Ile Lys Phe Asn Leu Arg Val Ser Ala Phe Phe Asn Lys
115 120 125

Lys His Ser Lys Lys Gly Ser Val Glu Leu Lys Glu Cys Asn Asn Asn
130 135 140

Asn Asn Asn Lys Glu Lys Glu Thr Ser Gln Lys Ile Glu Ile Leu Gln
145 150 155 160

Thr Lys Val Tyr Ala Lys Lys Cys Lys Phe Leu Thr Asn Tyr Tyr Thr
165 170 175

Lys Ile Leu

```

```

<210> 88
<211> 2775
<212> DNA
<213> Borrelia garinii

```

&lt;400&gt; 88

```

cggaatcaa gccacctaaa acaacttccc aaaagtttct caaaaaatat tatattcagc 60
agtaaattct ataagtcatt aattatttta tactattcaa cagtaaattc tataagtcac 120
taattattta atactattca gcagtaaatt ctataagtca ttaattattt aatactattc 180
agcagtaaat tctataagtc attaattatt taatactatt cagcagtaaa ttctataagt 240
cattaattat ttaatactat tcagcagtaa attctataag tcattaatta ttttaatacta 300
ttcagcagta aattctataa gtcattaatt caattaggta acggattctt agatgtattc 360
acctcttttg gtggattagt tgcagatgca ttgggggtta aagctgatcc aaaaaaatct 420
gatgtaaaaa cttattttga atctctagct aaaaaattag aagaaacaaa agatgggtta 480
actaagttgt ccaaaggtaa tgacgggtgat actggaaagg ctgggtgatgc tgggtgggct 540
ggtggtggcg ctagtgtctgc aggtggcgct ggtgggattg agggcgctat aacagagatt 600
agcaaattggt tagatgatat ggcaaaagct gctgcggaag ctgcaagtgc tgctactggc 660
aatgcagcaa ttggggatgt tgttaattgg aatgggtggag cagcaaaagg tgggtgatgcg 720
gagagtgtta atgggattgc taaggggata aaggggattg ttgatgctgc tgagaaggct 780
gatgcgaagg aaggggaagt ggatgtggct ggtgatgctg gtggggctgg tgggtggcgct 840
ggtgctgcag gtggcgctgg tgggattgag ggcgctataa cagagattag caaatgggta 900
gatgatattg caaaagctgc tgcggttgct gcaagtgtgc caagtgtgc tactggtaac 960
gcagcaattg gggatgttgt taatggtaac gatggagcag caaaagggtg tgatgcggcg 1020
agtgttaatg ggattgctaa ggggataaag gggattgttg atgctgctga gaaggctgat 1080
gcgaaggaa ggaagttgga tgtggctggt gatgctggtg agggtaacaa ggaatgctgg 1140
aagctgtttg tgaagaagaa tctggtgat gagggtgggt aagcaaatga tgctgggaag 1200
gctgctgctg cggttgctgc tgttagtggt gaggcagat taaaagcgat tgttgatgct 1260
gctgaggggt atgataagca gggtaagaag gctgcggat ctacaaatcc gattgaggcg 1320
gctattgggg gtgcggatgc ggtgctaat gctgagcgct ttaataagat gaagaaggat 1380
gatcagattg ctgctgctat ggttctgagg ggaatggcta aggatgggca gtttgctttg 1440
aaggatgatg ctgctgctca tgaagggact tgcaagtgc tctactggta gtacaacgat tggagatggt 1500
gctgcggaag ctgcaagtgc tgcaagtgc tctactggta gtacaacgat tggagatggt 1560
gttaagagtg gtgaggcaaa agatggtgat gcggcgagtg ttaatgggat tgctaagggg 1620
ataaagggga ttgttgatgc tgctgagaag gctgatgcga aggaaggga gttggatgtg 1680
gctggtgctg ctggtacgac taacgtgaat gttgggaagt tgtttgtgaa gaataatggt 1740
aatgaggggt gtgatgcaag tgatgctggg aaagctgctg ctgcggttgc tgctgtagt 1800
ggggagcaga tattaanaagc gattggtgat gctgctaaag atggtgataa gcaggggggt 1860
actgatgtaa aggatgctac aaatccgatt gaggcggtca ttgggggtac aaatgataat 1920
gatgctgcgg cgtttgctac tatgaagaag gatgatcaga ttgctgctgc tatggttctg 1980
aggggaatgg ctaaggatgg gcagtttgct ttgaaggatg atgctgctaa ggatgggtgat 2040
aaaacggggg ttgctgcgga tgctgaaaat ccgattgacg cggctatttg ggggtgcggat 2100
gctgatgctg cggcggttta taaggagggg atgaagaagg atgatcagat tgctgctgct 2160
atggttctga ggggaatggc taaggatggg cagtttgctt tgacgaataa tgctgctgct 2220
catgaaggga ctgttaagaa tgctgttgat atggcaaaag ctgctgcggg tgctgcaagt 2280
gctgctactg gcaatgcagc aattggggat gttgttaaga gtaatgggtg agcagcagca 2340
aaaggtgggt atgcggcgag tgttaattgg attgctaagg ggataaaggg gattgttgat 2400
gctgctgaga aggtgatgct gaaggaaggg aagttggatg tggctgggtg tgctggtgaa 2460
actaacaagg atgctgggaa gttgtttgtg aagaagaatg gtgatgatg tgggtgatgca 2520
ggtgatgctg ggaaggctgc tgctgcgggt gctgctgcta gtggggagca gatattaaaa 2580
gcgattgttg atgctgctaa agatggtgat aagacggggg ttactgatgt aaaggatgct 2640
acaaatccga ttgacgcggc tattgggggg agtgcggtat ctaatgctga ggcgtttgat 2700
aagatgaaga aggatgatca gattgctgct gctatggttc tgaggggaat ggctaaggat 2760
gggcagtttg ctttg 2775

```

&lt;210&gt; 89

&lt;211&gt; 2075

&lt;212&gt; DNA

<213> *Borrelia garinii*

&lt;400&gt; 89

```

ataaagggga ttgttgatgc tgctgagaag gctgatgcga aggaaggga gttggatgtg 60
gctggtgatg ctggtgaaac taacaaggat gctgggaagt tgtttgtgaa gaacaatgg 120
aatgaggggt gtgatgcaga tgatgctggg aaggctgctg ctgcggttgc tgctgtagt 180
ggggagcaga tattaanaagc gattgttgat gctgctaagg gtggtgataa gacgggtaag 240
aataatgtga aggatgctga aaatccgatt gaggcggtca ttgggagtag tgcggtgct 300
gatgctgcgg cgttttaataa ggaggggatg aagaaggatg atcagattgc tgctgctatg 360
gttctgaggg gaatggctaa ggatgggcag tttgctttga cgaatgatgc tgctgctcat 420
gaagggactg ttaagaatgc tgttgggagt gcaacaaata agaccgttgt tgctttggct 480

```

```

aacttggttc gaaagaccgt gcaagctggg ttgaagaagg ttggggatgt tgттаагаат 540
agtгaggcaa aagatggтga tgcggcgagt gtтаатggga ttgctaaggг gataаaggгг 600
attgtttgat ctgctgagaa ggctgatgcg aaggаaggga agttggatgt ggctggтgct 660
gctggтgaaa cтааcaaggа tgctgggaag ttgtttgtga agaаgaataa tgaggгtggt 720
gaagcaaatg atgctggгaa ggctgctgct gcgгttgctg ctgttagtgг ggagcagata 780
ttaaaagcga ttgttgatgc tgctaaggat ggtgatgata agcaggгтаа gaaggctgag 840
gatgctacaa atccgattga cgcggctatt gggggтgcag gtgcggгггc таатgctgct 900
gcggcgтттa атаатatgaa gaaggatgat cagattgctg ctgctatggт tctgaggгga 960
atggctaagg atgggcagtt tgctttgacg aataatgctc atactaatca таaggгgact 1020
gtтаагаатg ctgttgatat gacaaaagct gctgcggггг ctgcaagtgc tgcaagtgct 1080
gctactggta atgcagcaat tggggatgгt gtтаатggta atgatggagc agcaaaaggг 1140
ggtgatgcgg cgagtgtтаа tgggattgct aaggгgataa aggggattgt tgatgctgct 1200
gagaaggctg atgcgaaggа agggаagггг aatgtggггг gtgctgctgгг tgctgaggгг 1260
aacgaggctg ctgggaagct gtttgtgaag аагаатgctg gtgatcatgg тggтgaagca 1320
ggtgatgctg ggaggггггг tgctgcggгг gctgctgtта gtggggagca gatattaaaa 1380
gcgattgttg atgctgctaa ggatggгггг gataagcagg gтаагаaggc tgaggatgct 1440
gaaaatccga ttgacgcggc тattgggagт acggгггггг atgataatgc tgctgaggгг 1500
tttgctacta tgaagaaggа tgatcagatt gctgctgcta тggгггггг gggaatggгг 1560
aaggatgggc agtttgcttt gaaggatgct gctcatgata atcataaggг gactgtтааg 1620
aatgctgttg atataataaa ggctactgcg gttgctgcaa gtgctgctac тggtagtaca 1680
acgattgggg atgttgгтаа gaatggгггг gcaaaaggгг gtgaggггaa gagtgtтаат 1740
gggattgcta aggggataaa ggggattgгг gatgctgctg gaaaggгггг tgсgaaggaa 1800
gggaagгггг atgtggгггг tgctgctggг gaggгггггг aggгггггг gaagгггггг 1860
gtgtaaatta ctataggatt agaactagtg tacgatatga gtcctttggг тattttgcag 1920
ctgctaatag atttgaaata agtgaagггг ааатгггггг ttттаатggа acacatttta 1980
ttgctacaaa agagaaaggа atattatag attcactггг тttaaggггг cgtggaaaaa 2040
tatттгaaат aactтcaаag cgaatgtттa agctt 2075

```

&lt;210&gt; 90

&lt;211&gt; 552

&lt;212&gt; DNA

<213> *Borrelia garinii*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(552)

&lt;400&gt; 90

```

gaa ggg act gtt aag aat gct gtt gat atg gca aaa gct gct gcg gtt 48
Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val
  1             5             10            15

gct gca agt gct gct act ggc aat gca gca att ggg gat gtt gtt aag 96
Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys
          20             25             30

agt aat ggt gga gca gca gca aaa ggt ggt gat gcg gcg agt gtt aat 144
Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn
      35             40             45

ggg att gct aag ggg ata aag ggg att gtt gat gct gct gag aag gct 192
Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala
  50             55             60

gat gcg aag gaa ggg aag ttg gat gtg gct ggt gct gct ggt gaa act 240
Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr
  65             70             75             80

aac aag gat gct ggg aag ttg ttt gtg aag aag aat ggt gat gat ggt 288
Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly
          85             90             95

ggg gat gca ggt gat gct ggg aag gct gct gct gcg gtt gct gct gtt 336

```

Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val  
 100 105 110

agt ggg gag cag ata tta aaa gcg att gtt gat gct gct aaa gat ggt 384  
 Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
 115 120 125

gat aag acg ggg gtt act gat gta aag gat gct aca aat ccg att gac 432  
 Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp  
 130 135 140

gcg gct att ggg ggg agt gcg gat gct aat gct gag gcg ttt gat aag 480  
 Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys  
 145 150 155 160

atg aag aag gat gat cag att gct gct gct atg gtt ctg agg gga atg 528  
 Met Lys Lys Asp Asp Gln Ile Ala Ala Met Val Leu Arg Gly Met  
 165 170 175

gct aag gat ggg cag ttt gct ttg 552  
 Ala Lys Asp Gly Gln Phe Ala Leu  
 180

<210> 91  
 <211> 184  
 <212> PRT  
 <213> Borrelia garinii

<400> 91  
 Glu Gly Thr Val Lys Asn Ala Val Asp Met Ala Lys Ala Ala Val  
 1 5 10 15

Ala Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Lys  
 20 25 30

Ser Asn Gly Gly Ala Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn  
 35 40 45

Gly Ile Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala  
 50 55 60

Asp Ala Lys Glu Gly Lys Leu Asp Val Ala Gly Ala Ala Gly Glu Thr  
 65 70 75 80

Asn Lys Asp Ala Gly Lys Leu Phe Val Lys Lys Asn Gly Asp Asp Gly  
 85 90 95

Gly Asp Ala Gly Asp Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val  
 100 105 110

Ser Gly Glu Gln Ile Leu Lys Ala Ile Val Asp Ala Ala Lys Asp Gly  
 115 120 125

Asp Lys Thr Gly Val Thr Asp Val Lys Asp Ala Thr Asn Pro Ile Asp  
 130 135 140

Ala Ala Ile Gly Gly Ser Ala Asp Ala Asn Ala Glu Ala Phe Asp Lys  
 145 150 155 160

Met Lys Lys Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met  
 165 170 175

Ala Lys Asp Gly Gln Phe Ala Leu

180

<210> 92  
 <211> 420  
 <212> DNA  
 <213> *Borrelia garinii*

<220>  
 <221> CDS  
 <222> (1)..(420)

<400> 92  
 ata aag ggg att gtt gat gct gct gag aag gct gat gcg aag gaa ggg 48  
 Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly  
 1 5 10 15  
 aag ttg gat gtg gct ggt gat gct ggt gaa act aac aag gat gct ggg 96  
 Lys Leu Asp Val Ala Gly Asp Ala Glu Thr Asn Lys Asp Ala Gly  
 20 25 30  
 aag ttg ttt gtg aag aac aat ggt aat gag ggt ggt gat gca gat gat 144  
 Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly Gly Asp Ala Asp Asp  
 35 40 45  
 gct ggg aag gct gct gct gcg gtt gct gct gtt agt ggg gag cag ata 192  
 Ala Gly Lys Ala Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
 50 55 60  
 tta aaa gcg att gtt gat gct gct aag ggt ggt gat aag acg ggt aag 240  
 Leu Lys Ala Ile Val Asp Ala Ala Lys Gly Gly Asp Lys Thr Gly Lys  
 65 70 75 80  
 aat aat gtg aag gat gct gaa aat ccg att gag gcg gct att ggg agt 288  
 Asn Asn Val Lys Asp Ala Glu Asn Pro Ile Glu Ala Ala Ile Gly Ser  
 85 90 95  
 agt gcg gat gct gat gct gcg gcg ttt aat aag gag ggg atg aag aag 336  
 Ser Ala Asp Ala Asp Ala Ala Ala Phe Asn Lys Glu Gly Met Lys Lys  
 100 105 110  
 gat gat cag att gct gct gct atg gtt ctg agg gga atg gct aag gat 384  
 Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp  
 115 120 125  
 ggg cag ttt gct ttg acg aat gat gct gct gct cat 420  
 Gly Gln Phe Ala Leu Thr Asn Asp Ala Ala Ala His  
 130 135 140

<210> 93  
 <211> 140  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 93  
 Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala Lys Glu Gly  
 1 5 10 15  
 Lys Leu Asp Val Ala Gly Asp Ala Gly Glu Thr Asn Lys Asp Ala Gly  
 20 25 30  
 Lys Leu Phe Val Lys Asn Asn Gly Asn Glu Gly Gly Asp Ala Asp Asp  
 35 40 45

Ala Gly Lys Ala Ala Ala Val Ala Ala Val Ser Gly Glu Gln Ile  
 50 55 60

Leu Lys Ala Ile Val Asp Ala Ala Lys Gly Gly Asp Lys Thr Gly Lys  
 65 70 75 80

Asn Asn Val Lys Asp Ala Glu Asn Pro Ile Glu Ala Ala Ile Gly Ser  
 85 90 95

Ser Ala Asp Ala Asp Ala Ala Ala Phe Asn Lys Glu Gly Met Lys Lys  
 100 105 110

Asp Asp Gln Ile Ala Ala Ala Met Val Leu Arg Gly Met Ala Lys Asp  
 115 120 125

Gly Gln Phe Ala Leu Thr Asn Asp Ala Ala Ala His  
 130 135 140

<210> 94  
 <211> 942  
 <212> DNA  
 <213> *Borrelia garinii*

<400> 94  
 atgagaggat cgcatacaca tcaccatcac ggatccaagg ggactgttaa gaatgctgtt 60  
 gatatacaca aagctgctgc ggttgctgca agtgctgcaa gtgctgctac tggtaatgca 120  
 gcaattgggg atgttggttaa tggtaatgat ggagcagcaa aaggtggtga tgcggcgagt 180  
 gttaatggga ttgctaaggg gataaagggg attggtgatg ctgctgagaa ggctgatgctg 240  
 aaggaaggga agttgaatgt ggctggtgct gctggtgctg agggtaacga ggctgctggg 300  
 aagctgtttg tgaagaagaa tgctggtgat catggtggtg aagcaggtga tgctgggagg 360  
 gctgctgctg cgggtgctgc tgttagtggg gagcagatat taaaagcgat tgttgatgct 420  
 gctaaggatg gtggtgataa gcagggttaag aaggctgagg atgctgaaaa tccgattgac 480  
 gcggtctattg ggagtacggg tgcggatgat aatgctgctg aggcgtttgc tactatgaag 540  
 aaggatgatc agattgctgc tgctatggtt ctgaggggaa tggctaagga tgggcagttt 600  
 gctttgaagg atgctgctca tgataatcat ctgcagccaa gcttaattag ctgagcttgg 660  
 actcctgttg atagatccag taatgacctc agaactccat ctggatttgt tcagaacgct 720  
 cgggtgcccgc cgggcgtttt ttattggtga gaatccaagc tagcttggcg agattttcag 780  
 gagctaagga agctaaaatg gagaaaaaat cactggatat accaccgttg atatatccca 840  
 atggcatcgt aaagaacatt ttgaggcatt tcagtcagtt gctcaatgta cctataacca 900  
 gaccgttcag ctggatatta cggccttttt aaagaccgta ag 942

<210> 95  
 <211> 217  
 <212> PRT  
 <213> *Borrelia garinii*

<400> 95  
 Met Arg Gly Ser His His His His His His Gly Ser Lys Gly Thr Val  
 1 5 10 15

Lys Asn Ala Val Asp Met Thr Lys Ala Ala Val Ala Ala Ser Ala  
 20 25 30

Ala Ser Ala Ala Thr Gly Asn Ala Ala Ile Gly Asp Val Val Asn Gly  
 35 40 45

Asn Asp Gly Ala Ala Lys Gly Gly Asp Ala Ala Ser Val Asn Gly Ile  
 50 55 60

Ala Lys Gly Ile Lys Gly Ile Val Asp Ala Ala Glu Lys Ala Asp Ala  
 65 70 75 80

```

<400> 97
Met Arg Gly Ser His His His His His His Gly Ser Lys Ser Ala Val
  1                    5                10              15
Asp Glu Ala Ser Lys Trp Leu Glu Glu Met Ile Thr Ala Ala Gly Glu
      20                25              30
Ala Ala Thr Lys Gly Gly Thr Gly Glu Ala Ser Glu Lys Ile Gly Asp
      35                40              45

```

Val 50	Gly	Asp	Asn	Asn	His	Gly 55	Ala	Val	Ala	Asp	Ala 60	Asp	Ser	Val	Lys
Gly 65	Ile	Ala	Lys	Gly	Ile 70	Lys	Gly	Ile	Val	Asp 75	Ala	Ala	Gly	Lys	Ala 80
Phe	Gly	Lys	Asp	Gly 85	Ala	Leu	Lys	Asp	Val 90	Ala	Ala	Ala	Ala	Gly 95	Asp
Glu	Ala	Asn	Lys 100	Asp	Ala	Gly	Lys	Leu 105	Phe	Ala	Gly	Gln	Asp 110	Gly	Gly
Gly	Ala	Asp 115	Gly	Asp	Ile	Ala	Lys 120	Ala	Ala	Ala	Ala	Val 125	Thr	Ala	Val
Ser	Gly 130	Glu	Gln	Ile	Leu	Lys 135	Ala	Ile	Val	Glu	Ala 140	Ala	Gly	Asp	Lys
Ala 145	Asn	Gln	Val	Gly	Val 150	Lys	Ala	Ala	Gly	Ala 155	Ala	Thr	Asn	Pro	Ile 160
Ala	Ala	Ala	Ile	Gly 165	Thr	Asp	Asp	Asp	Asn 170	Ala	Ala	Ala	Phe	Asp 175	Lys
Asp	Glu	Met	Lys 180	Lys	Ser	Asn	Asp	Lys 185	Ile	Ala	Ala	Ala	Ile 190	Val	Leu
Arg	Gly 195	Val	Ala	Lys	Asp	Gly	Lys 200	Phe	Ala	Ala	Asn 205	Ala	Asn	Asp	Asn
Ser 210	Lys	Ala	Ser	Val	Leu	Gln 215	Pro	Ser	Leu	Ile	Ser 220				

```
<210> 98
<211> 26
<212> DNA
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

```
<400> 98
cggaattcac tcgccttact attatc
```

26

```
<210> 99
<211> 29
<212> DNA
<213> Artificial Sequence
```

<220>  
<223> Description of Artificial Sequence: Synthetic  
Primer

```
<400> 99
cgggatccga gagtgctggt gatgaggtt
```

29

```
<210> 100
<211> 35
<212> DNA
<213> Artificial Sequence
```



&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Primer

&lt;400&gt; 100

cgggatccaa gaggctgtg gatgaggcta gcaag

35

&lt;210&gt; 101

&lt;211&gt; 35

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Primer

&lt;400&gt; 101

ttctgcagca cactgcctt actattatca ttagc

35

&lt;210&gt; 102

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Primer

&lt;400&gt; 102

cgggatccgc tggtgggagt ygcaac

26

&lt;210&gt; 103

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Primer

&lt;400&gt; 103

aactgcagat tatcatgagc agcatccttc

30

&lt;210&gt; 104

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Primer

&lt;400&gt; 104

cgggatccaa ggggactgtt aagaatgctg ttg

33

&lt;210&gt; 105

&lt;211&gt; 34

&lt;212&gt; DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 105

ttctgcagat gattatcatg agcagcatcc ttca

34

<210> 106

<211> 17

<212> DNA

<213> *Borrelia burgdorferi*

<400> 106

tgagggggct attaagg

17

<210> 107

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Primer

<400> 107

ccggaattcc gg

12

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**